

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b>                  Omeprazole magnesium delayed-release tablet 20.6 mg (equivalent to 20 mg omeprazole)</p> <p><b>Use</b> • treats frequent heartburn (occurs 2 or more days a week) for 1 to 4 days for full effect</p> <p><b>Warnings</b>                  Allergy alert: Do not use if you are allergic to omeprazole. Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>• had heartburn over 3 months.</li> <li>• This may be a sign of a more serious condition.</li> <li>• heartburn with lightheadedness, sweating or dizziness</li> <li>• chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness</li> <li>• frequent chest pain</li> <li>• frequent wheezing, particularly with heartburn</li> <li>• unexplained weight loss</li> <li>• nausea or vomiting</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are taking</b></p> <ul style="list-style-type: none"> <li>• warfarin, clopidogrel, or clostrazolamide (blood-thinning medicine)</li> <li>• prescription antifungal or anti-yeast medicines</li> <li>• diltiazem (calcium channel blocker)</li> <li>• digoxin (heart medicine)</li> <li>• tacrolimus (immune system medicine)</li> <li>• prescription antiretrovirals (medicines for HIV infection)</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>• your heartburn continues or worsens</li> <li>• you need to take this product for more than 14 days</li> <li>• you need to take more than 1 course of treatment every 4 months</li> </ul> <p>If pregnant or breastfeeding, ask a health professional before use.</p>	<p><b>Drug Facts (continued)</b></p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b> • for adults 18 years of age and older</p> <ul style="list-style-type: none"> <li>• this product is to be used once a day (every 24 hours), every day for 14 days.</li> <li>• it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours</li> </ul> <p><b>14-Day Course of Treatment</b></p> <ul style="list-style-type: none"> <li>• swallow 1 tablet with a glass of water before eating in the morning</li> <li>• take every day for 14 days</li> <li>• do not take more than 1 tablet a day</li> <li>• do not use for more than 14 days unless directed by your doctor</li> <li>• swallow whole. Do not chew or crush tablets.</li> </ul> <p><b>Repeated 14-Day Courses (if needed)</b></p> <ul style="list-style-type: none"> <li>• you may repeat a 14-day course every 4 months</li> <li>• do not take for more than 14 days or more often than every 4 months unless directed by a doctor</li> <li>• children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.</li> </ul>
<p><b>Other information</b></p> <ul style="list-style-type: none"> <li>• read the directions and warnings before use</li> <li>• keep the carton. It contains important information.</li> <li>• store at 20-25°C (68-77°F) and protect from moisture</li> </ul> <p><b>Inactive ingredients</b> glyceryl monostearate, hydroxypropyl cellulose, hydroxypropyl methylcellulose, iron oxide, magnesium stearate, methacrylic acid copolymer, microcrystalline cellulose, paraffin, polyethylene glycol 6000, polyacrylate 80, polyvinylpyrrolidone, sodium alginin, fumarate, starch, sucrose, talc, titanium dioxide, triethylcitrate</p> <p><b>Questions?</b> 1-800-289-9181</p>	<p><b>Other information</b></p> <ul style="list-style-type: none"> <li>• avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables</li> <li>• Eat slowly and avoid big meals if overweight, lose weight</li> <li>• Quit smoking</li> </ul> <p><b>Tips for Managing Heartburn</b></p> <ul style="list-style-type: none"> <li>• Do not lie flat or bend over after eating</li> <li>• Do not wear tight-fitting clothing around the stomach</li> <li>• Do not eat before bedtime</li> <li>• Raise the head of your bed</li> </ul>



# Prilosec

omeprazole magnesium delayed-release tablets 20.6 mg / acid reducer

## OTC

**14 TABLETS**

*One 14-day course of treatment*

*Treats FREQUENT Heartburn!*

*Occurring 2 or More Days A Week*

NDC 37000-455-02

Dist. by Procter & Gamble, Cincinnati, OH 45202

Product of Sweden

XXXXXXX-XXX

If you are not satisfied with Prilosec OTC, simply return the UPC code from this package and original sales receipt within 60 days of purchase for a full refund. For offer details, visit [PrilosecOTC.guarantee.com](http://PrilosecOTC.guarantee.com).



0 57000 47774 7



SA 18C 2015216

Safety Feature—Do not use if tablet blister unit is open or torn.



Dist. by Procter & Gamble, Cincinnati, OH 45202  
 000 000 00



# Prilosec

omeprazole magnesium delayed-**OTC**<sup>®</sup>  
 release tablets 20.0mg / acid reducer

28 TABLETS

Two 14-day courses of treatment

Treats **FREQUENT Heartburn!**  
 Occurring 2 Or More Days A Week

80-959-4008 2304

90520506

Place store sticker here

If you are not satisfied with Prilosec OTC<sup>®</sup>, simply return the UPC Code from this package and original sales receipt within 60 days of purchase for a full refund. For offer details, visit [PrilosecOTCGuarantee.com](http://PrilosecOTCGuarantee.com).

**Drug Facts (continued)**

**Inactive ingredients**  
 glyceryl monostearate, hydroxypropyl cellulose, hypromellose, iron oxide, magnesium stearate, methacrylic acid copolymer, microcrystalline cellulose, paraffin, polyethylene glycol 6000, polyisobutene, polyvinylpyrrolidone, sodium stearyl fumarate, starch, sucrose, talc, titanium dioxide, triethyl citrate

**Questions?** 1-800-369-9181

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
  - Do not wear tight-fitting clothing around the stomach
  - Do not eat before bedtime
  - Raise the head of your bed
  - Avoid heartburn-causing foods such as rich/spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
  - Eat slowly and avoid big meals
  - If overweight, lose weight
  - Quit smoking



**Drug Facts**

**Active ingredient (in each tablet)**  
 Omeprazole magnesium delayed-release tablet 20.0 mg (equivalent to 20.0 mg omeprazole)..... Acid reducer

**Purpose**  
 Acid reducer

**Use**  
 • Treats frequent heartburn (occurs 2 or more days a week)  
 • Not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
**Always alert:** Do not use if you are allergic to omeprazole  
 Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools.  
 These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have**

- had heartburn over 3 months.
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain (particularly with heartburn)
- frequent weight loss
- nausea or vomiting
- stomach pain

**Ask a doctor or pharmacist before use if you are taking**

- warfarin, digoxin, or diclofenac (blood-thinning medicines)
- ceftriaxone, erythromycin, or anti-fungal medicines
- dexamethasone (steroid medicine)
- digoxin (heart medicine)
- baclofen (muscle spasm medicine)
- prescription antiretroviral medicines for HIV infection)

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

**Directions**

- For adults 18 years of age and older
- This product is to be used once a day (every 24 hours), every day for 14 days
- You may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- swallow whole. Do not chew or crush tablets.

**Repeat 14-Day Course (if needed)**

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor

**Other information**

- children under 18 years of age: ask a doctor; heartburn in children may sometimes be caused by a serious condition.
- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) and protect from moisture





FPO 128 code

QUIET ZONE 0 0000 00000 0

P&G  
100% GTIN  
VERSION A  
AREA GAUGE

0 0000 00000 0 QUIET ZONE





---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/

---

ANDREA LEONARD SEGAL  
08/29/2011