









*Indicated for the treatment of frequent heartburn\*  
\*occurring 2 or more days a week*

Drug Facts	Purpose
<b>Active ingredient (in each tablet)</b> Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium) Acid reducer	
<b>Use</b> • treats frequent heartburn (occurs 2 or more days a week) • not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	
<b>Warnings</b> <b>Allergy alert:</b> • Do not use if you are allergic to omeprazole. • Omeprazole may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If an allergic reaction occurs, stop use and seek medical help right away. <b>Do not use if you have:</b> • trouble or pain swallowing food, vomiting with blood, or bloody or black stools • heartburn with lightheadedness, sweating or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain These may be signs of a serious condition. See your doctor. <b>Ask a doctor before use if you have:</b> • had heartburn over 3 months. This may be a sign of a more serious condition. • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain <b>Ask a doctor or pharmacist before use if you are taking a prescription drug.</b> Acid reducers may interact with certain prescription drugs. <b>Stop use and ask a doctor if:</b> • your heartburn continues or worsens • you need to take this product for more than 14 days • you need to take more than 1 course of treatment every 4 months • you get diarrhea • you develop a rash or joint pain <b>If pregnant or breast-feeding,</b> ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> • for adults 18 years of age and older • this product is to be used once a day (every 24 hours), every day for 14 days • it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours	

See current Drug Facts NDC 37000-455-01  
Read Drug Facts label on back before trying

**FREE SAMPLE INSIDE**  
(NOT FOR SALE)



**Prilosec**  
omeprazole delayed-release tablets  
20 mg / acid reducer **OTC**

**omeprazole delayed-release tablets 20 mg / acid reducer**

**Treats FREQUENT Heartburn!**  
Occurring 2 Or More Days A Week

**2 TABLETS**

First two doses of a 14-day course of treatment  
Questions? Call 1-800-289-9181

**Experience the first couple of days of your 14-day course of treatment!**



**FREE SAMPLE OF**



**Prilosec**  
**OTC**

*It treats FREQUENT Heartburn!*

**Tips for Managing Heartburn** - Do not lie flat or bend over after eating - Do not wear tight-fitting clothing around the stomach - Do not eat before bedtime - Raise the head of your bed  
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables  
- Eat slowly and avoid big meals - If overweight, lose weight  
- Quit smoking

*Prilosec OTC is to be used once a day (every 24 hours), every day for 14 days. Read Drug Facts label before trying.*

**Drug Facts (continued)**

**14-Day Course of Treatment**  
• swallow 1 tablet with a glass of water before eating in the morning  
• take every day for 14 days  
• do not take more than 1 tablet a day  
• do not use for more than 14 days unless directed by your doctor  
• swallow whole. Do not chew or crush tablets.

**Repeated 14-Day Courses (if needed)**  
• you may repeat a 14-day course every 4 months  
• do not take for more than 14 days or more often than every 4 months unless directed by a doctor  
• children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

**Other information**  
• read the directions and warnings before use  
• keep the carton. It contains important information.  
• store at 20-25°C (68-77°F) and protect from moisture

**Inactive ingredients** glyceryl monostearate, hydroxypropyl cellulose, hypromellose, iron oxide, magnesium stearate, methacrylic acid copolymer, microcrystalline cellulose, paraffin, polyethylene glycol 6000, polysorbate 80, polyvinylpyrrolidone, sodium stearyl fumarate, starch, sucrose, talc, titanium dioxide, triethyl citrate

**Questions? 1-800-289-9181**

Dist. by Procter & Gamble, Cincinnati, OH 45202  
Product of Sweden

**P&G**  
www.pg.com  
Patents: www.pg.com/patents

**TREATS FREQUENT HEARTBURN**

90341736

**Drug Facts Specifications**

- The title Drug Facts 11 pt left justified, 11 pt leading
- Drug Facts (continued) 9 pt, bold, italic (continued in regular type, not bold or italic)
- Headings 9 pt, bold, italic, left justified, 10 pt leading
- Subheadings 6 pt, bold, left justified, 7-9 pt leading
- Text 7 pt, left justified, 0.5-1 pt leading
- 34 characters per inch
- Dark purple font printed on white background
- Bullets are solid circles, 9 pt
- Barlines separating each of the headings, entire width of the Drug Facts box, 1 pt
- Hairlines following titles and preceding subheadings, within 2 spaces on either side of the Drug Facts box, 0.5 pt













-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

JODY E GREEN  
10/04/2022 12:03:34 PM