

NDA 21-087/S-033

NDA 21-246/S-021

Page 23



Patient Information

TAMIFLU®

(oseltamivir phosphate)

R_x ONLY

This leaflet contains important information about TAMIFLU (TAM-ih-flew). Read it well before you begin treatment. This information does not take the place of talking with your healthcare professional about your medical condition or your treatment. This leaflet does not list all the benefits and risks of TAMIFLU. If you have any questions about TAMIFLU, ask your healthcare professional. Only your healthcare professional can determine if TAMIFLU is right for you.

What is TAMIFLU?

TAMIFLU attacks the influenza virus and stops it from spreading inside your body. TAMIFLU treats flu at its source, by attacking the virus that causes the flu, rather than simply masking symptoms.

TAMIFLU is for treating adults and children age 1 and older with the flu whose flu symptoms started within the last day or two. TAMIFLU can also reduce the chance of getting the flu in people age 1 and older who have a higher chance of getting the flu because they spend time with someone who has the flu. TAMIFLU can also reduce the chance of getting the flu if there is a flu outbreak in the community.

What is “Flu”?

“The flu” is an infection caused by the influenza virus. Flu symptoms include fever (usually 100°F to 103°F in adults, and sometimes higher in children) and problems such as cough, sore throat, runny or stuffy nose, headaches, muscle aches, fever, and extreme tiredness. Many people use the term “flu” to mean any combination of these symptoms, such as the common cold, but true influenza infection is often worse and may last longer than a cold.

Flu outbreaks happen about once a year, usually in the winter, when the influenza virus spreads widely in the community. Outside of those outbreaks, only a very tiny number of respiratory infections are caused by the influenza virus.

Should I get a flu shot?

TAMIFLU is not a substitute for a flu vaccination. You should continue to get a flu vaccination every year, according to your healthcare professional’s advice.

Who should not take TAMIFLU?

Do not take TAMIFLU if you are allergic to the main ingredient, oseltamivir phosphate, or to any other ingredients of TAMIFLU. Before starting treatment, make sure your healthcare professional knows if you take any other medicines, or are pregnant, planning to become pregnant, or breastfeeding. TAMIFLU is normally not recommended for use during pregnancy or nursing, as the effects on the

NDA 21-087/S-033
NDA 21-246/S-021
Page 24

unborn child or nursing infant are unknown. TAMIFLU is not recommended for use in children younger than 1 year of age.

Tell your healthcare professional if you have any type of kidney disease, heart disease, respiratory disease, or any serious health condition.

How should I take TAMIFLU?

It is important that you begin your treatment with TAMIFLU as soon as possible from the first appearance of your flu symptoms or soon after you are exposed to the flu. If you feel worse or develop new symptoms during treatment with TAMIFLU, or if your flu symptoms do not start to get better, you should contact your healthcare professional.

If you have the flu: Take TAMIFLU twice a day for 5 days, once in the morning and once in the evening. You should complete the entire treatment of 10 doses (capsules or liquid), even if you feel better.

To prevent the flu: If someone in your home has the flu, take TAMIFLU once a day for 10 days or for as long as prescribed. You can take TAMIFLU for up to 6 weeks if you are exposed to the flu because of an outbreak in your community. Follow your healthcare professional's advice on how long to take TAMIFLU.

TAMIFLU has not been studied in children 1 to 12 years of age for preventing flu during an outbreak in your community or for use for more than 10 days.

You can take TAMIFLU with food or without food. There is less chance of stomach upset if you take it with a light snack, milk, or a meal.

If you are taking TAMIFLU liquid, your pharmacist will give you a dosing dispenser marked with three possible doses. Follow your healthcare professional's instructions on which dose to take or how to combine them for the proper dose for you. In order to be sure you receive the proper dose, it is important that you use the dispenser provided. Review the instructions below on how to use the dispenser and ask your pharmacist if you have any questions. If you lose or damage the dispenser and cannot use it, contact your healthcare professional or pharmacist for advice on the proper dose.

If you forget to take your medicine, take the missed dose as soon as you remember, except if it is 2 hours or less before your next dose. Then continue to take TAMIFLU at the usual times. Do not take 2 doses at a time to make up for a missed dose. If you miss several doses, tell your healthcare professional and follow the advice given to you.

What are the possible side effects of TAMIFLU?

The most common side effects of TAMIFLU are nausea and vomiting. These are usually mild to moderate. They usually happen in the first 2 days of treatment. Taking TAMIFLU with food may reduce the chance of getting these side effects.

If you develop an allergic reaction or severe rash, stop taking TAMIFLU and contact your healthcare professional.

People with the flu, particularly children, may be at an increased risk of self-injury and confusion shortly after taking TAMIFLU and should be closely monitored for signs of unusual behavior. A

NDA 21-087/S-033
NDA 21-246/S-021
Page 25

healthcare professional should be contacted immediately if the patient taking TAMIFLU shows any signs of unusual behavior.

Before taking TAMIFLU, please let your healthcare provider know if you have received nasally administered influenza virus vaccine during the past two weeks.

If you notice any side effects not mentioned in this leaflet, or if you have any concerns about the side effects you get, tell your healthcare professional.

How and where should I store TAMIFLU?

TAMIFLU capsules should be stored at room temperature below 77°F (25°C) and kept in a dry place. Keep this medication out of reach of children.

TAMIFLU suspension should be stored under refrigeration at 36° to 46°F (2° to 8°C). Do not freeze.

General advice about prescription medicines:

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not use TAMIFLU for a condition for which it was not prescribed. Do not give TAMIFLU to other people, even if they have the same symptoms you have. It may not be right for them.

This leaflet summarizes the most important information about TAMIFLU. If you would like more information, talk with your healthcare professional. You can ask your pharmacist or healthcare professional for information about TAMIFLU that is written for health professionals.


DOSING INSTRUCTIONS FOR PATIENTS:

Please follow instructions carefully to ensure proper dosing of the oral suspension.



- Shake closed bottle well for about 5 seconds before each use.
- Remove child-resistant cap.
- Before inserting the tip of the oral dispenser into bottle adapter, push the plunger completely down toward the tip of the oral dispenser. Insert tip firmly into opening of the bottle adapter.
- Turn the entire unit (bottle and oral dispenser) upside down.
- Pull the plunger out slowly until the desired amount of medication is withdrawn into the oral dispenser (see figure). The 75 mg dose is obtained by filling the dispenser twice, once to the 30 mg graduation, and a second fill to the 45 mg graduation.

NDA 21-087/S-033
NDA 21-246/S-021
Page 26



- Turn the entire unit right side up and remove the oral dispenser slowly from the bottle.
- Dispense directly into mouth. Do not mix with any liquid prior to dispensing.
- Close bottle with child-resistant cap after each use.
- Disassemble oral dispenser, rinse under running tap water and air dry prior to next use.

Distributed by:



Pharmaceuticals

Roche Laboratories Inc.
340 Kingsland Street
Nutley, New Jersey 07110-1199

Licensors:

Gilead Sciences, Inc.
Foster City, California 94404

Revised: Month Year

Copyright © 1999-200x by Roche Laboratories Inc. All rights reserved.