

Contents: 1 mg treprostinil per mL.
Each mL also contains 0.5 mg sodium citrate dihydrate, 5.3 mg sodium chloride, 3 mg metacresol, 0.24 mg sodium hydroxide, and Water for Injection. Hydrochloric acid and sodium hydroxide may have been added to adjust pH.
Prior to intravenous infusion, Remodulin® may be diluted.
See Package Insert for Dosage and Administration.
United Therapeutics Corp., RTP, NC USA 27709

REMODULIN®
(treprostinil) Injection
20 mg/20 mL
(1 mg/mL)


For Subcutaneous and intravenous infusion

20 mL multidose vial

United Therapeutics

NDC 66302-101-01
Store at 25°C (77°F) with excursions permitted to 2-30°C (36-86°F).
Keep out of reach of children.

Rx Only	100350	Rev. 01/2023
LOT	XXXXXXXX	
EXP	MM-YYYY	



(b) (4)

Contents: 2.5 mg treprostinil per mL.
Each mL also contains 0.3 mg sodium citrate dihydrate, 5.3 mg sodium chloride, 3 mg metacresol, 0.32 mg sodium hydroxide, and Water for Injection. Hydrochloric acid and sodium hydroxide may have been added to adjust pH.
Prior to intravenous infusion, Remodulin™ may be diluted.
See Package Insert for Dosage and Administration.
United Therapeutics Corp., RTP, NC USA 27709

REMODULIN™
(treprostinil) Injection
50 mg/20 mL
(2.5 mg/mL)


For Subcutaneous and intravenous infusion

20 mL multidose vial
United Therapeutics

NDC 66302-102-01
Store at 25°C (77°F) with excursions permitted to 2-30°C (36-86°F).
Keep out of reach of children.

Rx Only 100351 Rev. 01/2023

LOT XXXXXXXX
EXP MM-YYYY



(b) (4)


Contents: 5 mg treprostinil per mL.
Each mL also contains 6.5 mg sodium citrate dihydrate, 5.3 mg sodium chloride, 3 mg metacresol, 0.62 mg sodium hydroxide, and Water for Injection. Hydrochloric acid and sodium hydroxide may have been added to adjust pH.
Prior to intravenous infusion, Remodulin® may be diluted.
See Package Insert for Dosage and Administration.
United Therapeutics Corp., RTP, NC USA 27709



100 mg/20 mL
(5 mg/mL)

For Subcutaneous and intravenous infusion

20 mL multidose vial



NDC 66302-105-01
Store at 25°C (77°F) with excursions permitted to 2-30°C (36-86°F).
Keep out of reach of children.

Rx Only	100352	Rev. 01/2023
LOT	XXXXXXXX	
EXP	MM-YYYY	

(b) (4)

Contents: 10 mg treprostinil per mL.
Each mL also contains 6.5 mg sodium citrate dihydrate, 4 mg sodium chloride, 3 mg metacresol, 1.2 mg sodium hydroxide, and Water for Injection. Hydrochloric acid and sodium hydroxide may have been added to adjust pH.
Prior to intravenous infusion, Remodu tr[®] may be diluted.
See Package Insert for Dosage and Administration.
United Therapeutics Corp., RTP, NC USA 27709

REMODULIN[®]
(treprostinil) Injection
200 mg/20 mL
(10 mg/mL)

For Subcutaneous and intravenous infusion


20 mL multidose vial

United Therapeutics

NDC 66302-110-01
Store at 25°C (77°F) with excursions permitted to 2-30°C (36-86°F).
Keep out of reach of children.

Rx Only 100353 Rev. 01/2023

LOT XXXXXXXX
EXP MM-YYYY



(b) (4)

Contents: 20 mg treprostinil per mL.
Each mL also contains 6.3 mg sodium citrate dihydrate, 4 mg sodium chloride, 3 mg metacresol, 2.25 mg sodium hydroxide, and Water for Injection. Hydrochloric acid and sodium hydroxide may have been added to adjust pH.
Prior to intravenous infusion, Remodu *in*® may be diluted.
See Package Insert for Dosage and Administration.
United Therapeutics Corp., RTP, NC USA 27709

REMODULIN[®]
(treprostinil) Injection

400 mg/20 mL
(20 mg/mL)

For Subcutaneous
and Intravenous Infusion


20 mL multidose vial

**United
Therapeutics**


NDC 65302-120-01

Store at 25°C (77°F) with excursions permitted to 2-30°C (36-86°F).
Keep out of reach of children.


Rx Only	101749	Rev. 01/2023
LOT	XXXXXXXX	
EXP	MM-YYYY	




(b) (4)




(b) (4)







(b) (4)




(b) (4)





(b) (4)



(b) (4)

