



Actual Size 100%
FDA59282-9

COVERED	REMOVED	ACTUAL IN FILE
Drug Facts Title	4.8 pt	6.5 pt
Drug Facts Content	8 pt	8 pt
Warnings	10 pt @ 14 pt	5 pt
Directions	10 pt	6 pt
Other Text	10 pt	5 pt
Bottom	10 pt	5 pt
Quantity	10 pt	10 pt
Net Wt	10 pt	10 pt
Box and Package	10 pt @ 10 pt	10 pt
Character for Box	10 pt @ 10 pt	10 pt

Do not use if seal on tube is broken or not visible. See crimp for lot number and expiration date. Store between 20° to 25°C (68° to 77°F). Questions? 1-866-060-0266. Bayer and Lotrimin Ultra are registered trademarks of Bayer. © 2021 Bayer. Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981. Product of Japan. **Bayer**

LOTTRIMIN ULTRA®
 butenafine hydrochloride cream 1% ANTIFUNGAL Net Wt 1.2g (0.42 oz)

Active ingredient Butenafine hydrochloride 1% **Purpose** Antifungal

Uses ■ cures most ringworm ■ cures most jock itch ■ cures most athlete's foot between the toes ■ relieves itching, burning, cracking, and scaling which accompany these conditions

Warnings For external use only

Do not use ■ on nails or scalp ■ in or near the mouth or the eyes ■ for vaginal yeast infections
When using this product do not get into the eyes. If eye contact occurs, rinse thoroughly with water.
Stop use and ask a doctor if too much irritation occurs or irritation gets worse.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ adults and children 12 years and older: ■ use the tip of the cap to break the seal and open the tube ■ wash the affected skin with soap and water and dry completely before applying ■ for ringworm and jock itch: apply once a day to affected skin for 2 weeks or as directed by a doctor ■ for athlete's foot between the toes: apply to affected skin between and around the toes twice a day for 1 week (morning and night), or once a day for 4 weeks, or as directed by a doctor. Wear well-fitting, ventilated shoes. Change shoes and socks at least once daily. ■ wash hands after each use ■ children under 12 years: ask a doctor



Actual Size 100%

FDA59303-4

ICL (Updated 2/27/19)		
CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt*	N/A
Drug Facts Continued	7 pt*	N/A
Headings	7 pt-bold italic*	7 pt
Subheadings	6 pt*	6 pt
Body Text	6 pt*	6 pt
Bullets	5 pt	5 pt
Hairlines	0.5 pt rule	0.5 pt
Barlines ¹	> 0.5 pt rule	N/A
Characters per inch rule is not applicable to ICL. *ICL minimum font size is 4.5 pt. < 0.5 pt leading may be used as long as ascenders and descenders do not touch. FDA recommends size 2.5 pt type		

Do not use if seal on tube is broken or not visible. See crimp for lot number and expiration date.
Store between 20° to 25°C (68° to 77°F). Questions? 1-866-360-3266
Bayer and Lotrimin Ultra are registered trademarks of Bayer.
© 2021 Bayer Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981 Product of Japan

Bayer



Active ingredient Butenafine hydrochloride 1% **Purpose** Antifungal

Uses ■ cures most ringworm ■ cures most jock itch ■ cures most athlete's foot between the toes
■ relieves itching, burning, cracking, and scaling which accompany these conditions

Warnings For external use only

Do not use ■ on nails or scalp ■ in or near the mouth or the eyes ■ for vaginal yeast infections

When using this product do not get into the eyes. If eye contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if too much irritation occurs or irritation gets worse

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
■ adults and children 12 years and older: ■ use the tip of the cap to break the seal and open the tube
■ wash the affected skin with soap and water and dry completely before applying ■ for ringworm and jock
itch: apply once a day to affected skin for 2 weeks or as directed by a doctor ■ for athlete's foot between
the toes: apply to affected skin between and around the toes twice a day for 1 week (morning and night), or
once a day for 4 weeks, or as directed by a doctor. Wear well-fitting, ventilated shoes. Change shoes and
socks at least once daily. ■ wash hands after each use ■ children under 12 years: ask a doctor



Actual Size 100%

FDA59304-2

ICL (updated 2/27/19)		
CATEGORY	Bayer Standard	ACTUAL IN FILE
Drug Facts Title	>7 pt ¹	N/A
Drug Facts Continued	7 pt ²	N/A
Headings	7 pt- bold italic ³	7 pt
Subheadings	6 pt ²	6 pt
Body Text	6 pt ²	6 pt
Bullets	5 pt	5 pt
Headlines ¹	0.5 pt rule	0.5 pt
Barlines ¹	> 0.5 pt rule	N/A

Characters per inch rule is not applicable to ICL.
¹ICL minimum font size is 4.5 pt.
²< 0.5 pt leading may be used as long as ascenders and descenders do not touch.
³ FDA recommends size 2.5 pt type

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
01/02/2024 02:14:48 PM