

**MONISTAT<sup>®</sup>**  
EXTERNAL VULVAR CREAM  
Miconazole Nitrate Cream (2%)  
**EXTERNAL USE ONLY**

Relieves External Vulvar Itching & Irritation  
Due to a Vaginal Yeast Infection

Net Wt. 0.53 oz. (15 grams)

For external vulvar use only. Do not take by mouth or use in eyes.




Keep out of reach of children. Stop use and ask a doctor if symptoms last more than 7 days. Before using, read the enclosed leaflet for complete Directions and Warnings. Directions: To Open: Use cap to puncture seal. Squeeze a small amount of cream onto your fingertip. Apply to skin outside the vagina (vulva) that itches and is irritated. Use 2 times daily for up to 7 days, as needed.

Tamper-Evident Feature: The tube opening should be sealed. Do not use if foil seal has been punctured.

Active Ingredient: miconazole nitrate 2%.

See end of tube for lot number and expiration date.  
Store at 20°-25°C (68°-77°F).

© 2020 Distributed by INSIGHT Pharmaceuticals, LLC, Tarrytown, NY 10591, a Prestige Consumer Healthcare company. MNU50XX501



7-DAY  
TREATMENT  
COMBINATION PACK

50% MORE  
EXTERNAL CREAM  
FREE

Cream

Cream with  
7 Disposable Applicators

plus External Cream  
for Itch Relief

\*LIMITED TIME OFFER

# MONISTAT<sup>®</sup> 7

Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

COMBINATION PACK

TAMPER-EVIDENT UNIT  
DO NOT USE IF SEAL OVER TUBE  
HAS BEEN PUNCTURED.

# MONISTAT<sup>®</sup> 7

Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

COMBINATION PACK

- ✓ CURES most vaginal YEAST INFECTIONS
- ✓ Relieves associated external itching and irritation

Net Wt. 1.59 oz (45 g) Tube + 0.53 oz (15 g) Tube + 7 Disposable Applicators

Exp  
Lot

MONISTAT<sup>®</sup> 7

Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

COMBINATION PACK



Dist. by Insight Pharmaceuticals LLC, Tarrytown, NY 10591  
A Prestige Consumer Healthcare company  
©2019 Trade dress is owned by Insight Pharmaceuticals LLC.  
All rights reserved. MNUS001108

If you have any  
questions or  
comments, please call  
1-877-MONISTAT  
(1-877-666-4782)

**Drug Facts (continued)**

**Stop use and ask a doctor if:**

- symptoms do not get better in 3 days
- symptoms last more than 7 days
- you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- before using this product read the enclosed consumer information leaflet for complete directions and information
- adults and children 12 years of age and over:
  - applicator: insert 1 applicatorful into the vagina at bedtime for 7 nights in a row. Throw applicator away after use.
  - external cream: squeeze a small amount of cream onto your fingertip. Apply the cream onto the itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days, as needed.
- children under 12 years of age: ask a doctor

**Other information**

- do not use if seal over tube opening has been punctured
- do not purchase if carton is open
- store at 20°-25°C (68°-77°F)
- keep this carton for future reference. It contains important information

**Inactive ingredients** benzonic acid, cetyl alcohol, isopropyl myristate, polysorbate 80, potassium hydroxide, propylene glycol, purified water, stearyl alcohol

**Questions?** If you have any questions or comments, please call 1-877-666-4782

**Drug Facts**

**Active ingredients**

Miconazole nitrate 2% (vaginal antifungal)  
Miconazole nitrate 2% (external cream)

**Uses**

- treats vaginal yeast infections
- relieves external itching and irritation due to a vaginal yeast infection

**Warnings**

**For vaginal use only**

**Do not use if you have never had a vaginal yeast infection diagnosed by a doctor.**

**Ask a doctor before use if you have**

- vaginal itching and discomfort for the first time
- lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.
- vaginal yeast infections often (such as once a month or in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes, or a weakened immune system.
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS

**Ask a doctor or pharmacist before use if you are taking the prescription blood thinning medicine warfarin, because bleeding or bruising may occur.**

**When using this product**

- do not use tampons, douches, spermicides or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).
- do not have vaginal intercourse
- mild increase in vaginal burning, itching or irritation may occur if you do not get complete relief ask a doctor before using another product

MONISTAT<sup>®</sup> 7  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal  
COMBINATION PACK

**3-DAY TREATMENT COMBINATION PACK**

**Ovule®** + **50% MORE EXTERNAL CREAM FREE\***

3 Ovule® Inserts + Plus External Cream for Itch Relief

\*LIMITED TIME OFFER

# MONISTAT® 3

COMBINATION PACK

Miconazole Nitrate Vaginal Insert (200 mg) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

**TAMPER-EVIDENT UNIT DO NOT USE IF PRINTED SEALED POUCH AND SEAL OVER TUBE ARE TORN, OPENED OR INCOMPLETELY SEALED**

# MONISTAT® 3

COMBINATION PACK

Miconazole Nitrate Vaginal Insert (200 mg) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

✔ CURES most vaginal YEAST INFECTIONS

✔ Relieves associated external itching and irritation

3 OVULE® Inserts Net Wt. 2.5 g each + 0.53 oz (15 g) Tube



3 63736 44860 8

Dist. by Insight Pharmaceuticals LLC, Tarrytown, NY 10591  
A Prestige Consumer Healthcare company  
©2019 Trade dress is owned by Insight Pharmaceuticals LLC.  
All rights reserved. MNUSO36103

If you have any questions or comments, please call  
**1-877-MONISTAT**  
(1-877-666-4782)



Exp  
Lot

**MONISTAT® 3**  
COMBINATION PACK  
Miconazole Nitrate Vaginal Insert (200 mg) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

**Drug Facts**

**Active ingredients**  
Miconazole nitrate (200 mg in each vaginal insert). Vaginal antifungal.

**Purpose**  
Symptoms do not get better in 3 days  
Symptoms last more than 7 days  
You get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge

**Directions**  
Before using this product read the enclosed consumer information leaflet for complete directions and information  
Adults and children 12 years of age and over:  
vaginal insert: with a disposable applicator, place the insert into the vagina at bedtime. Throw applicator away after use.  
external cream: squeeze a small amount of cream onto your fingertip. Apply the cream onto the itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days, as needed.  
children under 12 years of age: ask a doctor.

**Other information**  
Do not use if printed sealed pouches containing vaginal inserts are torn, open or incompletely sealed  
Do not use if seal over tube opening has been punctured  
Do not purchase if carton is open  
Keep this carton for future reference. It contains important information

**Inactive ingredients**  
vaginal insert: gelatin, glycerin, hydrogenated vegetable oil base, titanium dioxide  
external cream: benzoic acid, cetyl alcohol, isopropyl myristate, polyorbate 60, potassium hydroxide, propylene glycol, purified water, stearyl alcohol

**Questions?** If you have any questions or comments, please call 1-877-666-4782

**Warnings**  
Do not use if you have never had a vaginal yeast infection diagnosed by a doctor.  
Ask a doctor before use if you have vaginal itching and discomfort for the first time  
lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.  
vaginal yeast infections often (such as once a month or 3 in 6 months), you could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes, or a weakened immune system.  
been exposed to the human immunodeficiency virus (HIV) that causes AIDS  
Ask a doctor or pharmacist before use if you are taking the prescription blood thinning medicine warfarin, because bleeding or bruising may occur.  
When using this product  
do not use tampons, douches, spermicides or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).  
do not have vaginal intercourse  
do not increase in vaginal burning, itching or irritation may occur  
if you do not get complete relief ask a doctor before using another product

**Uses**  
relieves vaginal yeast infections  
relieves external itching and irritation due to a vaginal yeast infection

**3-DAY  
TREATMENT  
COMBINATION PACK**

**50% MORE  
EXTERNAL CREAM  
FREE\***

+

Vaginal Suppositories  
with 3 Applicators

plus External Cream  
for Itch Relief

\*LIMITED TIME OFFER

# MONISTAT<sup>®</sup> 3

COMBINATION PACK

Miconazole Nitrate Vaginal Suppositories (200 mg) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

**TAMPER-EVIDENT UNIT**  
DO NOT USE IF  
PRINTED SUPPOSITORY  
BLISTERS OR SEAL  
OVER TUBE ARE TORN,  
OPENED OR  
INCOMPLETELY  
SEALED

# MONISTAT<sup>®</sup> 3

COMBINATION PACK

Miconazole Nitrate Vaginal Suppositories (200 mg) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

✓ CURES most vaginal YEAST INFECTIONS

✓ Relieves associated external itching and irritation

**3 Suppositories Net Wt. 2.5 g each + 0.53 oz (15 g) Tube**

3 63736 44305 4

Dist. by Insight Pharmaceuticals LLC,  
Tarrytown, NY 10591  
A Prestige Consumer Healthcare company  
©2019 Trade dress is owned by  
Insight Pharmaceuticals LLC.  
All rights reserved. MNU5028103

If you have any  
questions or  
comments, please call  
**1-877-MONISTAT**  
(1-877-666-4782)

Exp  
Lot

**Drug Facts** (continued)

**Stop use and ask a doctor if:**

- symptoms do not get better in 3 days
- symptoms last more than 7 days
- you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- before using this product read the enclosed consumer information leaflet for complete directions and information
- adults and children 12 years of age and over:
  - suppositories: insert 1 suppository into the vagina at bedtime for three nights in a row. Throw applicator away after use.
  - external cream: squeeze a small amount of cream onto your fingertip. Apply the cream onto the itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days, as needed.
- children under 12 years of age: ask a doctor.

**Other Information**

- do not use if printed suppository blisters are torn, open or incompletely sealed
- do not use if seal over tube opening has been punctured
- do not purchase if carton is open
- do not use if carton is open
- store at 20°-25°C (68°-77°F)
- keep this carton for future reference. It contains important information

**Inactive ingredients**

suppository: hydrogenated vegetable oil base  
external cream: benzoin acid, cetyl alcohol, isopropyl myristate, polysorbate 60, potassium hydroxide, propylene glycol, purified water, stearyl alcohol

**Questions?** If you have any questions or comments, please call 1-877-666-4782

**Drug Facts**

**Active ingredients**

Miconazole nitrate 2% (external cream).....Vaginal antifungal  
Miconazole nitrate (200 mg in each suppository).....Vaginal antifungal

**Uses**

- treats vaginal yeast infections
- relieves external itching and irritation due to a vaginal yeast infection

**Warnings**

For vaginal use only

Do not use if you have never had a vaginal yeast infection diagnosed by a doctor.

**Ask a doctor before use if you have**

- vaginal itching and discomfort for the first time
- lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.
- vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes, or a weakened immune system.
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS

**Ask a doctor or pharmacist before use if you are taking the** prescription blood thinning medicine warfarin, because bleeding or bruising may occur.

**When using this product**

- do not use tampons, douches, spermicides or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).
- do not have vaginal intercourse
- mild increase in vaginal burning, itching or irritation may occur if you do not get complete relief ask a doctor before using another product

**Drug Facts**

**Purpose**

Miconazole nitrate (200 mg in each suppository).....Vaginal antifungal  
Miconazole nitrate 2% (external cream).....Vaginal antifungal

# MONISTAT<sup>®</sup> 3

COMBINATION PACK

Miconazole Nitrate Vaginal Suppositories (200 mg) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

3-DAY  
TREATMENT  
COMBINATION PACK

**50% MORE  
EXTERNAL CREAM  
FREE!**

**Prefilled Cream  
Applicator**

+

plus External Cream  
for Itch Relief

3 Ready-to-Use  
Prefilled Applicators

\*LIMITED TIME OFFER

**MONISTAT<sup>®</sup> 3**  
COMBINATION PACK  
Miconazole Nitrate Vaginal Cream (4%) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

TAMPER-EVIDENT  
UNIT DO NOT USE IF  
PRINTED APPLICATOR  
WRAPPER OR SEAL  
OVER TUBE ARE  
TORN, OPENED OR  
INCOMPLETELY  
SEALED

Exp  
Lot

**MONISTAT<sup>®</sup> 3**  
COMBINATION PACK  
Miconazole Nitrate Vaginal Cream (4%) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

**MONISTAT<sup>®</sup> 3**  
COMBINATION PACK  
Miconazole Nitrate Vaginal Cream (4%) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

✓ CURES most vaginal YEAST INFECTIONS  
✓ Relieves associated external itching and irritation

3 Prefilled Applicators Net Wt. 0.18 oz (5 g) each + 0.53 oz (15 g) Tube



3 63736 44300 9

Dist. by Insight Pharmaceuticals LLC, Tarrytown, NY 10591  
A Prestige Consumer Healthcare company  
©2019 Trade dress is owned by Insight Pharmaceuticals LLC.  
All rights reserved. MNUSO13106

If you have any  
questions or  
comments, please call  
**1-877-MONISTAT**  
(1-877-666-4782)

**Drug Facts**

**Active ingredient** Miconazole nitrate 4% Vaginal antifungal  
Miconazole nitrate 2% (external cream) Vaginal antifungal

**Uses**  
■ treats vaginal yeast infections  
■ relieves external itching and irritation due to a vaginal yeast infection

**Warnings**  
For vaginal use only  
Do not use if you have never had a vaginal yeast infection diagnosed by a doctor.  
Ask a doctor before use if you have  
■ vaginal itching and discomfort for the first time  
■ lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.  
■ vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes, or a weakened immune system.  
■ been exposed to the human immunodeficiency virus (HIV) that causes AIDS  
■ Ask a doctor or pharmacist before use if you are taking the prescription blood thinning medicine warfarin, because bleeding or bruising may occur.  
■ do not use tampons, douches, spermicides or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).  
■ do not have vaginal intercourse  
■ mild increase in vaginal burning, itching or irritation may occur if you do not get complete relief ask a doctor before using another product

**When using this product**  
■ do not use tampons, douches, spermicides or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).  
■ do not have vaginal intercourse  
■ mild increase in vaginal burning, itching or irritation may occur if you do not get complete relief ask a doctor before using another product

**Other Information**  
■ do not use if printed applicator wrapper is torn, open or incompletely sealed  
■ do not use if seal over tube opening has been punctured  
■ do not purchase if carton is open  
■ store at 20°-25°C (68°-77°F)  
■ Keep this carton for future reference. It contains important information

**Inactive ingredients** benzoic acid, cetyl alcohol, isopropyl myristate, poly sorbate 80, potassium hydroxide, propylene glycol, purified water, stearyl alcohol

**Questions? If you have any questions or comments,** please call 1-877-666-4782

**Drug Facts (continued)**

**Purpose**  
Vaginal antifungal

**Stop use and ask a doctor if:**  
■ symptoms do not get better in 3 days  
■ symptoms last more than 7 days  
■ you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge

**Directions**  
before using this product read the enclosed consumer information leaflet for complete directions and information  
■ adults and children 12 years of age and over:  
■ applicators: insert 1 applicator into the vagina at bedtime for 3 nights in a row. Throw applicator away after use.  
■ external cream: squeeze a small amount of cream onto your fingertip. Apply the cream onto the itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days, as needed.  
■ children under 12 years of age: ask a doctor

**Directions**  
before using this product read the enclosed consumer information leaflet for complete directions and information  
■ adults and children 12 years of age and over:  
■ applicators: insert 1 applicator into the vagina at bedtime for 3 nights in a row. Throw applicator away after use.  
■ external cream: squeeze a small amount of cream onto your fingertip. Apply the cream onto the itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days, as needed.  
■ children under 12 years of age: ask a doctor

**Other Information**  
■ do not use if printed applicator wrapper is torn, open or incompletely sealed  
■ do not use if seal over tube opening has been punctured  
■ do not purchase if carton is open  
■ store at 20°-25°C (68°-77°F)  
■ Keep this carton for future reference. It contains important information

**Inactive ingredients** benzoic acid, cetyl alcohol, isopropyl myristate, poly sorbate 80, potassium hydroxide, propylene glycol, purified water, stearyl alcohol

**Questions? If you have any questions or comments,** please call 1-877-666-4782



1-DAY  
TREATMENT  
COMBINATION PACK

50%  
MORE  
EXTERNAL CREAM  
FREE

Day or Night™ OVULE®  
Stays in Place

Ovule® Insert

plus External Cream  
for Itch Relief

\*LIMITED TIME OFFER

**MONISTAT 1**  
COMBINATION PACK  
Miconazole Nitrate Vaginal Insert (1200 mg) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

TAMPER-EVIDENT  
UNIT DO NOT USE  
IF PRINTED SEALED  
POUCH AND SEAL  
OVER TUBE ARE  
TORN, OPENED OR  
INCOMPLETELY  
SEALED

CRN-0904

**MONISTAT 1**  
COMBINATION PACK  
Miconazole Nitrate Vaginal Insert (1200 mg) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

MAXIMUM  
STRENGTH  
OVULE®

- ✓ CURES most vaginal YEAST INFECTIONS
- ✓ Relieves associated external itching and irritation

1 OVULE® Insert Net Wt. 2.6 g + 0.53 oz (15 g) Tube



Dist. by Insight Pharmaceuticals LLC, Tarrytown, NY 10591  
A Prestige Consumer Healthcare company  
©2019 Trade dress is owned by Insight Pharmaceuticals LLC.  
All rights reserved. MNUS019104

If you have any  
questions or  
comments, please call  
1-877-MONISTAT  
(1-877-666-4782)

NON-VARNISH AREA



LOT  
EXP

**MONISTAT 1**  
COMBINATION PACK  
Miconazole Nitrate Vaginal Insert (1200 mg) and  
Miconazole Nitrate Vaginal Cream (2%) — Vaginal Antifungal

**Drug Facts**

**Active Ingredients**  
Miconazole nitrate 1200 mg (in vaginal insert)..... Vaginal antifungal  
Miconazole nitrate 2% (external cream)..... Vaginal antifungal

**Purpose**  
or foul-smelling vaginal discharge  
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**  
before using this product read the enclosed consumer information leaflet for complete directions and information  
adults and children 12 years of age and over:  
vaginal insert: with the applicator, place the vaginal insert into the vagina. Throw away applicator.  
external cream: squeeze a small amount of cream onto your fingertip. Apply the cream onto the itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days as needed.  
children under 12 years of age: ask a doctor

**Other Information**  
do not use if sealed pouch is torn, open or incompletely sealed  
do not use if seal over tube opening has been punctured  
do not purchase if carton is open  
store at 20°-25°C (68°-77°F)  
this is a 1-dose treatment. Most women do not get complete relief of their symptoms in just 1 day. Most women get some improvement in 1 day and complete relief by 7 days.  
Keep this carton for future reference. It contains important information

**Inactive Ingredients**  
vaginal insert: gelatin, glycerin, leucithin, mineral oil, titanium dioxide, white petrolatum  
external cream: benzoic acid, cetyl alcohol, purified water, stearyl alcohol, 60, potassium hydroxide, propylene glycol, isopropyl myristate, polysorbate 80, potassium hydroxide, purified water, stearyl alcohol

**Questions?** If you have any questions or comments,  
please call 1-877-666-4782

**Warnings**  
Do not use if you have never had a vaginal yeast infection diagnosed by a doctor.  
Ask a doctor before use if you have:  
vaginal itching and discomfort for the first time  
lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.  
vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.  
been exposed to the human immunodeficiency virus (HIV) that causes AIDS.  
Ask a doctor or pharmacist before use if you are taking the prescription blood thinning medicine warfarin, because bleeding or bruising may occur.  
When using this product:  
do not use tampons, douches, spermicides or other vaginal products.  
Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).  
do not have vaginal intercourse  
do not increase in vaginal burning, itching or irritation may occur  
if you do not get complete relief ask a doctor before using another product  
Stop use and ask a doctor if:  
symptoms do not get better in 3 days  
you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting.

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

KAREN M MAHONEY  
07/22/2020 01:00:56 PM