

UNPRINTED AREA  
FOR LOT & EXP  
UNVARNISHED

# Advil

## COLD & SINUS

### LIQUI-GELS

Relieves Sinus Pressure, Nasal Congestion and Fever

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

# Advil

## COLD & SINUS

Non-Drowsy

**Ibuprofen 200 mg > Pain Reliever/Fever Reducer (NSAID)**  
**Pseudoephedrine HCl 30 mg > Nasal Decongestant**

Relieves Sinus Pressure,  
Nasal Congestion and Fever

### LIQUI-GELS

32 LIQUID-FILLED CAPSULES

**Advil** COLD & SINUS LIQUI-GELS

Relieves Sinus Pressure, Nasal Congestion and Fever

**Advil** COLD & SINUS LIQUI-GELS

Relieves Sinus Pressure, Nasal Congestion and Fever

**Advil** COLD & SINUS LIQUI-GELS

Relieves Sinus Pressure, Nasal Congestion and Fever

**Advil** COLD & SINUS LIQUI-GELS

Relieves Sinus Pressure, Nasal Congestion and Fever

**Advil** COLD & SINUS LIQUI-GELS

Relieves Sinus Pressure, Nasal Congestion and Fever

**Advil** COLD & SINUS LIQUI-GELS

Relieves Sinus Pressure, Nasal Congestion and Fever

**Advil** COLD & SINUS LIQUI-GELS

**Directions**  
Do not take more than directed. The smallest effective dose should be used. Adults and children 12 years of age and over: take 1 capsule every 4 to 6 hours while symptoms persist. If symptoms do not respond to 1 capsule, 2 capsules may be used. Do not use more than 6 capsules in any 24-hour period unless directed by a doctor. Children under 12 years of age: do not use.

**Drug Facts (continued)**

**Other information**  
Each capsule contains: potassium 20 mg. Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). Read all warnings and directions before use. Keep carton.

**Inactive ingredients**  
D&C yellow no. 10, FD&C red no. 40, glycol, potassium hydroxide, purified water, sorbitol sorbitan solution.

**Questions or comments?**  
Call weekdays from 9 AM to 5 PM EST toll free at 1-800-88-ADVIL.

**Drug Facts (continued)**

**Do not use**  
If you have ever had an allergic reaction to any other NSAID, or if you are taking aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Ask a doctor before use if you are**  
If you are taking a diuretic, due to an enlarged prostate gland, or had a stroke. You have a history of stomach problems, such as heartburn or fever reducers. You have problems or serious side effects from taking pain relievers. Stomach bleeding warning applies to you. Stomach bleeding warning applies to you. Stomach bleeding warning applies to you.

**When using this product**  
Take with food or milk if stomach upset occurs. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: vomit blood, feel faint, have bloody or black stools, have stomach pain that does not get better.

**Drug Facts (continued)**

**When using this product**  
Take with food or milk if stomach upset occurs. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: vomit blood, feel faint, have bloody or black stools, have stomach pain that does not get better.

**Drug Facts (continued)**

**Warnings**  
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others). Have 3 or more alcoholic drinks every day while using this product. Minor body aches and pains, nasal congestion, sinus pressure, headache, fever.

**Purposes**  
Sulfonated ibuprofen equal to 200 mg ibuprofen (NSAID). Pain reliever/fever reducer (present as the free acid and potassium salt). Pseudoephedrine HCl 30 mg. Nasal decongestant.

**Uses**  
Temporarily relieves these symptoms associated with the common cold or flu: temporarily relieves these symptoms associated with the common cold or flu: temporarily relieves these symptoms associated with the common cold or flu.

Product inside sealed in plastic blister with foil backing. Do Not Use if plastic blister or foil barrier is broken.

For most recent product information, visit [www.advil.com](http://www.advil.com). Distributed by: GSK Consumer Healthcare, Warren, NJ 07059. Trademarks owned or licensed by GSK. ©2021 GSK or licensor. LIQUI-GELS® is a trademark or registered trademark of Galeant Pharma Solutions.

000065783

FPO

3 0573 0184 32 8

000065783

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

VALERIE S PRATT  
11/19/2021 03:11:18 PM