

PRINTED AREA FOR DATE & LOT
LASER CODING (VARNISHED)

000070871

FPO

3 0573 0167 43 7

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredients (in each capsule)

Diphenhydramine hydrochloride 25 mg..... Nighttime sleep-aid

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)..... Pain reliever (present as the free acid and potassium salt)

nonsteroidal anti-inflammatory drug

Purposes

■ for relief of occasional sleeplessness when associated with minor aches and pains

■ helps you fall asleep and stay asleep

Uses

■ for relief of occasional sleeplessness when associated with minor aches and pains

■ helps you fall asleep and stay asleep

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ facial swelling ■ asthma (wheezing)

■ shock ■ skin redness ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

■ are age 60 or older

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

■ unless you have time for a full night's sleep

■ in children under 12 years of age

■ right before or after heart surgery

■ with any other product containing diphenhydramine, even one used on skin

■ if you have sleeplessness without pain

Ask a doctor before use if

■ stomach bleeding warning applies to you

■ you have problems or serious side effects from taking pain relievers or fever reducers

■ you have a history of stomach problems, such as heartburn

■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

■ you are taking a diuretic

■ you have a breathing problem such as emphysema or chronic bronchitis

■ you have diabetes

■ you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

■ taking sedatives or tranquilizers, or any other sleep-aid

■ under a doctor's care for any continuing medical illness

■ taking any other antihistamines

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin

■ taking any other drug

When using this product

■ drowsiness will occur

■ do not drive a motor vehicle or operate machinery while with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:

■ vomit blood ■ have bloody or black stools

■ have stomach pain that does not get better

■ you have symptoms of heart problems or stroke:

■ chest pain ■ trouble breathing

■ weakness in one part or side of body

■ slurred speech ■ leg swelling

■ pain gets worse or lasts more than 10 days

■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness

■ redness or swelling is present in the painful area

■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed

■ adults and children 12 years and over: take 2 capsules at bedtime

■ do not take more than 2 capsules in 24 hours

Other information


■ each capsule contains: potassium 20 mg

■ read all warnings and directions before use. Keep carton, store at 20-25°C (68-77°F)

■ avoid excessive heat above 40°C (104°F)

■ protect from light

Inactive ingredients: D&C red no. 33, FD&C blue no. 1, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution



Solubilized Ibuprofen, 200 mg / Diphenhydramine HCl, 25 mg Pain Reliever (NSAID)/Nighttime Sleep-Aid

40 Liquid Filled Capsules

Drug Facts (continued)

Questions or comments?

Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

For most recent product information, visit www.advil.com

©2021 GSK Consumer Healthcare, Warren, NJ 07059

Trademarks are owned by or licensed to the GSK group of companies

Patents: www.products.com

LIQUI-GELS® is a trademark or registered trademark of Catalent Pharma Solutions.

000070871

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredients (in each capsule)

Diphenhydramine hydrochloride 25 mg..... Nighttime sleep-aid
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*..... Pain reliever (present as the free acid and potassium salt), nonsteroidal anti-inflammatory drug

Purposes

■ for relief of occasional sleeplessness when associated with minor aches and pains
■ helps you fall asleep and stay asleep

Uses

■ for relief of occasional sleeplessness when associated with minor aches and pains
■ helps you fall asleep and stay asleep

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin redness ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinner (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product.
■ take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ unless you have time for a full night's sleep
■ in children under 12 years of age
■ right before or after heart surgery
■ with any other product containing diphenhydramine, even one used on skin
■ if you have sleeplessness without pain

Ask a doctor before use if

■ stomach bleeding warning applies to you
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you are taking a diuretic
■ you have a breathing problem such as emphysema or chronic bronchitis
■ you have glaucoma
■ you have trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

■ taking sedatives or tranquilizers, or any other sleep-aid
■ under a doctor's care for any continuing medical illness
■ taking any other anti-inflammatories
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
■ taking any other drug

When using this product

■ avoid alcoholic drinks
■ do not drive a motor vehicle or operate machinery
■ take with food or milk if stomach upset occurs

Drug Facts (continued)

You use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke:
■ chest pain ■ trouble breathing
■ sudden speech ■ leg swelling
■ pain gets worse or lasts more than 10 days
■ sleeplessness persists continuously for more than 2 weeks; insomnia may be a symptom of a serious underlying medical illness.
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not take more than 2 capsules in 24 hours
■ adults and children 12 years and over: take 2 capsules at bedtime

Other information

■ each capsule contains: potassium 20 mg
■ read all warnings and directions before use. Keep carton.
■ store at 20-25°C (68-77°F)
■ avoid excessive heat above 40°C (104°F)
■ protect from light.

Inactive ingredients D&C red no. 33, FD&C blue no. 1, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

PRINTED AREA FOR DATE & LOT
LASER CODING (VARNISHED)

UNPRINTED AREA
FOR DATE & LOT
(UNVARNISHED)

25% MORE FREE

Advil PM LIQUI-GELS

Solubilized Ibuprofen, 200 mg/
Diphenhydramine HCl, 25 mg
Pain Reliever (NSAID)/Nighttime Sleep-Aid

40 50
Liquid Filled Capsules

10 Free LIQUI-GELS

For most recent product information, visit www.Advill.com
©2021 GSK Consumer Healthcare, Warren, NJ 07099
Distributed by: GSK Consumer Healthcare, Warren, NJ 07099
Trademarks are owned by or licensed to the GSK group of companies
Patents: www.productpatents.com
LIQUI-GELS® is a trademark or registered trademark of Catalent Pharma Solutions.

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Questions or comments?
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

000070872

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



PRINTED AREA FOR DATE & LOT
LASER CODING (VARNISHED)

000070873

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



PRINTED AREA FOR DATE & LOT
LASER CODING (VARNISHED)

000070874

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
01/11/2022 12:12:48 PM