


FRONT



BACK

LASER ENCODING AREA FOR LOT AND EXPIRATION INFORMATION. PRINTED IN ONE SOLID COLOR, BLACK OR DARK BLUE.

BACK

**208495**


MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• BARLINES, HAIRLINES	N/A

TOP (BACK)

**Warnings**

- **Ask your doctor before use** if you are pregnant, under a doctor's care for any continuing medical illness, age 60 or over, taking any other drug or have stomach problems.
- This product may cause a **severe allergic reaction**, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
- **Do not use** this product if you have ever had an allergic reaction to any pain reliever/fever reducer; unless you have time for a full night's sleep; in children under 12 years or with any other product containing diphenhydramine.
- **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you
  - are age 60 or older
  - have had stomach ulcers or bleeding problems
  - take a blood thinning (anticoagulant) or steroid drug
  - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
  - have 3 or more alcoholic drinks every day while using this product
  - take more or for a longer time than directed.
- **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

COMPOSITE



BASE

- **When using this product** drowsiness will occur. Avoid alcoholic drinks. Do not drive a motor vehicle or operate machinery. Take with food or milk if stomach upset occurs.
- Directions** ■ **do not take more than directed**
- adults and children 12 years and over: take 2 capsules at bedtime
- do not take more than 2 capsules in 24 hours
- Each capsule contains: **potassium 20 mg**
- Store at 20-25°C (68-77°F)
- Avoid excessive heat above 40°C (104°F). Protect from light.
- Questions or comments?** Call weekdays 9 AM to 5 PM EST at **1-800-88-ADVIL**

For most recent product information, visit [www.Advii.com](http://www.Advii.com)  
Dist. by: Haleon, Warren, NJ 07059  
Trademarks owned or licensed by Haleon.  
©2023 Haleon or licensor.  
LIQUI-GELS is a trademark or registered trademark of Catalent Pharma Solutions.

**DO NOT USE unless vial was sealed in unbroken plastic blister on cardboard card.**

208497

TOP (FACE)

**Solubilized Ibuprofen, 200 mg/  
Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)  
/Nighttime Sleep-Aid**

**8 Liquid Filled Capsules**

**READ AND KEEP CARD FOR COMPLETE WARNINGS AND INFORMATION**

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**OPEN HERE TO VIEW MORE PRODUCT INFORMATION**

LOT EXP AREA  
NO VARNISH

208497

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE	.5 pt

FRONT



BACK



↑  
LOT & EXP AREA

208499

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	7.0 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each capsule)</b></p> <p>Diphenhydramine hydrochloride 25 mg.....Nighttime sleep-aid Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)* .....Pain reliever (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug</p>		<p><b>Purposes</b></p> <p>■ for relief of occasional sleeplessness when associated with minor aches and pains ■ helps you fall asleep and stay asleep</p>	<p><b>Drug Facts (continued)</b></p> <p>Symptoms may include:</p> <ul style="list-style-type: none"> <li>■ hives ■ facial swelling</li> <li>■ asthma (wheezing)</li> <li>■ shock ■ skin reddening</li> <li>■ rash ■ blisters</li> </ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> <li>■ are age 60 or older</li> <li>■ have had stomach ulcers or bleeding problems</li> <li>■ take a blood thinning (anticoagulant) or steroid drug</li> <li>■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> <li>■ have 3 or more alcoholic drinks every day while using this product</li> <li>■ take more or for a longer time than directed</li> </ul>
<p><b>Warnings</b></p> <p><b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.</p>			

208500

LIFT HERE

PANEL 1

<p><b>Drug Facts (continued)</b></p> <p><b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p><b>Do not use</b> ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ unless you have time for a full night's sleep ■ in children under 12 years of age ■ right before or after heart surgery ■ with any other product containing diphenhydramine, even one used on skin ■ if you have sleeplessness without pain</p> <p><b>Ask a doctor before use if</b> ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers</p>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ you have a history of stomach problems, such as heartburn</li> <li>■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke</li> <li>■ you are taking a diuretic</li> <li>■ you have a breathing problem such as emphysema or chronic bronchitis</li> <li>■ you have glaucoma</li> <li>■ you have trouble urinating due to an enlarged prostate gland</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>■ taking sedatives or tranquilizers, or any other sleep-aid</li> <li>■ under a doctor's care for any continuing medical illness</li> <li>■ taking any other antihistamines</li> <li>■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin</li> <li>■ taking any other drug</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>■ drowsiness will occur</li> <li>■ avoid alcoholic drinks</li> </ul>
---	---

PANEL 2

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ do not drive a motor vehicle or operate machinery</li> <li>■ take with food or milk if stomach upset occurs</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> <li>■ feel faint</li> <li>■ vomit blood</li> <li>■ have bloody or black stools</li> <li>■ have stomach pain that does not get better</li> </ul> </li> <li>■ you have symptoms of heart problems or stroke: <ul style="list-style-type: none"> <li>■ chest pain</li> <li>■ trouble breathing</li> <li>■ weakness in one part or side of body</li> <li>■ slurred speech</li> <li>■ leg swelling</li> </ul> </li> <li>■ pain gets worse or lasts more than 10 days</li> <li>■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.</li> <li>■ redness or swelling is present in the painful area</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ any new symptoms appear</li> </ul> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ do not take more than directed</li> <li>■ adults and children 12 years and over: take 2 capsules at bedtime</li> <li>■ do not take more than 2 capsules in 24 hours</li> </ul>
--	--


BASE PANEL

208500

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



FRONT



LOT & EXP AREA

BACK

**208498**

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	N/A
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**Drug Facts** (continued)

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ unless you have time for a full night's sleep ■ in children under 12 years of age
- right before or after heart surgery ■ with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

**Ask a doctor before use if**

- stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis ■ you have glaucoma ■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other antihistamines ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product**

- drowsiness will occur ■ avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk, if stomach upset occurs

**Drug Facts** (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed ■ adults and children 12 years and over: take 2 capsules at bedtime
- do not take more than 2 capsules in 24 hours

**Other information**

- each capsule contains: potassium 20 mg
- read all warnings and directions before use.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)
- protect from light

**Inactive ingredients**

D&C red no. 33, FD&C blue no. 1, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Questions or comments?**  
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
Dist. by: Haleon, Warren, NJ 07059  
Trademarks owned or licensed by Haleon.  
©2023 Haleon or licensor.  
LIQUI-GELS is a trademark or registered trademark of Catalent Pharma Solutions.

208502

**81% FPO**

3 05736 55631 0

**DO NOT USE IF SEAL AROUND CAP IS BROKEN OR MISSING. ADVIL SAFETY SEALED.**

**Advil PM LIQUI-GELS minis** 8 Liquid Filled Capsules

Solubilized Ibuprofen, 200 mg/  
Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID) / Nighttime Sleep-Aid

**Drug Facts**

**Active ingredients (in each capsule)**      **Purposes**

Diphenhydramine hydrochloride 25 mg..... Nighttime sleep-aid

Solubilized ibuprofen equal to 200 mg  
ibuprofen (NSAID) ..... Pain reliever  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**LIET HERE for More Drug Facts**

**Drug Facts** (continued)

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint ■ vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain ■ trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

ADVIL SAFETY SEALED ADVIL SAFETY SEALED ADVIL SAFETY SEALED  
TEAR HERE TO OPEN

208502

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**DRUG FACTS (CONTINUED)**

**Drug Facts**  
**Active ingredients (in each capsule)**  
Diphenhydramine hydrochloride 25 mg, Nighttime sleep-aid  
Solubilized ibuprofen equal to  
200 mg ibuprofen (NSAID) (present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Purposes**  
Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ for relief of occasional sleeplessness when associated with minor aches and pains ■ helps you fall asleep and stay asleep

**Warnings**  
**Alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, skin redness, rash, and blisters. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Drug Facts (continued)**  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any of the pain reliever/fever reducer ■ unless you have time for a full night's sleep ■ in children under 12 years of age ■ right before or after heart surgery ■ with any other product containing diphenhydramine, even one used on skin ■ if you have sleeplessness without pain

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have a breathing problem such as emphysema or chronic bronchitis ■ you have diabetes ■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers, or any other sleep-aid ■ under a doctor's care for any continuing medical illness ■ taking any other antihistamines ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

**When using this product** ■ drowsiness will occur ■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery ■ take with food or milk if stomach upset occurs

**Drug Facts (continued)**  
**Stomach bleeding warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any of the pain reliever/fever reducer ■ unless you have time for a full night's sleep ■ in children under 12 years of age ■ right before or after heart surgery ■ with any other product containing diphenhydramine, even one used on skin ■ if you have sleeplessness without pain

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have a breathing problem such as emphysema or chronic bronchitis ■ you have diabetes ■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers, or any other sleep-aid ■ under a doctor's care for any continuing medical illness ■ taking any other antihistamines ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

**When using this product** ■ drowsiness will occur ■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery ■ take with food or milk if stomach upset occurs

**Drug Facts (continued)**  
**Stomach bleeding warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any of the pain reliever/fever reducer ■ unless you have time for a full night's sleep ■ in children under 12 years of age ■ right before or after heart surgery ■ with any other product containing diphenhydramine, even one used on skin ■ if you have sleeplessness without pain

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have a breathing problem such as emphysema or chronic bronchitis ■ you have diabetes ■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers, or any other sleep-aid ■ under a doctor's care for any continuing medical illness ■ taking any other antihistamines ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

**When using this product** ■ drowsiness will occur ■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery ■ take with food or milk if stomach upset occurs


UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

208515

3 05736 55756 0

FPO



**Advil<sup>®</sup> PM**  
**Liqui-Gels<sup>®</sup> minis**

Solubilized Ibuprofen, 200 mg/  
Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid


**20**  
Liquid Filled Capsules

\*Compared to Advil PM Liqui-Gels

**208515**


DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP (FACE)  
PANEL 1




FOR LOT & EXPIRATION AREA,  
UNVARNISHED AND PRINTED  
IN BLACK OR DARK BLUE.

BASE  
PANEL 3




TOP (BACK)  
PANEL 2





208513

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt



FOR LOT & EXPIRATION AREA,  
UNVARNISHED AND PRINTED  
IN BLACK OR DARK BLUE.



208525

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt

**UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)**

**PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)**

208526

**DRUG FACTS MODIFIED TEXT DEFINED**

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**DRUG FACTS (continued)**

**Active ingredients (in each capsule)** **Purposes**  
Diphenhydramine hydrochloride 25 mg, ... Nighttime sleep-aid  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID) \* ..... Pain reliever  
(present as the free acid and potassium salt)  
Nonsteroidal anti-inflammatory drug

**Uses** ■ For relief of occasional sleeplessness when associated with minor aches and pains ■ Helps you fall asleep and stay asleep

**Warnings**  
**Allegry alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
■ skin reddening ■ rash ■ hiccups  
■ dizziness ■ fainting  
If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Drug Facts (continued)**  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ If you have ever had an allergic reaction to any other pain reliever/fever reducer ■ unless you have time for a full night's sleep ■ in children under 12 years of age ■ right before or after heart surgery ■ with any other product containing diphenhydramine, even one used on skin ■ if you have sleeplessness without pain

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have a breathing problem such as emphysema, or chronic bronchitis ■ you have glaucoma ■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** ■ taking statins or tranquilizers, or any other sleep-aid ■ taking a doctor's care for any continuing medical illness ■ taking any other antihistamines ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

**When using this product** ■ drowsiness will occur ■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery ■ shake with food or milk if stomach upset occurs

**Drug Facts (continued)**  
**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding:  
■ heart burn ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing ■ weakness in one part of side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ sleeplessness persists continuously for more than 2 weeks ■ nausea may be a symptom of a serious underlying medical illness ■ redness or swelling is present in the genital area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed ■ adults and children 12 years and over: take 2 capsules at bedtime ■ do not take more than 2 capsules in 24 hours

**Other information** ■ each capsule contains potassium 20 mg ■ read all warnings and directions before use. Keep carton. ■ store at 20°-25° C (68°-77° F)

**10 FREE Liquid Filled Capsules**

**Advil<sup>®</sup> PM LIQUI-GELS<sup>®</sup> minis**

Solubilized Ibuprofen, 200 mg /  
Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid

**20 30**  
Liquid Filled Capsules

\*Compared to Advil PM Liqui-Gels

**Do Not Use if seal under bottle cap is broken or missing.**

For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
©2024 Haleson group of companies or its licensor.  
Distributed by Haleson Warren, MI 48090  
Haleson group of companies.  
Trademarks are owned by or licensed to the  
Liqui-Gels and Advil trademarks or registered trademarks  
of Calient Prime Solutions.


**Questions or comments?**  
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

**Inactive ingredients** D&C red no. 33, FD&C blue no. 1, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution

**Drug Facts (continued)**  
■ protect from light  
■ avoid excessive heat above 40°C (104°F)




**TOP (FACE)  
PANEL 1**




FOR LOT & EXPIRATION AREA,  
UNVARNISHED AND PRINTED  
IN BLACK OR DARK BLUE.

**BASE  
PANEL 3**




**TOP (BACK)  
PANEL 2**





**208516**

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.3 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt



FOR LOT & EXPIRATION AREA,  
UNVARNISHED AND PRINTED  
IN BLACK OR DARK BLUE.



208527

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt



**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
*(In each capsule)*

**Purposes**  
Diphenhydramine  
hydrochloride 25 mg ..... Nighttime sleep-aid  
Solubilized ibuprofen equal to  
200 mg ibuprofen (NSAID) ..... Pain reliever  
(present as the free acid and potassium salt)  
nonsteroidal anti-inflammatory drug

**Uses**  
■ for relief of occasional sleeplessness when associated  
with minor aches and pains  
■ helps you fall asleep and stay asleep

**Warnings**  
**Allergic alert:** ibuprofen may cause a severe allergic  
reaction, especially in people allergic to aspirin.  
Symptoms may include:  
■ hives ■ facial swelling ■ rash ■ hives  
■ sneez ■ skin redness ■ asthma (wheezing)  
If an allergic reaction occurs, stop use and seek medical help  
right away.  
**Stomach bleeding warning:** This product contains an  
NSAID, which may cause severe stomach bleeding.  
The chance is higher if you  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or  
nonprescription NSAIDs (aspirin, ibuprofen, naproxen,  
or others)  
■ have 3 or more alcoholic drinks every day while using  
this product  
■ take more or for a longer time than directed

**Drug Facts (continued)**  
**Heart attack and stroke warning:** NSAIDs, except aspirin,  
increase the risk of heart attack, heart failure, and stroke.  
These can be fatal. The risk is higher if you use more than  
directed or for longer than directed.  
**Do not use**  
■ if you have ever had an allergic reaction to any other pain  
reliever/fever reducer  
■ unless you have time for a full night's sleep  
■ in children under 12 years of age  
■ right before or after heart surgery  
■ with any other product containing diphenhydramine, even  
one used on skin  
■ if you have sleeplessness without pain

**Ask a doctor before use if:**  
■ stomach bleeding warning applies to you  
■ you have problems or serious side effects from taking  
pain relievers or fever reducers  
■ you have a history of stomach problems, such as  
heartburn  
■ you have high blood pressure, heart disease, liver  
cirrhosis, kidney disease, asthma, or had a stroke  
■ you are taking a diuretic  
■ you have a breathing problem such as emphysema or  
chronic bronchitis  
■ you have glaucoma  
■ you have trouble urinating due to an enlarged prostate  
gland

**Ask a doctor or pharmacist before use if you are:**  
■ taking sedatives or tranquilizers, or any other sleep-aid  
■ under a doctor's care for any continuing medical illness  
■ being any other antihistamines  
■ being aspirin for heart attack or stroke, because ibuprofen  
may decrease the benefit of aspirin  
■ taking any other drug

**When using this product:**  
■ avoid alcoholic drinks  
■ do not drive a motor vehicle or operate machinery  
take with food or milk if stomach upset occurs

**Drug Facts (continued)**  
**Drug Facts (continued)**  
**Stop use and ask a doctor if:**  
■ you experience any of the following signs of  
stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing  
■ weakness in one part of body  
■ dizziness or lightheadedness  
■ pain gets worse or lasts more than 10 days  
■ sleeplessness persists continuously for more than  
2 weeks. Insomnia may be a symptom of a serious  
underlying medical illness.  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before  
use.** It is especially important not to use ibuprofen at 20 weeks  
or later in pregnancy unless directly ordered to do so by a  
doctor because it may cause problems in the unborn child or  
complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical  
help or contact a Poison Control Center right away.

**Directions** ■ **do not take more than directed**  
■ do not take more than 2 capsules in 24 hours  
■ adults and children 12 years and over: take 2 capsules  
at bedtime

**Other information**  
■ each capsule contains potassium 20 mg  
■ read all warnings and directions before use. Keep carton,  
sore at 20-25°C (68-77°F)  
■ avoid excessive heat above 40°C (104°F)  
■ protect from light

**Inactive ingredients** D&C red no. 33, D&C blue no. 1,  
gelatin, medium-chain triglycerides, pharmaceutical ink,  
polyethylene glycol, potassium hydroxide, purified water,  
sodium stannate solution

208519

**FPO**

615952

1 6 5 5 7 3 6 5 5 7 5 1 3

**PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)**

**UNPRINTED AREA  
FOR DATE & LOT  
(UNVARNISHED)**

HALEON

**NEW** Smaller Capsule  
**Same Strength\***

# Advil<sup>®</sup> PM

## LIQUI-GELS<sup>®</sup>

Solubilized Ibuprofen, 200 mg /  
Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)/Nighttime Sleep-Aid

**Actual Size**

**minis**

**80**  
**Liquid Filled Capsules**

\*Compared to Advil PM Liqui-Gels

For most recent product information, visit [www.advil.com](http://www.advil.com)  
©2024 Haleon group of companies or its licensor.  
Distributed by: Haleon, Warren, NJ 07059  
Haleon group of companies or its licensor.  
Trademarks are owned by or licensed to the  
Haleon group of companies.

**Do Not Use if seal under bottle cap imprinted with  
"SEALED for YOUR PROTECTION" is broken or missing.**

Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

**Questions or comments?**

**Drug Facts (continued)**

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Reference ID: 5443434

**HALEON**

# Advil<sup>®</sup>PM

## LIQUI-GELS<sup>®</sup>

**minis**

Solubilized Ibuprofen, 200 mg /  
 Diphenhydramine HCl, 25 mg  
**Pain Reliever (NSAID) /  
 Nighttime Sleep-Aid**

**80**  
Liquid Filled  
Capsules

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

- **Ask your doctor before use** if you are pregnant, under a doctor's care for any continuing medical illness, age 60 or over, taking any other drug or have stomach problems.
- This product may cause a **severe allergic reaction**, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
- **Do not use** this product if you have ever had an allergic reaction to any pain reliever/fever reducer; unless you have time for a full night's sleep; in children under 12 years or with any other product containing diphenhydramine.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**When using this product** drowsiness will occur. Avoid alcoholic drinks. Do not drive a motor vehicle or operate machinery. Take with food or milk if stomach upset occurs

**Directions** ■ **do not take more than directed**

- adults and children 12 years and over: take 2 capsules at bedtime
- do not take more than 2 capsules in 24 hours

Each capsule contains: **potassium 20 mg**  
 Store at 20-25°C (68-77°F)  
 Avoid excessive heat above 40°C (104°F). Protect from light.

**Questions or comments?**  
 Call weekdays 9 AM to 5 PM EST at **1-800-88-ADVIL**  
 For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
 Dist. by: Haleon, Warren, NJ 07059  
 Trademarks owned or licensed by Haleon. ©2023 Haleon or licensor.  
 LIQUI-GELS is a trademark or registered trademark of Catalent Pharma Solutions.

208518

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

208518

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5.25 pt
• SUBHEADINGS/BODY TEXT	5.25 pt
• LEADING	5.5 pt
• BULLETS	4.25 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt



**HALEON**

**Advil<sup>®</sup> PM**  
**LIQUI-GELS<sup>®</sup>**

Solubilized Ibuprofen, 200 mg /  
 Diphenhydramine HCl, 25 mg  
**Pain Reliever (NSAID) /  
 Nighttime Sleep-Aid**

**minis**

**100**  
 Liquid Filled  
 Capsules

**READ AND KEEP CARTON FOR COMPLETE  
 WARNINGS AND INFORMATION**

**Do Not Use** if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

**Uses**  
 ■ for relief of occasional sleeplessness when associated with minor aches and pains  
 ■ helps you fall asleep and stay asleep

**Warnings**  
 ■ **Ask your doctor before use** if you are pregnant, under a doctor's care for any continuing medical illness, age 60 or over, taking any other drug or have stomach problems.  
 ■ This product may cause a **severe allergic reaction**, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.  
 ■ **Do not use** this product if you have ever had an allergic reaction to any pain reliever/fever reducer; unless you have time for a full night's sleep; in children under 12 years or with any other product containing diphenhydramine.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**When using this product** drowsiness will occur. Avoid alcoholic drinks.  
 Do not drive a motor vehicle or operate machinery.  
 Take with food or milk if stomach upset occurs

**Directions** ■ **do not take more than directed**  
 ■ adults and children 12 years and over: take 2 capsules at bedtime ■ do not take more than 2 capsules in 24 hours  
 Each capsule contains: **potassium 20 mg**  
 Store at 20-25°C (68-77°F)  
 Avoid excessive heat above 40°C (104°F). Protect from light.

**Questions or comments?**  
 Call weekdays 9 AM to 5 PM EST at **1-800-88-ADVIL**  
 For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
 Dist. by: Haleon, Warren, NJ 07059  
 Trademarks owned or licensed by Haleon. ©2023 Haleon or licensor.  
 LIQUI-GELS is a trademark or registered trademark of Catalent Pharma Solutions.

208529

UNVARNISHED AND PRINTED  
 IN BLACK OR DARK BLUE FOR  
 LOT & EXPIRATION AREA.

208529

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5.25 pt
• SUBHEADINGS/BODY TEXT	5.25 pt
• LEADING	5.5 pt
• BULLETS	4.25 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**(In each capsule)**

**Active Ingredients**  
Diphenhydramine hydrochloride 25 mg  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
(present as the free acid and potassium salt)  
nonsteroidal anti-inflammatory drug

**Purposes**  
Nighttime sleep-aid  
Pain reliever

**Uses**  
for relief of occasional sleeplessness when associated with minor aches and pains  
highly you fall asleep and stay asleep

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
swelling of the face, lips, throat, and tongue  
hives  
skin redness  
rash  
blistering  
If an allergic reaction occurs, stop use and seek medical help (right away).  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
are age 60 or older  
have had stomach ulcers or bleeding problems  
take a blood thinning (anticoagulant) or steroid drug  
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
have 3 or more alcoholic drinks every day while using this product  
take more or for a longer time than directed

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
if you have ever had an allergic reaction to any other pain reliever/fever reducer  
unless you have time for a full night's sleep  
in children under 12 years of age  
right before or after heart surgery  
with any other product containing diphenhydramine, even one used on skin  
if you have sleeplessness without pain

**Ask a doctor before use if:**  
stomach bleeding warning applies to you  
you have problems or serious side effects from taking pain relievers or fever reducers  
you have a history of stomach problems, such as heartburn  
you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
you are taking a diuretic  
you have a breathing problem such as emphysema or chronic bronchitis  
you have glaucoma  
you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are:**  
taking sedatives or tranquilizers, or any other sleep-aid  
under a doctor's care for any continuing medical illness  
taking any other anti-inflammatories  
taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin  
taking any other drug

**When using this product:**  
avoid alcoholic drinks  
do not drive a motor vehicle or operate machinery  
take with food or milk if stomach upset occurs

**Drug Facts (continued)**

**Stop use and ask a doctor if:**  
you experience any of the following signs of stomach bleeding:  
black stool  
vomit blood  
have bloody or black stools  
ever stomach pain that does not get better  
you have symptoms of heart problems or stroke:  
chest pain  
trouble breathing  
weakness in one part or side of body  
slurred speech  
egg swelling  
pain gets worse or lasts more than 10 days  
sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.  
redness or swelling is present in the painful area  
any new symptoms appear

**If pregnant or breast-feeding:** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
**do not take more than directed**  
adults and children 12 years and over: take 2 capsules at bedtime  
do not take more than 2 capsules in 24 hours

**Other information**  
each capsule contains: potassium 20 mg  
read all warnings and directions before use. Keep carton.  
store at 20-25°C (68-77°F)  
avoid excessive heat above 40°C (104°F)  
protect from light

**Inactive ingredients:** D&C red no. 33, D&C blue no. 1, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sodium solution

208530

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

UNPRINTED AREA FOR DATE & LOT FOR DATE & LOT (UNVARNISHED)

**20 FREE Liquid Filled Capsules**

**Smaller Capsule Same Strength\***

**HALEON**

**Advil<sup>®</sup> PM LIQUI-GELS<sup>®</sup> minis**

Solubilized Ibuprofen, 200 mg / Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)/Nighttime Sleep-Aid

**80 100**  
Liquid Filled Capsules

\*Compared to Advil PM Liqui-Gels

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**HALEON**

**Advil<sup>®</sup> PM**  
**LIQUI-GELS<sup>®</sup>**

Solubilized Ibuprofen, 200 mg /  
 Diphenhydramine HCl, 25 mg  
 Pain Reliever (NSAID) /  
 Nighttime Sleep-Aid

**minis**

**120**  
 Liquid Filled  
 Capsules

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Do Not Use** if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

- **Ask your doctor before use** if you are pregnant, under a doctor's care for any continuing medical illness, age 60 or over, taking any other drug or have stomach problems.
- This product may cause a **severe allergic reaction**, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
- **Do not use** this product if you have ever had an allergic reaction to any pain reliever/fever reducer; unless you have time for a full night's sleep; in children under 12 years or with any other product containing diphenhydramine.

**Directions**

- **do not take more than directed**
- adults and children 12 years and over: take 2 capsules at bedtime
- do not take more than 2 capsules in 24 hours
- Each capsule contains: **potassium 20 mg**
- Store at 20-25°C (68-77°F)
- Avoid excessive heat above 40°C (104°F). Protect from light.

**Questions or comments?**  
 Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL  
 For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
 Dist. by: Haleon, Warren, NJ 07059  
 Trademarks owned or licensed by Haleon.  
 ©2023 Haleon or licensor.  
 LIQUI-GELS is a trademark or registered trademark of Catalent Pharma Solutions.

208520

UNVARNISHED AND PRINTED  
 IN BLACK OR DARK BLUE FOR  
 LOT & EXPIRATION AREA.

**208520**

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	5.25 pt
• SUBHEADINGS/BODY TEXT	5.25 pt
• LEADING	5.75 pt
• BULLETS	4.25 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredients (in each capsule)**

Diphenhydramine hydrochloride, 25 mg..... Nighttime sleep-aid  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*..... Pain reliever  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Purposes**

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- swollen
- skin redness
- itching

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you:

- are age 65 or older
- are taking stomach ulcers or bleeding problems medicine
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take a liver drug (including prescription or herbal)
- take NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more than 3 more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other antihistamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

**When using this product**

- do not take more than directed
- do not take with alcohol
- do not take with food or milk if stomach upset occurs

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- you have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part of side of body
  - slurred speech
  - stained speech
- your pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- adults and children 12 years and over: take 2 capsules at bedtime
- do not take more than 2 capsules in 24 hours

**Other information**

- each capsule contains potassium 20 mg
- read all warnings and directions before use. Keep carton, store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)
- protect from light


**Inactive ingredients:** D&C red no. 33, FD&C blue no. 1, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sodium sorbitan solution

**Questions or comments?**  
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

UNPRINTED AREA FOR DATE & LOT CODING (UNVARNISHED)

8 22925 55622 3  
FPO  
3 05736 55622 8  
129802



**Advil<sup>®</sup> PM<sup>®</sup> LIQUI-GELS<sup>®</sup> minis**

Solubilized Ibuprofen, 200 mg / Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)/ Nighttime Sleep-Aid

120 Liquid Filled Capsules

\*Compared to Advil PM Liqui-Gels

For most recent product information, visit [www.AdvIL.com](http://www.AdvIL.com)  
Distributed by: Haleon, Warren, NJ 07059  
Trademarks are owned by or licensed to the Haleon group of companies.  
©2024 Haleon group of companies or its licensor. LIQUI-GELS is a trademark or registered trademark of Galant Pharma Solutions.

208521	
DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
*(In each capsule)* ..... **Purposes**  
Diphenhydramine hydrochloride 25 mg..... Nighttime sleep-aid  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID) ..... Pain reliever  
..... Pain reliever  
..... Transdermal anti-inflammatory drug

**Uses**  
■ for relief of occasional sleeplessness when associated with minor aches and pains  
■ helps you fall asleep and stay asleep

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.  
Symptoms may include:  
■ hives  
■ facial swelling  
■ chest pain (wheezing)  
■ asthma (wheezing)  
■ skin redness/itching

■ if an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.  
The chance is higher if you  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ have taken aspirin, NSAIDs, or other pain relievers (e.g., ibuprofen or naproxen), or other NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ unless you have time for a full night's sleep  
■ in children under 12 years of age  
■ with any other product containing diphenhydramine, even one used on skin  
■ if you have sleeplessness without pain

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you  
■ you have problems or serious side effects from taking pain relievers or fever reducers  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
■ you have a bleeding problem such as emphysema or chronic bronchitis  
■ you have glaucoma  
■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**  
■ taking sedatives or tranquilizers, or any other sleep-aid under a doctor's care for any continuing medical illness  
■ taking any other antihistamines  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
■ taking any other drug

**When using this product**  
■ drowsiness will occur  
■ do not drive a motor vehicle or operate machinery  
■ take with food or milk if stomach upset occurs  
**Stop use and ask a doctor if**  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint

**Drug Facts (continued)**

■ vomit blood  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain  
■ trouble breathing  
■ trouble speaking  
■ pain gets worse or lasts more than 10 days  
■ sleeplessness persists continuously for more than 2 weeks.  
Insomnia may be a symptom of a serious underlying medical illness.  
■ dizziness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not take more than directed  
■ adults and children 12 years and over: take 2 capsules at bedtime  
■ do not take more than 2 capsules in 24 hours

**Other information**  
■ **inactive ingredients** 20 mg  
■ read all warnings and directions before use. Keep carton.  
■ store at 20-25°C (68-77°F)  
■ avoid excessive heat above 40°C (104°F)  
■ protect from light

**Questions or comments?**  
Call weekdays 9 AM to 9 PM EST at 1-800-88-ADUI

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

UNPRINTED AREA FOR DATE & LOT LASER CODING (UNVARNISHED)

208523

**HALEON**

**NEW Smaller Capsule**  
**Same Strength\***

# Advil<sup>®</sup> PM

## LIQUI-GELS<sup>®</sup>

Solubilized Ibuprofen, 200 mg/  
Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)/Nighttime Sleep-Aid

**Actual Size**

**minis**

**160**  
**Liquid Filled Capsules**


\*Compared to Advil PM Liqui-Gels

LIQUI-GELS is a trademark or registered trademark of Calant Pharma Solutions.  
©2024 Haleon group of companies or its licensor.  
LIQUI-GELS is a trademark or registered trademark of Haleon group of companies.  
Distributed by: Haleon, Warren, NJ 07059  
For most recent product information, visit [www.Advil.com](http://www.Advil.com)

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

**208523**

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	7.0 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



**HALÉON**

**Advil<sup>®</sup> PM**  
**LIQUI·GELS<sup>®</sup>**

Solubilized Ibuprofen, 200 mg /  
 Diphenhydramine HCl, 25 mg  
**Pain Reliever (NSAID) /  
 Nighttime Sleep-Aid**

**minis**

**160**  
**Liquid Filled  
 Capsules**

**READ AND KEEP CARTON FOR COMPLETE  
 WARNINGS AND INFORMATION**

**Do Not Use** if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

- **Ask your doctor before use** if you are pregnant, under a doctor's care for any continuing medical illness, age 60 or over, taking any other drug or have stomach problems.
- This product may cause a **severe allergic reaction**, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
- **Do not use** this product if you have ever had an allergic reaction to any pain reliever/fever reducer, unless you have time for a full night's sleep; in children under 12 years or with any other product containing diphenhydramine.

■ **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.

■ **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

■ **When using this product** drowsiness will occur. Avoid alcoholic drinks. Do not drive a motor vehicle or operate machinery. Take with food or milk if stomach upset occurs

**Directions** ■ **do not take more than directed**

- adults and children 12 years and over: take 2 capsules at bedtime
- do not take more than 2 capsules in 24 hours

Each capsule contains: **potassium 20 mg** Store at 20-25°C (68-77°F) Avoid excessive heat above 40°C (104°F). Protect from light.

**Questions or comments?**  
 Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL  
 For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
 Dist. by: Haleon, Warren, NJ 07059  
 ©2023 Haleon or licensor. Trademarks owned or licensed by Haleon.  
 LIQUI·GELS is a trademark or registered trademark of Catalent Pharma Solutions.

3/16" X 3/16" DATA MATRIX CODE (ECC 200), PRINTED IN BLACK OR DARK BLUE ON UNPRINTED WHITE BACKGROUND.

208522

**208522**

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	6 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	.5 pt

TOP LAYER FACE

UNVARNISHED & PRINTED AREA FOR LOT & EXP. (LINES DO NOT PRINT)

TOP LAYER BACK

<p><b>Drug Facts (continued)</b></p> <p><b>Warnings</b>  <b>Allergy alert:</b> Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:          ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock          ■ skin reddening ■ rash ■ blisters          If an allergic reaction occurs, stop use and seek medical help right away.  <b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you          ■ are age 60 or older          ■ have had stomach ulcers or bleeding problems          ■ take a blood thinning (anticoagulant) or steroid drug          ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]          ■ have 3 or more alcoholic drinks every day while using this product          ■ take more or for a longer time than directed  <b>Heart attack and stroke warning:</b> NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  <b>Do not use</b>          ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer          ■ unless you have time for a full night's sleep          ■ in children under 12 years of age          ■ right before or after heart surgery          ■ with any other product containing diphenhydramine, even one used on skin          ■ if you have sleeplessness without pain  <b>Ask a doctor before use if</b>          ■ stomach bleeding warning applies to you</p>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ you have problems or serious side effects from taking pain relievers or fever reducers</li> <li>■ you have a history of stomach problems, such as heartburn</li> <li>■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke</li> <li>■ you are taking a diuretic</li> <li>■ you have a breathing problem such as emphysema or chronic bronchitis</li> <li>■ you have glaucoma ■ you have trouble urinating due to an enlarged prostate gland</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b>          ■ taking sedatives or tranquilizers, or any other sleep-aid          ■ under a doctor's care for any continuing medical illness          ■ taking any other antihistamines          ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin          ■ taking any other drug</p> <p><b>When using this product</b>          ■ drowsiness will occur ■ avoid alcoholic drinks          ■ do not drive a motor vehicle or operate machinery          ■ take with food or milk if stomach upset occurs</p> <p><b>Stop use and ask a doctor if</b>          ■ you experience any of the following signs of stomach bleeding:          ■ feel faint ■ vomit blood ■ have bloody or black stools          ■ have stomach pain that does not get better          ■ you have symptoms of heart problems or stroke:          ■ chest pain ■ trouble breathing ■ weakness in one part or side of body          ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days          ■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.          ■ redness or swelling is present in the painful area          ■ any new symptoms appear</p>
---	--

BASE LAYER

<p><b>Drug Facts (continued)</b></p> <p><b>if pregnant or breast-feeding,</b> ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b>          ■ do not take more than directed          ■ adults and children 12 years and over: take 2 capsules at bedtime          ■ do not take more than 2 capsules in 24 hours</p> <p><b>Other information</b>          ■ each capsule contains: potassium 20 mg          ■ read all warnings and directions before use.          ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)          ■ protect from light</p> <p><b>Inactive ingredients</b>          D&amp;C red no. 33, FD&amp;C blue no. 1, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution</p> <p><b>Questions or comments?</b> Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL</p>	<h1 style="font-size: 48pt; margin: 0;">Advil®</h1> <p style="font-size: 10pt; margin: 10px 0 0 0;">For most recent product information, visit <a href="http://www.Advil.com">www.Advil.com</a>          Distributed by: Haleon, Warren, NJ 07059          ©2024 Haleon group of companies or its licensor.          Trademarks are owned by or licensed to the Haleon group of companies.          LIQUI-GELS is a trademark or registered trademark of          Catalent Pharma Solutions.</p>
--	--

208524

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



**50% MORE FREE**

**10 Free LIQUI-GELS®**

**Advil<sup>®</sup> PM LIQUI-GELS<sup>®</sup>**

Solubilized Ibuprofen, 200 mg/  
Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid

Actual Size  
**20/30**  
Liquid Filled Capsules

**HALEON**

**READ AND KEEP CAPTION FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active ingredients (in each capsule)** **Purpose(s)**  
Diphenhydramine hydrochloride 25 mg ... Nighttime sleep-aid  
Solubilized ibuprofen equal to ..... Pain reliever  
(present as the free acid and potassium salt)  
Nonsteroidal anti-inflammatory drug

**Uses** ■ For relief of occasional sleeplessness what is associated with minor aches and pains ■ helps you fall asleep and stay asleep

**Warnings**  
**Alert:** ■ Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.  
Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing)  
■ stomach pain ■ skin redness ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs (including prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or by a longer time than directed

**Drug Facts (continued)**  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ unless you have time for a full night's sleep ■ in children under 12 years of age ■ right before or after heart surgery ■ with any other product containing diphenhydramine, even one used on skin ■ if you have sleeplessness without pain  
**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have a breathing problem such as emphysema or chronic bronchitis ■ you have glaucoma ■ you have trouble urinating due to an enlarged prostate gland  
**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers, or any other sleep-aid ■ under a doctor's care for any continuing medical illness ■ taking any other antihistamines ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product** ■ drowsiness will occur ■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery ■ take with food or milk if stomach upset occurs

**Drug Facts (continued)**  
**Spun use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ mouth/hood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ sleeplessness persists continuously for more than 2 weeks ■ insomnia may be a symptom of a serious underlying medical illness. ■ redness or swelling is present in the painful area  
**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed ■ adults and children 12 years and over: take 2 capsules at bedtime ■ do not take more than 2 capsules in 24 hours

**Other information** ■ each capsule contains potassium ■ 20 mg ■ read all other directions before use:  
Keep out of reach of children. ■ store at 20°-25°C (68°-77°F)

**Drug Facts (continued)**  
■ avoid excessive heat above 40°C (104°F)  
■ protect from light  
**Inactive ingredients** D&C red no. 33, FD&C blue no. 1, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution  
**Questions or comments?**  
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL  
**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**  
For most recent product information, visit [www.Advil.com](http://www.Advil.com)  
Distributed by: Haleson, Warren, NJ 07059  
©2024 Haleson or licensee.  
Patents: [www.products.com](http://www.products.com)  
Liqui-Gels® is a trademark or registered trademark of Haleson Pharma Solutions.  
NPA015500

UNPRINTED AREA  
FOR DATE & LOT  
(UNVARNISHED)

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

**NPA015500**

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**(in each capsule)**

**Purposes**  
Diphenhydramine hydrochloride 25 mg..... Nighttime sleep-aid  
Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*..... Pain reliever (present as the free acid and potassium salt), nonsteroidal anti-inflammatory drug

**Uses**  
■ for relief of occasional sleeplessness when associated with minor aches and pains  
■ helps you fall asleep and stay asleep

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing)  
■ shock ■ skin redness ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product.  
■ take more or for a longer time than directed

**Drug Facts (continued)**  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ unless you have time for a full night's sleep  
■ in children under 12 years of age  
■ right before or after heart surgery  
■ with any other product containing diphenhydramine, even one used on skin  
■ if you have sleeplessness without pain

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you  
■ you have problems or serious side effects from taking pain relievers or fever reducers  
■ you have a history of stomach problems, such as heartburn, high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
■ you are taking a diuretic  
■ you have a breathing problem such as emphysema or chronic bronchitis  
■ you have glaucoma  
■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**  
■ taking sedatives or tranquilizers, or any other sleep-aid  
■ under a doctor's care for any continuing medical illness  
■ taking any other antihistamines  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin  
■ taking any other drug

**When using this product** ■ avoid alcoholic drinks  
■ drowsiness will occur  
■ do not drive a motor vehicle or operate machinery  
■ take with food or milk if stomach upset occurs

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

For most recent product information, visit [www.Advii.com](http://www.Advii.com)  
©2024 HALEON or licensor  
Trademarks owned or licensed by HALEON.  
Patents: [www.productpatents.com](http://www.productpatents.com)  
LQUV-6EL5@ is a trademark or registered trademark of Catalent Pharma Solutions.

**Questions or comments?**  
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

**Drug Facts (continued)**

UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

**Drug Facts (continued)**

**Side use and ask a doctor if**  
■ you experience any of the following signs of stomach bleeding:  
■ feel faint ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ have symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing  
■ slowed speech ■ leg swelling  
■ pain gets worse or lasts more than 10 days  
■ sleeplessness persists continuously for more than 2 weeks; insomnia may be a symptom of a serious underlying medical illness.  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery. In case of overdose, get medical help or contact a Poison Control Center right away.

**Keep out of reach of children.**

**Directions** ■ **do not take more than directed**  
■ adults and children 12 years and over: take 2 capsules at bedtime  
■ do not take more than 2 capsules in 24 hours

**Other information**  
■ each capsule contains: potassium 20 mg  
■ read all warnings and directions before use. Keep carton.  
■ store at 20-25°C (68-77°F)  
■ avoid excessive heat above 40°C (104°F)  
■ protect from light.

**Inactive ingredients** D&C red no. 33, FD&C blue no. 1, gelatin, medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution



**Advil<sup>®</sup> PM<sup>®</sup>**  
**LIQUI-GELS<sup>®</sup>**  
Solubilized Ibuprofen, 200 mg /  
Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)/Nighttime Sleep-Aid

**40**  
Liquid Filled Capsules

HALEON

**NPA015501**

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredients (in each capsule)**      **Purposes**

Diphenhydramine  
hydrochloride 25 mg..... Nighttime sleep-aid  
Solubilized ibuprofen equal to  
200 mg ibuprofen (NSAID) \*..... Pain reliever  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- have had stomach ulcers or bleeding problems
- are age 60 or older
- have had stomach (anticoagulant) or steroid drug
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other antihistamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

**When using this product**

- avoid alcoholic drinks
- drowsiness will occur
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

**Drug Facts (continued)**

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

For most recent product information, visit [www.Advii.com](http://www.Advii.com)  
Distributed by: Haleon, Warren, NJ 07059  
©2024 Haleon or licensor  
Patents: [www.producpats.com](http://www.producpats.com)  
Liqui-GELs® is a trademark or registered trademark of Catalent Pharma Solutions.

Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

**Questions or comments?**

UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

NPA015495

**25% MORE FREE**

**10 Free LIQUI-GELS®**

**Advil PM LIQUI-GELS®**

Solubilized Ibuprofen, 200 mg / Diphenhydramine HCl, 25 mg  
Pain Reliever (NSAID)/Nighttime Sleep-Aid

Actual Size **40 50** Liquid Filled Capsules

HALEON

**NPA015495**

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



UNPRINTED AREA FOR DATE & LOT CODING (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

**NPA015496**

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



UNPRINTED AREA FOR DATE & LOT CODING (UNVARNISHED)

PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)

**NPA015498**

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
09/09/2024 05:16:48 PM