



FRONT

000071099

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	N/A

BACK



AREA FOR LOT AND EXP  
TO BE PRINTED IN DARK COLOR  
26 mm x 5.5 mm

SEAL AREA FOR LOT & EXP  
25 mm x 4.5 mm

000068372

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	N/A

**Advil PM**

**Drug Facts** (continued)

- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ **do not take more than directed**

- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours

**Other information**

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

**Inactive ingredients** calcium stearate, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue no. 2 aluminum lake, glyceryl dibehenate, hypromellose, lactose monohydrate, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** Call weekdays 9 AM to 5 PM EST at **1-800-88-ADVIL**

For most recent product information, visit [www.Advils.com](http://www.Advils.com).  
Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
©2021 GSK group of companies or its licensor. Trademarks are owned by or licensed to the GSK group of companies. Patents: [www.productpats.com](http://www.productpats.com) 000071082

**Advil PM**

Ibuprofen, 200 mg /  
Diphenhydramine citrate, 38 mg  
Pain Reliever (NSAID) /  
Nighttime Sleep-Aid

**Two 2 Count Packets**

**4 Coated Caplets\***  
\*Capsule Shaped Tablets

**Two 2ct PACKETS**

Do Not Use if pouch is broken or torn.

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredients (in each caplet)**

Diphenhydramine citrate 38 mg..... Nighttime sleep-aid  
Ibuprofen 200 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Purposes**

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Drug Facts** (continued)

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other antihistamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better

**FPO**

3 0573 0164 14 6

LOT CODE & EXP. DATE

000071082

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**Drug Facts (continued)**

**Uses**  
 ■ for relief of occasional sleeplessness when associated with minor aches and pains ■ helps you fall asleep and stay asleep


**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ unless you have time for a full night's sleep ■ in children under 12 years of age ■ right before or after heart surgery ■ with any other product containing diphenhydramine, even one used on skin ■ if you have sleeplessness without pain

**Ask a doctor before use if**  
 ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have a breathing problem such as emphysema or chronic bronchitis ■ you have glaucoma ■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**  
 ■ taking sedatives or tranquilizers, or any other sleep-aid ■ under a doctor's care for any continuing medical illness ■ taking any other antihistamines ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

**When using this product**  
 ■ drowsiness will occur ■ avoid alcoholic drinks ■ do not drive a motor vehicle or operate machinery ■ take with food or milk if stomach upset occurs



**Drug Facts (continued)**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not take more than directed ■ adults and children 12 years and over: take 2 caplets at bedtime ■ do not take more than 2 caplets in 24 hours

**Other information**  
 ■ read all warnings and directions before use. ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
 calcium stearate, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue no. 2, aluminum lake, glyceryl dibehenate, hypromellose, lactose monohydrate, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?**  
 Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

For most recent product information, visit [www.Advil.com](http://www.Advil.com).  
 Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensor  
 Trademarks owned or licensed by GSK  
 Patents: [www.productpats.com](http://www.productpats.com)

000068377

**81% FPO**

3 0573 0164 09 2

DO NOT USE IF SEAL AROUND CAP IS BROKEN OR MISSING. ADVIL SAFETY SEALED.

**Advil PM**  
 Ibuprofen, 200 mg / Diphenhydramine citrate, 38 mg  
 Pain Reliever (NSAID) / Nighttime Sleep-Aid

**8 Coated Caplets\***  
 \*Capsule-Shaped Tablets

**Drug Facts**

Active ingredients (in each caplet)	Purposes
Diphenhydramine citrate 38 mg	Nighttime sleep-aid
Ibuprofen 200 mg (NSAID)*	Pain reliever
*nonsteroidal anti-inflammatory drug	

LIFT HERE for More Drug Facts

1.125" X 0.4"  
 UNVARNISHED &  
 UNPRINTED AREA  
 FOR LOT & EXP  
 (LINES DO NOT PRINT)

ADVIL SAFETY SEALED ADVIL SAFETY SEALED ADVIL SAFETY SEALED

TEAR HERE TO OPEN

**Drug Facts (continued)**

**Stop use and ask a doctor if**  
 ■ you experience any of the following signs of stomach bleeding:  
 ■ feel faint ■ vomit blood  
 ■ have bloody or black stools  
 ■ have stomach pain that does not get better  
 ■ you have symptoms of heart problems or stroke:  
 ■ chest pain ■ trouble breathing  
 ■ weakness in one part or side of body  
 ■ slurred speech  
 ■ leg swelling  
 ■ pain gets worse or lasts more than 10 days  
 ■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

000068377

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**Advil PM**

**Travel Size**

**Advil PM**

Ibuprofen, 200 mg /  
Diphenhydramine citrate, 38 mg  
Pain Reliever (NSAID) /  
Nighttime Sleep-Aid

Six 2 Count Packets

12  
Coated Caplets\*  
\*Capsule-Shaped Tablets

**Drug Facts (continued)**

- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours

**Other information**

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F)

**Inactive ingredients** calcium stearate, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue no. 2 aluminum lake, glyceryl dibehenate, hypromellose, lactose monohydrate, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?** Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
©2021 GSK group of companies or its licensor. Trademarks are owned by or licensed to the GSK group of companies. Patents: [www.productpats.com](http://www.productpats.com)  
For most recent product information, visit [www.Advil.com](http://www.Advil.com).

**Do Not Use if pouch is broken or torn.**

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredients (in each caplet)**

Diphenhydramine citrate 38 mg..... Nighttime sleep-aid  
Ibuprofen 200 mg (NSAID)\* ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Purposes**

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Drug Facts (continued)**

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer unless you have time for a full night's sleep
- in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other antihistamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- drowsiness will occur
- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better

000071091

FPO

3 0573 0164 12 2

000071091

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

**UNPRINTED AREA FOR DATE & LOT (UNVARNISHED)**

**PRINTED AREA FOR DATE & LOT LASER CODING (VARNISHED)**

Patents: [www.productipats.com](http://www.productipats.com)  
000068375

3 0573 0164 20 7

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active ingredients (in each caplet)** **Purposes**  
Diphenhydramine citrate 38 mg..... Nighttime sleep-aid  
Ibuprofen 200 mg (NSAID)\*..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ For relief of occasional sleeplessness when associated with minor aches and pains ■ helps you fall asleep and stay asleep

**Warnings**  
**Major alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma wheezing ■ shock ■ skin redness ■ rash ■ blisters

**Skin reactions:** ■ If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Drug Facts (continued)**  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risks are higher if you use more than directed or for longer than directed.

**Do not use** ■ If you have ever had an allergic reaction to any one pain reliever never reduce ■ unless you have one for a full night's sleep ■ in children under 12 years of age ■ right before or after heart surgery ■ with any other product ■ if you have sleeplessness without pain

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have a breathing problem such as emphysema or chronic bronchitis ■ you have glaucoma ■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers, or any other sleep-aid ■ under a doctor's care for any continuing medical illness ■ taking any other antihistamines ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

**When using this product** ■ avoid alcoholic drinks ■ drowsiness will occur

**Drug Facts (continued)**  
■ avoid excessive heat above 40°C (104°F)

**Inactive ingredients** calcium stearate, camellia wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue no. 2 aluminum lake, glyceryl dibehenate, hypromellose, lactose monohydrate, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

**Questions or comments?**  
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

For most recent product information, visit [www.AdvIL.com](http://www.AdvIL.com).  
Distributed by GSK Consumer Healthcare, Warren, NJ 07059  
©2021 GSK or licensee. Trademarks owned or licensed by GSK.

**Drug Facts (continued)**  
■ do not drive a motor vehicle or operate machinery ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ you have stomach pain that does not get better

■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days

■ sleeplessness persists continuously for more than 2 weeks.

■ insomnia may be a symptom of a serious underlying medical illness. ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed ■ adults and children 12 years and over: take 2 caplets at bedtime ■ do not take more than 2 caplets in 24 hours

**Other information** ■ read all warnings and directions before use. Keep carton. ■ store at 20-25°C (68-77°F)

# Advil PM

Ibuprofen, 200 mg/  
Diphenhydramine citrate, 38 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid



000068375

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active ingredients (in each caplet)** **Purposes**  
Ibuprofen 200 mg (NSAID)\*..... Nighttime sleep-aid  
Diphenhydramine citrate 38 mg..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ For relief of occasional sleeplessness when associated with minor aches and pains ■ Helps you fall asleep and stay asleep

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

000071092

3 0573 0164 21 4

FPO

**Drug Facts (continued)**  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ If you have ever had an allergic reaction to any other pain reliever/fever reducer ■ unless you have time for a full night's sleep ■ in children under 12 years of age ■ right before or after heart surgery ■ with any other product containing diphenhydramine, even one used on skin ■ if you have sleeplessness without pain

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver or kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have a breathing problem such as emphysema or chronic bronchitis ■ you have glaucoma ■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** ■ taking sedatives or tranquilizers, or any other sleep-aid ■ under a doctor's care for any continuing medical illness ■ taking any other antihistamines ■ taking aspirin or heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ taking any other drug

**When using this product** ■ avoid alcoholic drinks ■ drowsiness will occur ■ do not drive a motor vehicle or operate machinery

**Drug Facts (continued)**  
**Stomach upset occurs** ■ Take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs or symptoms: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. ■ redness or swelling is present in the painful area ■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless specifically directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ **do not take more than directed** ■ adults and children 12 years and over, take 2 caplets at bedtime ■ do not take more than 2 caplets in 24 hours

**Other information** ■ read all warnings and directions before use. Keep carton. ■ store at 20-25°C (68-77°F)

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

For most recent product information, visit [www.Advii.com](http://www.Advii.com)  
©2021 GSK group of companies or its licensor  
Distributed by GSK Consumer Healthcare, Merck, NJ 07099

Patents: [www.products.com](http://www.products.com)  
Trademarks are owned by or licensed to the GSK group of companies

Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

**Questions or comments?**  
starch glycolate, stearic acid, titanium dioxide  
glycol, pregelatinized starch, sodium lauryl sulfate, sodium cellulose, pharmaceutical ink, polydextrose, polyethylene glycol, hypromellose, lactose monohydrate, microcrystalline cellulose, PEG 400, croscarmellose sodium, FD&C blue no. 2 aluminum lake, glyceryl dibehenate, colloidal silicon dioxide, corn starch, croscarmellose sodium, calcium stearate, carmellose sodium, calcium stearate, camellia wax

**Inactive ingredients**

**Drug Facts (continued)**  
■ avoid excessive heat above 40°C (104°F)

**50% MORE FREE**

**Advil PM**

Ibuprofen, 200 mg /  
Diphenhydramine citrate, 38 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid


**20 30**  
Coated Caplets\*  
\*Capsule-Shaped Tablets

000071092

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

000071084



3 0573 0164 30 6

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**  
**Active ingredients (in each caplet)** **Purposes**  
Ibuprofen 200 mg (NSAID).....Nighttime sleep-aid  
Diphenhydramine citrate 38 mg.....Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ for relief of occasional sleeplessness when associated with or acts and pains ■ helps you fall asleep and stay asleep

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing aspirin or steroid drug ■ take other drugs containing naproxen, rofecoxib, or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed


**Drug Facts (continued)**  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ unless you have been told by a health care provider ■ in children under 12 years of age ■ right before or after heart surgery ■ with any other product containing diphenhydramine, even one used on skin ■ if you have sleeplessness without pain.  
**Ask a doctor before use if:** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a breathing problem such as emphysema or chronic bronchitis ■ you have glaucoma ■ you have trouble urinating due to an enlarged prostate gland.  
**Ask a doctor or pharmacist before use if you are:** ■ taking medicines or herbal products, or any other sleep-aid ■ taking a doctor's care for any continuing medical illness ■ taking any other antihistamines ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug.  
**When using this product:** ■ drowsiness will occur ■ avoid alcohol drinks ■ do not drive a motor vehicle or operate machinery ■ take with food or milk if stomach upset occurs

**Drug Facts (continued)**  
**Stop use and ask a doctor if:** ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: chest pain in one part of chest ■ cold sweating ■ shortness in one part of chest ■ dizziness ■ blurred speech ■ fast breathing 10 days ■ painless sores or blisters on tongue ■ you have had 2 weeks or longer of a simple cold or sore throat ■ you have a simple cold or sore throat or swelling is present in the painful area ■ any new symptoms appear.  
**Important safety information:** Ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless directly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not take more than directed ■ adults and children 12 years and over: take 2 caplets at bedtime ■ do not take more than 2 caplets in 24 hours

**Other information** ■ read all warnings and directions before use. Keep carton. ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

**Advil PM**  
Ibuprofen, 200 mg /  
Diphenhydramine citrate, 38 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid



**40**  
Coated Caplets\*  
\*Capsule-Shaped Tablets

**Drug Facts (continued)**  
**Inactive ingredients**  
corn starch, croscarmellose sodium, FD&C blue no. 2, calcium stearate, carnauba wax, colloidal silicon dioxide, starch glycolate, stearic acid, titanium dioxide

**Questions or comments?**  
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL  
For most recent product information, visit [www.Advill.com](http://www.Advill.com).  
Patients: [www.productcaps.com](http://www.productcaps.com)

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.**

000071084

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

# Advil PM

Ibuprofen, 200 mg /  
Diphenhydramine citrate, 38 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid

**50**  
Coated Caplets\*  
\*Capsule-Shaped Tablets

**Drug Facts (continued)**

**Warnings** **Alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  hives  facial swelling  asthma (wheezing)  shock  skin redness/itching  rash  blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you  are age 60 or older  have had stomach ulcers or bleeding problems  take a blood thinning (anticoagulant) or steroid drug  take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  have 3 or more alcoholic drinks every day while using this product  take more or for a longer time than directed

**When using this product:**  avoid alcoholic drinks  do not drive a motor vehicle or operate machinery  take with food or milk if stomach upset occurs

**Drug Facts (continued)**

**Stop use and ask a doctor if:**  you experience any of the following signs of stomach bleeding:  feel faint  vomit blood  have bloody or black stools  have stomach pain that does not get better  you have symptoms of heart problems or stroke: trouble breathing weakness in one part or side of body slurred speech  big swelling pain gets worse or lasts more than 10 days  sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.  redness or swelling is present in the painful area  any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  do not take more than directed  adults and children 12 years and over: take 2 caplets at bedtime  do not take more than 2 caplets in 24 hours

**Other information**  read all warnings and directions before use. Keep carton.  store at 20-25°C (68-77°F)  avoid excessive heat above 40°C (104°F)

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use:**  if you have ever had an allergic reaction to any other pain reliever/fever reducer  unless you have first had a full night's sleep  in children under 12 years of age  right before or after heart surgery  with any other product containing diphenhydramine, even one used or soon to be used for sleeplessness without pain relief

**Ask a doctor before use if:**  stomach bleeding warning applies to you  you have problems or serious side effects from taking pain relievers or fever reducers  you have a history of stomach problems, such as heartburn  you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  you are taking a diuretic  you have a breathing problem such as emphysema or chronic bronchitis  you have glaucoma  you have a urinary tract or enlarged prostate gland

**Ask a doctor or pharmacist before use if you are:**  taking sedatives or tranquilizers, or any other sleep-aid  under a doctor's care for any continuing medical illness  taking any other antihistamines  taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  taking any other drug

**When using this product:**  drowsiness will occur  avoid alcoholic drinks  do not drive a motor vehicle or operate machinery  take with food or milk if stomach upset occurs

**Drug Facts (continued)**

**Active ingredients:** calcium stearate, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue no. 2, aluminum lake, glyceryl dibehenate, hypromellose, lactose monohydrate, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, potassium sulfate, sodium starch glycolate, stearyl acid, titanium dioxide

**Questions or comments?**  
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

**Do Not Use if seal under bottle cap imprinted with "SCALED FOR YOUR PROTECTION" is broken or missing.**

Distributed by GSK Consumer Healthcare, Warren, NJ 07069  
©2021 GSK group of companies or its licensor. Trademarks are owned by or licensed to the GSK group of companies.

UNPRINTED AREA  
FOR DATE & LOT  
(UNVARNISHED)

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

000071085

FPO

3 0573 0164 33 7

For most recent product information, visit [www.Advil.com](http://www.Advil.com).  
Patients: [www.producingtablets.com](http://www.producingtablets.com)

000071085

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredients (In each caplet)**

Diphenhydramine citrate 38 mg  
Ibuprofen, 200 mg (NSAID)  
\*nonsteroidal anti-inflammatory drug

**Purposes**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Uses**

- for relief of occasional sleeplessness when associated with minor aches and pains
- helps you fall asleep and stay asleep

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- swelling of the face, lips, tongue, or throat
- skin redness
- itching
- difficulty breathing
- stomach pain
- diarrhea
- nausea
- vomiting
- stomach bleeding or black stools
- blood in stool
- blood in urine
- blood in vomit
- black or tarry stools
- bleeding or bruising
- unusually easy bruising or bleeding
- unusually heavy menstrual bleeding
- unusually heavy nosebleeds
- unusually heavy bleeding from cuts or scrapes
- unusually heavy bleeding from gums
- unusually heavy bleeding from nose
- unusually heavy bleeding from mouth
- unusually heavy bleeding from skin
- unusually heavy bleeding from wounds
- unusually heavy bleeding from surgery
- unusually heavy bleeding from dental work
- unusually heavy bleeding from any other source

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning medication or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other amphetamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

**When using this product**

- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

**Drug Facts (continued)**

**Sup use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - red pain
  - vomiting blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part of side of body
  - slurred speech
  - eg swelling
- your pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks
- insomnia may be a symptom of a serious underlying medical illness:
  - redness or swelling is present in the painful area
  - any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours

**Other information**

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients:** calcium stearate, cannaflax wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue no. 2 aluminum lake, glyceryl dibehenate, hypromellose, lactose monohydrate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, povidone, polyvinyl alcohol, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

**Advil PM**

Ibuprofen, 200 mg/  
Diphenhydramine citrate, 38 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid

**80**  
Coated Caplets\*  
\*Capsule-Shaped Tablets

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- unless you have time for a full night's sleep in children under 12 years of age
- right before or after heart surgery
- with any other product containing diphenhydramine, even one used on skin
- if you have sleeplessness without pain

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have glaucoma
- you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers, or any other sleep-aid
- under a doctor's care for any continuing medical illness
- taking any other amphetamines
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

**When using this product**

- avoid alcoholic drinks
- do not drive a motor vehicle or operate machinery
- take with food or milk if stomach upset occurs

**Drug Facts (continued)**

**Sup use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - red pain
  - vomiting blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part of side of body
  - slurred speech
  - eg swelling
- your pain gets worse or lasts more than 10 days
- sleeplessness persists continuously for more than 2 weeks
- insomnia may be a symptom of a serious underlying medical illness:
  - redness or swelling is present in the painful area
  - any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults and children 12 years and over: take 2 caplets at bedtime
- do not take more than 2 caplets in 24 hours

**Other information**

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients:** calcium stearate, cannaflax wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue no. 2 aluminum lake, glyceryl dibehenate, hypromellose, lactose monohydrate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, povidone, polyvinyl alcohol, pregelatinized starch, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

000071088

3 0573 0164 40 5

000071088

FPO

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

UNPRINTED AREA  
FOR DATE & LOT  
(UNVARNISHED)

000071088

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



TOP  
PANEL 1

FOR LOT & EXPIRATION AREA,  
UNVARNISHED AND PRINTED  
IN BLACK OR DARK BLUE.

BASE  
PANEL 3

RELEASED BACK  
PANEL 2

000068376

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP  
PANEL 1

3/16" X 3/16" DATA MATRIX CODE (ECC 200), PRINTED IN BLACK OR DARK BLUE ON UNPRINTED WHITE BACKGROUND.

FOR LOT & EXPIRATION AREA. UNVARNISHED AND PRINTED IN BLACK OR DARK BLUE.

BASE  
PANEL 3

RELEASED BACK  
PANEL 2

000068376

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



**Advil PM**  
Ibuprofen, 200 mg /  
Diphenhydramine citrate, 38 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid

120  
Coated Caplets\*  
Capsule-Shaped Tablets

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION**

**Drug Facts**

**Active ingredients (in each caplet)**  
Diphenhydramine citrate 38 mg, ..... Nighttime sleep-aid  
Ibuprofen 200 mg (NSAID)\*, ..... Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses**  
■ for relief of occasional sleeplessness when associated with minor aches and pains  
■ helps you fall asleep and stay asleep

**Warnings**  
**Alert: Heart:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
■ skin reddening ■ rash  
■ an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**  
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ unless you have time for a full night's sleep  
■ in children under 12 years of age  
■ right before or after heart surgery

PRINTED AREA FOR DATE & LOT  
LASER CODING (VARNISHED)

3 0573 0164 43 6



FPO  
000071089

For US Patent or Application status see [www.productcapts.com](http://www.productcapts.com)

**Drug Facts (continued)**

■ with any other product containing diphenhydramine, even one used on skin  
■ if you have sleeplessness without pain

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you  
■ you have problems or serious side effects from taking pain relievers or fever reducers  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver/cirrhosis, kidney disease, asthma, or had a stroke  
■ you are taking a diuretic  
■ you have a breathing problem such as emphysema or chronic bronchitis  
■ you have glaucoma  
■ you have trouble urinating due to an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**  
■ taking sedatives or tranquilizers, or any other sleep-aid  
■ under a doctor's care for any continuing medical illness  
■ taking any other antihistamines  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
■ taking any other drug

**When using this product:**  
■ drowsiness will occur ■ avoid alcoholic drinks  
■ do not drive a motor vehicle or operate machinery  
■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**  
■ you experience any of the following signs of stomach bleeding:  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing  
■ weakness in one part or side of body  
■ slurred speech  
■ leg swelling  
■ pain gets worse or lasts more than 10 days

**Drug Facts (continued)**

■ sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not take more than directed  
■ adults and children 12 years and over, take 2 caplets at bedtime  
■ do not take more than 2 caplets in 24 hours

**Other information**  
■ read all warnings and directions before use. Keep carton.  
■ store at 20-25°C (68-77°F)  
■ avoid excessive heat above 40°C (104°F)

**Inactive ingredients:** calcium stearate, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue no. 2 aluminum lake, glyceryl dibehenate, hypromellose, lactose monohydrate, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyethylene glycol, pregelatinized starch, sodium lauryl sulfate, sodium stearoyl glycolate, stearic acid, titanium dioxide

**Questions or comments?**  
Call weekdays 9 AM to 5 PM EST at 1-800-88-ADVIL

**Do Not Use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing**

For most recent product information, visit [www.advil.com](http://www.advil.com).  
Distributed by: SSK Consumer Healthcare, Warren, NJ 07059  
©2017 GSK Group of companies or its licensor  
Trademarks are owned by or licensed to the GSK group of companies

128 HD CODE

128 HD CODE

**Advil PM**  
Ibuprofen, 200 mg /  
Diphenhydramine citrate, 38 mg  
Pain Reliever (NSAID)/  
Nighttime Sleep-Aid

120  
Coated Caplets\*  
\*Capsule-Shaped Tablets

128 HD CODE

000071089

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces





-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

VALERIE S PRATT  
01/10/2022 04:55:14 PM