

**Drug Facts** (continued)  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.  
**Directions**  
 • do not take more than directed  
 • adults:  
 • take 2 capsules with a glass of water  
 • if symptoms persist or worsen, ask your doctor  
 • do not take more than 2 capsules in 24 hours, unless directed by a doctor  
 • do not take longer than 10 days, unless directed by a doctor (see Warnings)  
 • under 18 years of age: ask a doctor  
**Other information**  
 • read all warnings and directions before use. Keep carton  
 • store at 20-25°C (68-77°F)

**Drug Facts** (continued)  
 • avoid high humidity and excessive heat above 40°C (104°F)  
 • do not use if Tamper-Evident seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing  
**Inactive ingredients**  
 gelatin, mannitol, pharmaceutical ink, polyethylene glycol, povidone, purified water, sorbitan, sorbitol, vitamin E polyethylene glycol succinate and may contain FD&C Blue #1.  
**Questions or Comments?**  
 call toll free 1-800-447-1140

**Drug Facts**  
**Active ingredient (in each capsule)** Purpose  
 Ibuprofen 200 mg (NSAID)\* .....Pain reliever  
 \*nonsteroidal anti-inflammatory drug  
**Use** • treats migraine  
**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 • hives • facial swelling • asthma (wheezing)  
 • shock • skin reddening • rash • blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:  
 • are age 60 or older  
 • have had stomach ulcers or bleeding problems

• take a blood thinning (anticoagulant) or steroid drug  
 • take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]  
 • have 3 or more alcoholic drinks every day while using this product  
 • take more or for a longer time than directed  
**Do not use**  
 • if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 • right before or after heart surgery  
**Ask a doctor before use if you have**  
 • never had migraines diagnosed by a health professional  
 • a headache that is different from your usual migraines  
 • the worst headache of your life  
 • fever and stiff neck  
 • headaches beginning after or caused by head injury, exertion, coughing or bending  
 • experienced your first headache after the age of 50  
 • daily headaches  
 • a migraine so severe as to require bed rest

**Drug Facts** (continued)  
 • problems or serious side effects from taking pain relievers or fever reducers  
 • stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain  
 • ulcers  
 • bleeding problems  
 • high blood pressure  
 • heart or kidney disease  
 • taken a diuretic  
 • reached age 60 or older  
**Ask a doctor or pharmacist before use if you are**  
 • taking any other drug containing an NSAID (prescription or nonprescription)  
 • taking a blood thinning (anticoagulant) or steroid drug  
 • under a doctor's care for any serious condition  
 • taking any other drug  
**When using this product**  
 • take with food or milk if stomach upset occurs

**Drug Facts** (continued)  
 • long term continuous use may increase the risk of heart attack or stroke  
**Stop use and ask a doctor if**  
 • migraine pain is not relieved or gets worse after first dose  
 • you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.  
 • pain gets worse or lasts more than 10 days  
 • stomach pain or upset gets worse or lasts  
 • any new symptoms appear  
**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

NDC 010888-0000-00  
**Ibuprofen Capsules, 200 mg**  
 Pain Reliever(NSAID)  
 40 Softgels\*  
 \*liquid filled capsules  
 KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

**IP**

**DIE 0437C**  
 9.4375 x 8.59375 Flat  
 2.03125 x 2.03125 x 3.875

**Pharmalabel**  
 7915 INDUSTRIAL VILLAGE ROAD, GREENSBORO, NC 27409  
 PHONE 336-855-7142 • FAX 336-854-3753  
 www.pharmalabel.com

Customer: BANNER  
 Product Number: MIGRAINE 200MG 40CT  
 Revision Level: REV 11/05  
 Size: DIE 0437C  
 Artist: JLS PP# 25528BANN

Proof Date: 11/15/05 Q.C.  
 Proofread: JLS Artist  
 1st Proof  2nd Proof  3rd Proof  4th Proof  Other  
 Revise and resubmit proof

PHARMALABEL provides this proof to represent the image prepared for you according to the instructions on the PHARMALABEL. PHARMALABEL is not responsible for errors or omissions unless clearly indicated on this proof.  
 This proof does not accurately represent the colors that will print in the finished product. Any colors that are referenced on this proof are representative only. The finished product will be printed to match the referenced Pantone Color or any approved match standard provided by the customer.

Approved By: \_\_\_\_\_  
 Date: \_\_\_\_\_

BLACK  
 VARNISH

**DIE 0437C**  
9.4375 x 8.59375 Flat  
2.03125 x 2.03125 x 3.875

NDC 010888-0000-00

**See new warnings information**

# Migraine Relief

## Ibuprofen Capsules, 200 mg

### Pain Reliever (NSAID)

40 Softgels\*  
\*liquid filled capsules

**Ibuprofen Capsules, 200 mg**  
Pain Reliever(NSAID)  
40 Softgels\*  
\*liquid filled capsules

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

**Drug Facts** (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed
- adults:
  - take 2 capsules with a glass of water
  - if symptoms persist or worsen, ask your doctor
- do not take more than 2 capsules in 24 hours, unless directed by a doctor
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- under 18 years of age: ask a doctor

**Other information**

- read all warnings and directions before use. Keep carton
- store at 20-25°C (68-77°F)

**Drug Facts** (continued)

- avoid high humidity and excessive heat above 40°C (104°F)
- do not use if Tamper-Evident seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing

**Inactive ingredients**

gelatin, mannitol, pharmaceutical ink, polyethylene glycol, povidone, purified water, sorbitan, sorbitol, vitamin E polyethylene glycol succinate and may contain FD&C Blue #1.

**Questions or Comments?**  
call toll free 1-800-447-1140

**Drug Facts**

**Active ingredient (in each capsule) Purpose**  
Ibuprofen 200 mg (NSAID)\* .....Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Use** • treats migraine

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives • facial swelling • asthma (wheezing)
- shock • skin reddening • rash • blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if you have**

- never had migraines diagnosed by a health professional
- a headache that is different from your usual migraines
- the worst headache of your life
- fever and stiff neck
- headaches beginning after or caused by head injury, exertion, coughing or bending
- experienced your first headache after the age of 50
- daily headaches
- a migraine so severe as to require bed rest

**Drug Facts** (continued)

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- high blood pressure
- heart or kidney disease
- taken a diuretic
- reached age 60 or older

**Ask a doctor or pharmacist before use if you are**

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs


**Drug Facts** (continued)

- long term continuous use may increase the risk of heart attack or stroke

**Stop use and ask a doctor if**


- migraine pain is not relieved or gets worse after first dose
- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- pain gets worse or lasts more than 10 days
- stomach pain or upset gets worse or lasts
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.



KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Rev # 11/05  
Lot No:  
Exp. Date:



Manufactured by: Banner Pharmacaps Inc.  
4125 Premier Drive, High Point, NC 27265



Customer: **BANNER** Proof Date: 11/15/05  
Product Number: **MIGRAINE 200MG 40CT** Proofread: **JLS** Artist: **JLS** Q.C.  
Revision Level: **REV 11/05**  
Size: **DIE 0437C**  
Artist: **JLS** PP# **25528BANN**

1st Proof  2nd Proof  3rd Proof  4th Proof  Other

**Revise and resubmit proof**

PHARMALABEL provides this proof to represent the image prepared for you according to the instructions indicated on this proof. PHARMALABEL is not responsible for errors or omissions unless clearly indicated on this proof. PHARMALABEL represents the colors that will print in the finished product. Any colors that are referenced on this proof are representative only. The finished product will be printed to match the referenced Pantone Color or any approved match standard provided by the customer.

Approved By: \_\_\_\_\_ Date: \_\_\_\_\_

Customer: **BANNER** Product Number: **MIGRAINE 200MG 40CT** Revision Level: **REV 11/05** Size: **DIE 0437C** Artist: **JLS** PP# **25528BANN**

BLACK VARNISH



-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Andrea Segal  
3/13/2006 10:26:52 AM