


NOT OFFICIAL SIZE/DIELINE




Label Size : 1.5" x 4.5"
 Track : 19/02/2017, 22/02/2017



CATEGORY	Art Production DRUG FACTS INFO.		
	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	6 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	39

	Font	Size
Drug Facts	helvetica bold Oblique	10 pt.
Headings	helvetica bold Oblique	8 pt.
Subheadings	helvetica bold condensed	8 pt.
Body Text	helvetica medium condensed	6 pt.
Leading	N/A	0.5 pt
Bullets	Zapf Dingbats	5 pt.
Barline	N/A	2.5 pt.
Hairlines	N/A	0.5 pt.

Carton Size : 3.0628" x 2.1253" x 1.5934"
Track : 19/02/2017, 22/02/2017



NOT OFFICIAL SIZE/DIELINE

NDC 69452-178-26

**Ibuprofen Capsules,
200 mg**
Pain Reliever/Fever Reducer (NSAID)

200 Capsules

Drug Facts

Active ingredient (in each capsule)
Ibuprofen 200 mg (NSAID).....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - minor pain of arthritis
 - headache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- skin swelling
- asthma (wheezing)
- shock
- facial swelling
- rash
- blisters

If an allergic reaction occurs, seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take 4 or more of a long-term medicine

Heart attack warning: NSAIDs, except aspirin, increase the risk of heart attack, stroke, and death. These risks can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Manufactured for: **Briomp Pharma Inc.**
 600 Alexander Road,
 Princeton, NJ 08540
 Rev # 02/17
 Lot No.:
 Exp. Date:
PEEL HERE

Drug Facts (continued)

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part of side of body
 - slurred speech
 - big swollen vein
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless a health professional directs you to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over:
 - take 1 capsule every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 capsule, 2 capsules may be used
 - do not exceed 6 capsules in 24 hours, unless directed by a doctor
 - children under 12 years: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)
- do not use if the cap is broken or the seal is broken or missing
- Your Protection: Is broken or missing
- read all warnings and directions before use. Keep carton.

Inactive ingredients
FD&C Blue #1, gelatin, hypromellose, mannitol, polyethylene glycol, propylene glycol, povidone, purified water, sorbitan, sorbitol, titanium dioxide, vitamin E polyethylene glycol succinate


Questions or comments? call toll free 1-888-236-2466

Label Size : 3" x 7"
Track : 19/02/2017, 22/02/2017

Colors (b) (4)

	Font	Size
Drug Facts	helvetica bold Oblique	10 pt.
Headings	helvetica bold Oblique	8 pt.
Subheadings	helvetica bold condensed	8 pt.
Body Text	helvetica medium condensed	6 pt.
Leading	N/A	0.5 pt
Bullets	Zapf Dingbats	5 pt.
Barline	N/A	2.5 pt.
Hairlines	N/A	0.5 pt.

NOT OFFICIAL SIZE/DIELINE



Font	Size
Drug Facts	10 pt.
Headings	8 pt.
Subheadings	8 pt.
Body Text	6 pt.
Leading	0.5 pt
Bullets	5 pt.
Barline	2.5 pt.
Hairlines	0.5 pt.

Colors



Carton Size : 2.031" x 2.031" x 3.875"
Track : 19/02/2017, 22/02/2017