


# NOT OFFICIAL SIZE/DIELINE



Carton Size : 3.0628" x 2.1253" x 1.5934"  
Track : 19/02/2017, 22/02/2017, 28/05/2021

	Font	Size
Drug Facts	helvetica bold Oblique	10 pt.
Headings	helvetica bold Oblique	8 pt.
Subheadings	helvetica bold condensed	8 pt.
Body Text	helvetica medium condensed	6 pt.
Leading	N/A	0.5 pt
Bullets	Zapf Dingbats	5 pt.
Barline	N/A	2.5 pt.
Hairlines	N/A	0.5 pt.



(b) (4)

# NOT OFFICIAL SIZE/DIELINE

NDC 69452-178-26

**Ibuprofen Capsules,  
200 mg**  
Pain Reliever/Fever Reducer (NSAID)

200 Capsules

**Drug Facts**

**Active ingredient (in each capsule)**  
Ibuprofen 200 mg (NSAID)\*.....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Purposes**

- Temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever
- toothache

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- shock
- facial swelling
- skin reddening
- asthma (wheezing)
- rash
- blisters

If an allergic reaction occurs, seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

Manufactured for: **Briopharma Inc.**  
 600 Alexander Road,  
 Princeton, NJ 08540  
 Rev # 05/21  
 Lot No.:  
 Exp. Date:  
**PEEL HERE**

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part of side of body
  - slurred speech
  - big swollen eye
- gain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding:** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact Poison Control Center right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- store at 20°-25° C (68°-77°F)
- avoid light and excessive heat above 40°C (104°F)
- do not use if the cap is loose, the inner bottle cap is imprinted with "Sealed for Your Protection" is broken or missing
- read all warnings and directions before use. Keep carton.

**Inactive ingredients**  
FD&C Blue #1, gelatin, hypromellose, mannitol, polyethylene glycol, propylene glycol, povidone, purified water, sorbitan, sorbitol, titanium dioxide, vitamin E polyethylene glycol succinate

**Questions or comments?** call toll free 1-888-235-2466

Label Size : 3" x 7"  
Track : 19/02/2017, 22/02/2017, 28/05/2021



	Font	Size
Drug Facts	helvetica bold Oblique	10 pt.
Headings	helvetica bold Oblique	8 pt.
Subheadings	helvetica bold condensed	8 pt.
Body Text	helvetica medium condensed	6 pt.
Leading	N/A	0.5 pt
Bullets	Zapf Dingbats	5 pt.
Barline	N/A	2.5 pt.
Hairlines	N/A	0.5 pt.

NOT OFFICIAL SIZE/DIELINE

(b) (4)

9.4375 x 8.59375 Flat  
2.03125 x 2.03125 x 3.875

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Rev # 05/21  
Lot No.:  
Exp. Date:

3  
694452178263

Manufactured for: **Bionpharma Inc.**  
 600 Alexander Road, Princeton, NJ 08540  
 Manufactured by: **Pathoan Softgels Inc.**  
 Highpoint, NC, 27265 USA

NDC 69452-178-26

## Ibuprofen Capsules, 200 mg

### Pain Reliever/Fever Reducer (NSAID)

200 Capsules

\*Liquid filled capsules  
Softgels\*

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Ibuprofen Capsules, 200 mg  
Pain Reliever/Fever Reducer (NSAID)

200 Capsules

\*Liquid filled capsules  
Softgels\*

**Drug Facts (continued)**

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen at 20 weeks or later of pregnancy. Ibuprofen may cause complications during delivery. It may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ **do not take more than directed** ■ **the smallest effective dose should be used**

- adults and children 12 years and over: ■ take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information** ■ store at 20-25°C (68-77°F); ■ avoid high humidity and excessive heat above 40°C (104°F)

- do not use if Tamper-Evident seal under bottle cap imprinted with Sealed for Your Protection is broken or missing
- read all warnings and directions before use. Keep carton.

**Inactive ingredients** FD&C Blue #1, gelatin, hypromellose, mannitol, polyethylene glycol, propylene glycol, potassium phosphate, sorbitol, sodium tetraborate, titanium dioxide, vitamin E polyethylene glycol succinate

**Questions or comments? call toll free 1-888-235-2466**

**Drug Facts**

**Active ingredient (in each capsule)** Ibuprofen 200 mg (NSAID) .....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Purposes**

- temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis
- toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin redness ■ rash ■ hiccups

**If an allergic reaction occurs, stop use and seek medical help, right away.**

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance of stomach bleeding is greater with longer use, higher doses, or if you have had stomach ulcers or bleeding problems. ■ take a blood thinning (anticoagulant) or steroid drug

- have had stomach ulcers or bleeding problems
- take other drugs combining prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Drug Facts (continued)**

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product** ■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 3 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear



Carton Size : 2.031" x 2.031" x 3.875"  
 Track : 19/02/2017, 22/02/2017, 28/05/2021

	Font	Size
Drug Facts	helvetica bold Oblique	10 pt.
Headings	helvetica bold Oblique	8 pt.
Subheadings	helvetica bold condensed	8 pt.
Body Text	helvetica medium condensed	6 pt.
Leading	N/A	0.5 pt
Bullets	Zapf Dingbats	5 pt.
Barline	N/A	2.5 pt.
Hairlines	N/A	0.5 pt.

-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

VALERIE S PRATT  
10/21/2021 01:19:36 PM