



PMS-186C PMS-116C PMS-301C Black PMS-300C

Sep. 05, 2002
 Revised by I.Horii/Fujisawa Japan

CrossTech-C45422-Proof 1
 Form M03-PS:C1
 Fujisawa-P.O. 21321-Job No. 62922
 3/10/05-jc-Q4.11

Fonts: Helvetica (A)



NDC 0469-3250-10

Mycamine™
(micafungin sodium)
for injection

50 mg/vial

50 mg X 10 Vials

325010



NDC 0469-3250-10

Mycamine™
(micafungin sodium)
for injection
For Intravenous Infusion Only

50 mg/vial

Once reconstituted with 5 mL of 0.9% sodium chloride injection
(without bacteriostatic agent), each mL contains 10 mg/mL.
Requires further dilution prior to infusion. See package insert.

Single Use Vial. Discard unused portion.

Rx Only

50 mg X 10 Vials

325010



Mycamine™
(micafungin sodium)
for injection

50 mg/vial

Made in Japan
Mfd. for:
Fujisawa Healthcare, Inc.
Deerfield, IL 60015-2548

50 mg X 10 Vials



NDC 0469-3250-10

Mycamine™
(micafungin sodium)
for injection
For Intravenous Infusion Only

50 mg/vial

Sterile, Nonpyrogenic
Each vial contains: Micafungin sodium 50 mg;
lactose, NF 200 mg with citric acid and/or sodium
hydroxide for pH adjustment.
Store at 25°C (77°F) excursions 15°-30°C (59°-86°F).
[See USP Controlled Room Temperature] Protect from light.

50 mg X 10 Vials

325010



(01) 2 0304693 25010 5




Mycamine™
(micafungin sodium)
for injection

50 mg/vial



Preservative free. Discard unused portion.

50 mg X 10 Vials



2 0 0 8 P A

PMS-186C PMS-116C PMS-301C PMS-300C Black



2 0 0 8



PMS-186C


PMS-116C

PMS-301C

PMS-300C

PMS-WHITE
BACKGROUND

Black



~~CrossTech~~ B45420-Proof A7
Form M01-PS:B1
Fujisawa-P.O. 21321-Job No. 40927
2/24/05-hc-Q4.11

Fonts: Helvetica (A)
Typesmiths Pi Font

