


TEAR BELOW

CONTENTS: One swab premoistened with 1 mL of a solution of 3.15% (w/w) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. **DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm). Tear open packet and remove swab. Do not unfold swab. Prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds. Allow the prepped area to air dry for 30 seconds. Do not blot or wipe dry. Discard after a single use. **WARNINGS: For external use only. Flammable, keep away from fire or flame.** Do not use with electrocautery procedures. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

LOT

EXP. 

2NB108

NDC 10819-1080-1 REORDER NO. B10800

Prevantics®
(Chlorhexidine Gluconate (3.15%)*
and Isopropyl Alcohol (70%) Swab)
ANTISEPTIC


SWAB

*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

FOR EXTERNAL USE ONLY

- Use for the preparation of the patient's skin prior to an injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only
- Latex free

IMPORTANT: See carton for complete **Drug Facts** and safety information.
One swab, 0.034 FL. oz. (1 mL) Each


(01)10310819000014

†PDI

Professional Disposables, International, Inc.
Orangeburg, NY 10962-1376 USA
1-800-899-0423 www.pdiinc.com

2NB108_02_07_14.pdf

NET CONTENTS: 10 BOXES OF 100 INDIVIDUAL SWABS
REORDER NO. B10800
NDC 10819-1080-1

†Prevantics® Swab

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
 Antiseptic

*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

Drug Facts		Drug Facts (continued)	
Active ingredients	Purpose	Directions	
Chlorhexidine gluconate 3.15% w/v.....	Antiseptic	<ul style="list-style-type: none"> • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm) • tear open packet and remove swab. Do not unfold swab. • prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds. • allow the prepped area to air dry for 30 seconds • do not blot or wipe dry • discard after a single use 	
Use For the preparation of the patient's skin prior to an injection	Warnings	Other information	
For external use only	<ul style="list-style-type: none"> • on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch 	<ul style="list-style-type: none"> • store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only 	
Flammable, keep away from fire or flame	Do not use	Inactive ingredient	
Do not use with electrocautery procedures	<ul style="list-style-type: none"> • on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch 	USP purified water	
When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	Questions? Call 1-800-999-6423, M-F 9:00 am-5:00 pm EST	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.			

Each swab contains 0.034 fl. oz. (1 mL)

Professional Disposables International, Inc.
 Orangeburg, NY 10962-1376 USA
 1-800-999-6423 www.pdfhc.com



(01)30310819000018



8HB10800

8HB10800_02_10_14.pdf


REORDER NO. B11400
NDC 10819-1080-2

Prevantics® Swab

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Antiseptic

* equivalent to 167 mg of Chlorhexidine Gluconate per pouch
Contains: 3000 Individual Swabs, 0.034 fl. oz. (1 mL) Each

<p>Drug Facts</p> <p>Active ingredients Chlorhexidine gluconate 3.15% w/v. Antiseptic Isopropyl alcohol 70% v/v. Antiseptic</p> <p>Use For the preparation of the patient's skin prior to an injection.</p> <p>Warnings For external use only Flammable, keep away from fire or flame Do not use with electrocautery procedures</p> <p>Do not use</p> <ul style="list-style-type: none"> • on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol • for lumbor puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch <p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm) • tear open packet and remove swab. Do not unfold swab. • prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds. • allow the prepped area to air dry for 30 seconds • do not blot or wipe dry • discard after a single use <p>Other information</p> <ul style="list-style-type: none"> • store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only <p>Inactive ingredient USP purified water</p> <p>Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>
--	--



(01)30310819000865

PDI
 Professional Disposables International, Inc.
 Orangeburg, NY 10962-1376 USA
 1-800-999-6423 www.pdihc.com

8HB10800_02_10_14.pdf

TEAR BELOW

CONTENTS: One swab premoistened with 1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. **DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm). Tear open packet and remove swab. Do not unfold swab. Prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin being prepared back and forth 30 seconds for 3 swabs. Allow the prep site to dry for 30 seconds. Do not blow-dry the prep site. Discard after single use. **WARNINGS:** For external use only. Flammable, keep away from fire or flame. Do not use with electrocautery procedures. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

LOT

EXP

2MB114

NDC 10819-1080-2 REORDER NO. B11400

Prevantics®
(Chlorhexidine Gluconate (3.15%)[†] and Isopropyl Alcohol (70%) Swab)
ANTISEPTIC

SWAB

*equivalent to 1.67 mg of Chlorhexidine Gluconate per pouch
FOR EXTERNAL USE ONLY

- Use for the preparation of the patient's skin prior to an injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only
- Latex free

IMPORTANT: See insert for complete **Drug Facts** and safety information.
One Swab, 0.034 fl. oz. (1 mL) Each

PDI

(01)103108190000861

2MB114_02_07_14.pdf

<p>*Prevantics® Swab REORDER NO. B11400 Chlorhexidine Gluconate (3.15%)* NDC 10819-1080-2 and Isopropyl Alcohol (70% Swab) Antiseptic *equivalent to 167 mg of Chlorhexidine Gluconate per pouch</p>	
<p>Drug Facts</p>	
<p>Active ingredients</p> <p>Chlorhexidine gluconate 3.15% w/v..... Antiseptic Isopropyl alcohol 70% v/v..... Antiseptic</p>	<p>Purpose</p> <p>Antiseptic</p>
<p>Use For the preparation of the patient's skin prior to an infection</p>	
<p>Warnings</p> <p>For external use only</p> <p>Flammable, keep away from fire or flame</p> <p>Do not use with electrocautery procedures</p>	
<p>Do not use</p> <ul style="list-style-type: none"> on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol for lumbal puncture or in contact with the meninges on open skin wounds or as a general skin cleanser under occlusive patch 	
<p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p>	
<p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶</p>	

<p>Drug Facts (continued)</p>	
<p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	
<p>Directions</p> <ul style="list-style-type: none"> use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm) tear open packet and remove swab. Do not unfold swab. prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds. allow the prepped area to air dry for 30 seconds. do not blot or wipe dry. discard after a single use 	
<p>Other information</p> <ul style="list-style-type: none"> store between 68-77° F (20-25° C) avoid freezing and excessive heat above 104° F (40° C) for hospital and professional use only 	
<p>Inactive ingredient</p> <p>USP purified water</p>	
<p>Questions? Call 1-800-999-6423, M-F 9:00 am-5:00 pm EST)</p>	
<p>Each Swab contains 0.034 fl. oz. (1 mL)</p>	
<p>PDI Professional Disposables International, Inc. Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdic.com</p>	
<p>5LB1141S</p>	

5LB1141S_02_10_14.pdf

NDC 10819-4078-1 REORDER NO. 540750

† **Prevantics**[®]
[Chlorhexidine Gluconate (3.15%)*
and Isopropyl Alcohol (70%) Swab]
PATIENT PREOPERATIVE SKIN PREPARATION

*equivalent to 248 mg of Chlorhexidine Gluconate per pouch
FOR EXTERNAL USE ONLY • Use for the preparation of the patient's skin prior to surgery or injection
of any kind. Do not use for the preparation of the patient's skin for any other purpose. See
professional use only. **IMPORTANT:** See carton for complete **Drug Facts** and safety information.
One Swabstick, 0.054 fl. oz. (1.6 mL) Each

SWABSTICK
Professional Disposables International, Inc.
1-800-999-8423 www.pdihc.com


† **pdi**

CONTENTS: One swabstick premoistened with 1.6 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. **DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burn. Maximum treatment area for one swabstick is 100 cm² (10.76 in²). Do not use on the face. Do not use on the face of patients under 12 years of age. Do not use to expose the swabstick. Remove the swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in a circular motion. Do not use on the face. Do not use on the face of patients under 12 years of age. Do not use on the face and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one and one half (1.5) minutes. Do not blot or wipe dry. Discard after a single use. **WARNINGS: For external use only. Flammable, keep away from fire or flame.** Do not use with electrocautery procedures. **Keep out of reach of children.**

Center Right-way,
Professional Disposables International, Inc.
1-800-999-8423 www.pdihc.com

LOT

THIS SIDE UP
PEEL APART HERE



(01)10310819000021

2MS407

2MS407_02_07_14.pdf

NET CONTENTS: 10 BOXES OF 50 INDIVIDUAL SWABSTICKS
REORDER NO. S40750
NDC 10819-4075-1

PDI Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation

*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

Drug Facts	
Active ingredients Chlorhexidine gluconate 3.15% w/v Antiseptic Isopropyl alcohol 70% v/v Antiseptic	Purpose Antiseptic Antiseptic
Use For the preparation of the patient's skin prior to surgery or injection	
Warnings For external use only Flammable, keep away from fire or flame Do not use with electrocautery procedures	
Do not use • on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch	
When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Drug Facts (continued)	
Directions • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm) • peel apart the pouch to expose the swabstick • remove the swabstick from package. Do not touch the foam tip. • prior to surgery or injection, apply swabstick to the procedure site • place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute • turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute • allow the prepped area to air dry for one and one half minutes (1.5 minutes) • do not blot or wipe dry • discard after a single use	
Other information • store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only	
Inactive ingredient USP purified water	
Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	

Each swabstick contains 0.054 fl. oz. (1.6 mL)

PDI Professional Disposables International, Inc.
Orangeburg, NY 10962-1376 USA
1-800-999-6423 www.pdiinc.com



8HS40750

8HS40750_02_10_14.pdf


REORDER NO. S32450
NDC 10819-4075-2

Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation

*equivalent to 268 mg of Chlorhexidine Gluconate per pouch
Contains: 500 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each

8LS32450



(01)30310819000872

Drug Facts	Drug Facts (continued)
<p>Active ingredients</p> <p>Chlorhexidine gluconate 3.15% w/v Antiseptic Isopropyl alcohol 70% v/v Antiseptic</p> <p>Use For the preparation of the patient's skin prior to surgery or injection</p> <p>Warnings</p> <p>For external use only Flammable, keep away from fire or flame Do not use with electrocautery procedures</p> <p>Do not use</p> <ul style="list-style-type: none"> • on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol • for lumbal puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch <p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and ask a doctor: if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p>Directions</p> <ul style="list-style-type: none"> • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm) • tear open the pouch to expose the swabstick • remove the swabstick from package. Do not touch the foam tip. • prior to surgery or injection, apply swabstick to the procedure site • place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute • turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute • allow the prepped area to air dry for one and one half minutes (1.5 minutes) • do not blot or wipe dry • discard after a single use <p>Other information</p> <ul style="list-style-type: none"> • store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only <p>Inactive ingredient USP-purified water</p> <p>Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>

†PDI Professional Disposables International, Inc.
Orangeburg, NY 10962-1376 USA
1-800-999-6423 www.pdfic.com

8LS32450_02_10_14.pdf

NDC 10819-4075-2 REORDER NO. 53240
Preventics[®]
 (Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab)
PATIENT PREOPERATIVE SKIN PREPARATION

*equivalent to 3.68 mg of Chlorhexidine Gluconate per patch

FOR EXTERNAL USE ONLY • Use for the preparation of the patient's skin prior to surgery or injection

- Latex free • Helps reduce bacteria that potentially can cause skin infection • For hospital and professional use only • **IMPORTANT:** See insert for complete **Drug Facts** and safety information.

One Swabstick, 0.654 fl.oz. (1.9 mL) Each

SWABSTICK

CONTENTS: One swabstick, premixed with 1.6 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl Alcohol. **Preventics** is a sterile, single-use swabstick. These products may cause irritation or chemical burns. Maximum treatment area for one swabstick is approximately 4 in. x 4 in. (10 cm x 10 cm) or moist site: 2 in. x 5 in. (5 cm x 12.5 cm). Tear open the pouch to expose the swabstick. Turn the swabstick over (unused side of the foam tip) and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (used side of the foam tip) and repeat the procedure by prepping one (1) minute. Allow the skin to air dry for one (1) minute. Do not use on mucous membranes. Do not use with electrocautery procedures. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

Professional: Deposite ea Innovations, Inc.
 Orangeburg, NY 10962, USA • 1-800-899-6423 www.pdhc.com

THIS SIDE UP ↑

TEAR OPEN HERE ↑

LOT

EX3



(01)10310819000078

2MS324

2MS324_02_10_14.pdf

REORDER NO. S32450
 NDC 10819-4075-2

+Prevantics[®] Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
 Patient preoperative skin preparation

*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

Drug Facts		Drug Facts (continued)	
Active ingredients	Purpose	Do not use	
Chlorhexidine gluconate 3.15% w/v	Antiseptic	• on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol	
Isopropyl alcohol 70% v/v	Antiseptic	• for lumbar puncture or in contact with the meninges	
Use	For the preparation of the patient's skin prior to surgery or injection	• on open skin wounds or as a general skin cleanser	
Warnings	For external use only Flammable, keep away from fire or flame Do not use with electrocautery procedures	• under occlusive patch	
		When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	

Drug Facts (continued)		Drug Facts (continued)	
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.		• place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.		• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute	
Directions		• allow the prepped area to air dry for one and one half minutes (1.5 minutes)	
• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.		• do not blot or wipe dry	
• maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)		• discard after a single use	
• tear open the pouch to expose the swabstick		Other information	
• remove the swabstick from package. Do not touch the foam tip.		• store between 68-77°F (20-25°C)	
• prior to surgery or injection, apply swabstick to the procedure site		• avoid freezing and excessive heat above 104°F (40°C)	
		• for hospital and professional use only	
		Inactive ingredient	
		USP purified water	
		Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	

Each swabstick contains 0.054 fl. oz. (1.6 mL)

PDI Professional Disposables International, Inc.
 Orangeburg, NY 10962-1376 USA
 1-800-999-6423 www.pdihc.com

5LS324IS



5LS324IS_02_10_14.pdf

5JS41950

30 Individual Maxi Swabsticks, 0.172 fl. oz. (5.1 mL) Each

PDI

Preventics

MAXI SWABSTICK

30 Individual Maxi Swabsticks, 0.172 fl. oz. (5.1 mL) Each

PDI

30 Individual Maxi Swabsticks, 0.172 fl. oz. (5.1 mL) Each

PDI

Preventics

MAXI SWABSTICK

30 Individual Maxi Swabsticks, 0.172 fl. oz. (5.1 mL) Each

PDI

30 Individual Maxi Swabsticks, 0.172 fl. oz. (5.1 mL) Each

PDI

Preventics

MAXI SWABSTICK

30 Individual Maxi Swabsticks, 0.172 fl. oz. (5.1 mL) Each

PDI

Drug Facts

Indications

• For use on the skin to help prevent the spread of infection.

Warnings

• Do not use if you are allergic to chlorhexidine gluconate or any of the other ingredients.

Directions

• Apply to the skin to be treated.

Other information

• Contains 30 individual Maxi Swabsticks, 0.172 fl. oz. (5.1 mL) Each.

Drug Facts (continued)

Directions

• For use on the skin to help prevent the spread of infection.

Warnings

• Do not use if you are allergic to chlorhexidine gluconate or any of the other ingredients.

Directions

• Apply to the skin to be treated.

Other information

• Contains 30 individual Maxi Swabsticks, 0.172 fl. oz. (5.1 mL) Each.

Drug Facts (continued)

Directions

• For use on the skin to help prevent the spread of infection.

Warnings

• Do not use if you are allergic to chlorhexidine gluconate or any of the other ingredients.

Directions

• Apply to the skin to be treated.

Other information

• Contains 30 individual Maxi Swabsticks, 0.172 fl. oz. (5.1 mL) Each.

NDC 1081-4076-1
EFC0808 NO. 54169

Prevantics®

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
PATIENT PREOPERATIVE SKIN PREPARATION

*equivalent to 85 mg of Chlorhexidine Gluconate per pouch

FOR EXTERNAL USE ONLY

- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only • Latex free

IMPORTANT: See carton for complete *Drug Facts* and safety information.

One Maxi Swabstick, 0.172 fl. oz. (5.1 mL) Each

LOT ↑

THIS SIDE UP
PEEL APART HERE

EX ↑

MAXI SWABSTICK

†PDI

CONTENTS: One Maxi Swabstick premoistened with 5.1 mL of a solution of 3.15%(w/v) Chlorhexidine gluconate, 70%(v/v) Isopropyl alcohol, and USP purified water.

DIRECTIONS: Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one maxi swabstick is approximately: dry site: 7 by 7 inches (18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm). Peel apart the pouch to expose the swabstick. Remove swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one and one half minutes (1.5 minutes). Do not blot or wipe dry. Discard after a single use.

WARNINGS: For external use only. Flammable, keep away from fire or flame. Do not use with electrocautery procedures. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Prevantra, Diprads as Information, Inc.
Changshu, NJ 07021-1278 USA 1-800-939-6423 www.pdi.com
(571) 031-0581 80000338

2MS419

2MS419_02_10_14.pdf

NET CONTENTS: 10 BOXES OF 30 INDIVIDUAL SWABSTICKS
 REORDER NO. S41950
 NDC 10819-4075-4

Prevantics® Maxi Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
 Patient preoperative skin preparation

*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

8HS41950

Drug Facts (continued)	
Drug Facts	Directions
Active Ingredients Chlorhexidine gluconate 3.15% w/v Antiseptic Isopropyl alcohol 70% v/v Antiseptic	<ul style="list-style-type: none"> use with care in premature infants or in infants under 2 months of age. These products may cause irritation or chemical burns. maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm) peel apart the pouch, to expose the swabstick remove the swabstick from package. Do not touch the foam tip. prior to surgery or injection, apply swabstick to the procedure site place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute allow the prepped area to air dry for one and one half minutes (1.5 minutes) do not blot or wipe dry discard after a single use
Use For the preparation of the patient's skin prior to surgery or injection	
Warnings For external use only Flammable, keep away from fire or flame Do not use with electrocautery procedures	
Do not use • on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol • for lumbur puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch	
When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
	Other information • store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only
	Inactive ingredient USP purified water
	Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)

Each Maxi Swabstick contains 0.172 fl. oz. (5.1 mL)

Professional Disposables International, Inc.
 Orangetburg, NY 10962-1376 USA
 1-800-999-6423 www.pdinc.com



(01)30310819000032



8HS41950_02_10_14.pdf

REORDER NO. S27350
 NDC 10819-4075-3

Prevantics® Maxi Swabstick
 [Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
 Patient preoperative skin preparation

*equivalent to 855 mg of Chlorhexidine Gluconate per pouch
Contains: 300 Individual Swabsticks, 0.172 fl. oz. (5.1 mL) Each

Drug Facts

Active ingredients
 Chlorhexidine gluconate 3.15% w/v Antiseptic
 Isopropyl alcohol 70% v/v Antiseptic

Use For the preparation of the patient's skin prior to surgery or injection

Warnings
For external use only
Flammable, keep away from fire or flame
 Do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions


- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swab is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)
- tear open the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply swabstick to the procedure site
- place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute
- turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute
- allow the prepped area to air dry for one and one half minutes (1.5 minutes)
- do not blot or wipe dry
- discard after a single use

Other information


- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient
 USP purified water

Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)



(01)30310819000889



Professional Disposables International, Inc.
 Orangeburg, NY 10962-1376 USA
 1-800-999-6423 www.pdihcc.com

8MS27350_02_10_14.pdf

MEC 10118-4016-3
RE-ORDER NO. 52710

Preventics®
(Chlorhexidine Gluconate (3.15%) * and Isopropyl Alcohol (70%) Swab)
PATIENT PREOPERATIVE SKIN PREPARATION

*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

FOR EXTERNAL USE ONLY

- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only • Latex free

IMPORTANT: See insert for complete *Drug Facts* and safety information.

One Maxi Swabstick, 0.172 fl.oz. (5.1 mL) Each

MAXI SWABSTICK

**THIS SIDE UP
TEAR OPEN HERE**

CONTENTS: One Maxi Swabstick, premoistened with 5.1 mL of a solution of 0.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water.

DIRECTIONS: Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one maxi swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm). Peel apart the pouch to expose the swabstick. Remove swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one and one half minutes (1.5 minutes). Do not blot or wipe dry. Discard after a single use.

WARNINGS: For external use only. Flammable, keep away from fire or flame. Do not use with electrocautery procedures. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Professional Disposables International, Inc.
Orangeburg, NY 10962-1976 USA 1-800-999-6423 www.pdih.com

011011011000088

2NS273

2NS273_02_10_14.pdf

REORDER NO. S27350
 NDC 10819-4075-3

+Prevantics[®] Maxi Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
 Patient preoperative skin preparation

*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

Drug Facts		Drug Facts (continued)	
Active ingredients	Purpose	Do not use	
Chlorhexidine gluconate 3.15% w/v.	Antiseptic	• on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol	
Isopropyl alcohol 70% v/v.	Antiseptic	• for lumbar puncture or in contact with the meninges	
Use For the preparation of the patient's skin prior to surgery or injection		• on open skin wounds or as a general skin cleanser	
Warnings		• under occlusive patch	
For external use only		When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
Flammable, keep away from fire or flame			
Do not use with electrocautery procedures			

Drug Facts (continued)		Drug Facts (continued)	
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.		• place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.		• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute	
Directions		• allow the prepped area to air dry for one and one half minutes (1.5 minutes)	
• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.		• do not blot or wipe dry	
• maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)		• discard after a single use	
• tear open the pouch to expose the swabstick		Other information	
• remove the swabstick from package. Do not touch the foam tip.		• store between 68-77°F (20-25°C)	
• prior to surgery or injection, apply swabstick to the procedure site		• avoid freezing and excessive heat above 104°F (40°C)	
		• for hospital and professional use only	
		Inactive ingredient	
		USP purified water	
		Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	

Each Maxi Swabstick contains 0.172 fl. oz. (5.1 mL)

+PDI Professional Disposables International, Inc.
 Orangeburg, NY 10962-1376 USA
 1-800-999-6423 www.pdihc.com

5LS273IS




Prevantics® Swabstick
 [Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
 Patient preoperative skin preparation

REORDER NO. S42850
 NDC 10819-4075-5

*equivalent to 268 mg of Chlorhexidine Gluconate per pouch
Contains: 500 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each

Drug Facts	Drug Facts (continued)
<p>Active ingredients</p> <p>Chlorhexidine gluconate 3.15% w/v Antiseptic Isopropyl alcohol 70% v/v Antiseptic</p> <p>Use For the preparation of the patient's skin prior to surgery or injection</p> <p>Warnings</p> <p>For external use only Flammable, keep away from fire or flame Do not use with electrocautery procedures</p> <p>Do not use</p> <ul style="list-style-type: none"> • on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch <p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p>Directions</p> <ul style="list-style-type: none"> • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm) • tear open the pouch to expose the swabstick • remove the swabstick from package. Do not touch the foam tip. • prior to surgery or injection, apply swabstick to the procedure site • place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute • turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute • allow the prepped area to air dry for one and one half minutes (1.5 minutes) • do not blot or wipe dry • discard after a single use <p>Other information</p> <ul style="list-style-type: none"> • store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only <p>Inactive ingredient USP purified water</p> <p>Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>

PDI Professional Disposables International, Inc.
 Orangeburg, NY 10962-1376 USA
 1-800-999-6423 www.pdih.com



(01)3 0310819000995

8LS428CK_02_10_14.pdf

NDC 10819-4075-5 REORDER NO. 542850

***Pprevanitics®**
[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70% Swab)]
PATIENT PREOPERATIVE SKIN PREPARATION

*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

FOR EXTERNAL USE ONLY • Use for the preparation of the patient's skin prior to surgery or injection • Latex free • Helps reduce bacteria that potentially can cause skin infection

IMPORTANT: See insert for complete **Drug Facts** and safety information.

One Swabstick, 0.054 fl. oz. (1.6 mL) Each

SWABSTICK



(01)10310819000391

TEAR OPEN HERE

EXP

LOT

CONTENTS: One swabstick premoistened with 1.6 mL of a solution of 3.15% (w/w) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. **DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swabstick is approximately: dry site : 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm). Tear open the pouch to expose the swabstick. Remove the swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one minute. **WARNINGS:** For **external use only. Flammable. Keep away from fire or flame.** Do not use with electrocautery procedures. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

Professional, Disposable, International, Inc.
Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdlhc.com

***PDI**



2MS428

2MS428_02_10_14_Annotated.pdf

REORDER NO. S42850
 NDC 10819-4075-5

+Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
 Patient preoperative skin preparation

*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

Drug Facts		Drug Facts (continued)	
Active ingredients	Purpose	Do not use	
Chlorhexidine gluconate 3.15% w/v	Antiseptic	<ul style="list-style-type: none"> on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol for lumbar puncture or in contact with the meninges on open skin wounds or as a general skin cleanser under occlusive patch 	
Isopropyl alcohol 70% v/v	Antiseptic	When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
Use For the preparation of the patient's skin prior to surgery or injection			
Warnings			
For external use only Flammable, keep away from fire or flame Do not use with electrocautery procedures			

Drug Facts (continued)		Drug Facts (continued)	
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.		<ul style="list-style-type: none"> place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute allow the prepped area to air dry for one and one half minutes (1.5 minutes) do not blot or wipe dry discard after a single use 	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.		Other information <ul style="list-style-type: none"> store between 68-77°F (20-25°C) avoid freezing and excessive heat above 104°F (40°C) for hospital and professional use only 	
Directions <ul style="list-style-type: none"> use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm) tear open the pouch to expose the swabstick remove the swabstick from package. Do not touch the foam tip. prior to surgery or injection, apply swabstick to the procedure site 		Inactive ingredient USP purified water	
		Questions? Call 1-800-999-6423 , M-F (9:00 am-5:00 pm EST)	

Each swabstick contains 0.054 fl. oz. (1.6 mL)

+PDI Professional Disposables International, Inc.
 Orangeburg, NY 10962-1376 USA
 1-800-999-6423 www.pdihc.com

5LS428IS



This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
03/24/2014