

5HB10800\_05\_20\_14Draft

NDC 10818-1080-1

**Preventics® Swab**

[Chlorhexidine Gluconate (3.15%)<sup>\*</sup> and Isopropyl Alcohol (70%) Swab]  
 Antiseptic  
 Non-sterile Solution  
 Contains: 100 Individual Swabs, 0.034 fl. oz. (1 mL) Each

PDI®

RECORDER NO. B10800

NDC 10818-1080-1

**Preventics® Swab**

[Chlorhexidine Gluconate (3.15%)<sup>\*</sup> and Isopropyl Alcohol (70%) Swab]  
 Antiseptic  
 Non-sterile Solution  
 equivalent to 167 mg of Chlorhexidine Gluconate per pouch  
 For external use only  
 Single step, broad spectrum, persistent antiseptic that significantly reduces the number of bacteria on the skin  
 Use for the preparation of the patient's skin prior to an injection  
 Helps reduce bacteria that potentially can cause skin infection  
 Latex free  
 For hospital and professional use only  
 Contains: 100 Individual Swabs, 0.034 fl. oz. (1 mL) Each

PDI®

RECORDER NO. B10800

NDC 10818-1080-1

**Preventics® Swab**

Do not use if pouch is torn or if contents are exposed to chlorhexidine gluconate

PDI®

www.pdi.com

(813) 203-7600

(813) 203-7600

NDC 10818-1080-1

**Preventics® Swab**

[Chlorhexidine Gluconate (3.15%)<sup>\*</sup> and Isopropyl Alcohol (70%) Swab]  
 Antiseptic  
 Non-sterile Solution  
 Contains: 100 Individual Swabs, 0.034 fl. oz. (1 mL) Each

PDI®

RECORDER NO. B10800

NDC 10818-1080-1

**Preventics® Swab**

[Chlorhexidine Gluconate (3.15%)<sup>\*</sup> and Isopropyl Alcohol (70%) Swab]  
 Antiseptic  
 Non-sterile Solution  
 equivalent to 167 mg of Chlorhexidine Gluconate per pouch  
 For external use only  
 Single step, broad spectrum, persistent antiseptic that significantly reduces the number of bacteria on the skin  
 Use for the preparation of the patient's skin prior to an injection  
 Helps reduce bacteria that potentially can cause skin infection  
 Latex free  
 For hospital and professional use only  
 Contains: 100 Individual Swabs, 0.034 fl. oz. (1 mL) Each

PDI®

RECORDER NO. B10800

NDC 10818-1080-1

**Preventics® Swab**

Do not use if pouch is torn or if contents are exposed to chlorhexidine gluconate

PDI®

www.pdi.com

(813) 203-7600

(813) 203-7600

5HB10800\_05\_20\_14Draft\_Clean

**NET CONTENTS: 10 BOXES OF 100 INDIVIDUAL SWABS**  
**REORDER NO. B10800**  
**NDC 10819-1080-1**

**Prevantics® Swab**  
 [Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]  
 Antiseptic

Non-sterile Solution

\*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

<p><b>Drug Facts</b></p> <p><b>Active ingredients</b>                  Chlorhexidine gluconate 3.15% w/v..... Antiseptic                  Isopropyl alcohol 70% w/v..... Antiseptic</p> <p><b>Use</b> For the preparation of the patient's skin prior to an injection</p> <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>■ For external use only</li> <li>■ Flammable, keep away from fire or flame</li> <li>■ Do not use with electrocautery procedures</li> </ul> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol</li> <li>■ for lumbar puncture or in contact with the meninges</li> <li>■ on open skin wounds or as a general skin cleanser</li> <li>■ under occlusive patch</li> </ul> <p><b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p><b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away. ▶</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>■ maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)</li> <li>■ tear open packet and remove swab. Do not unfold swab.</li> <li>■ prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.</li> <li>■ allow the prepped area to air dry for 30 seconds</li> <li>■ do not blot or wipe dry</li> <li>■ discard after a single use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 68-77°F (20-25°C)</li> <li>■ avoid freezing and excessive heat above 104°F (40°C)</li> <li>■ for hospital and professional use only</li> </ul> <p><b>Inactive ingredient</b>                  USP purified water</p> <p><b>Questions?</b> Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>
---	--



Each swab contains 0.034 fl. oz. (1 mL)

**PDI** Professional Disposables International, Inc., Orangeburg, NY 10962-1376  
 USA • 1-800-999-6423 • [www.pdihc.com](http://www.pdihc.com)

8GB10800\_05\_20\_14draft\_Clean

REORDER NO. B11400  
 NDC 10819-1080-2

# Prevantics® Swab

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]

Antiseptic

Non-sterile Solution

Packaging is sterile when in sterile kit \*equivalent to 167 mg of Chlorhexidine Gluconate per pouch  
 Contains: 3000 Individual Swabs, 0.034 fl. oz. (1 mL) Each

Drug Facts (continued)	
<b>Drug Facts</b>	<b>Directions</b>
<b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v.....Antiseptic Isopropyl alcohol 70% v/v.....Antiseptic	<ul style="list-style-type: none"> <li>use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)</li> <li>tear open packet and remove swab. Do not unfold swab.</li> <li>prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.</li> <li>allow the prepped area to air dry for 30 seconds</li> <li>do not blot or wipe dry</li> <li>discard after a single use</li> </ul>
<b>Use</b> For the preparation of the patient's skin prior to an injection	<b>Other information</b>
<b>Warnings</b>	<ul style="list-style-type: none"> <li>store between 68-77°F (20-25°C)</li> <li>avoid freezing and excessive heat above 104°F (40°C)</li> <li>for hospital and professional use only</li> </ul>
<ul style="list-style-type: none"> <li>For external use only</li> <li>Flammable, keep away from fire or flame</li> <li>Do not use with electrocautery procedures</li> </ul>	<b>Inactive ingredient</b>
<ul style="list-style-type: none"> <li>Do not use</li> <li>on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol</li> <li>for lumbar puncture or in contact with the meninges</li> <li>on open skin wounds or as a general skin cleanser</li> <li>under occlusive patch</li> </ul>	USP-purified water
<p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right a way and contact a doctor.</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right a way.</p>	<b>Questions?</b> Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)



**PDI** Professional Disposables International, Inc., Orangeburg, NY 10962-1376  
 USA • 1-800-999-6423 • [www.pdiinc.com](http://www.pdiinc.com)

8KB11400\_05\_20\_14Draft\_Clean


**Preventics® Swab**  
 (Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab) Antiseptic Non-sterile Solution  
 Packaging is sterile when in sterile kit  
 \*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

REORDER NO. B11400  
 NDC 10819-1080-2

<b>Drug Facts</b>
<b>Active Ingredients</b> Chlorhexidine gluconate 3.15% w/v.....Antiseptic Isopropyl alcohol 70% w/v.....Antiseptic
<b>Use</b> For the preparation of the patient's skin prior to an injection
<b>Warnings</b> <ul style="list-style-type: none"> <li>For external use only</li> <li>Flammable, keep away from fire or flame</li> <li>Do not use with electrocautery procedures</li> </ul> Do not use <ul style="list-style-type: none"> <li>on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol</li> <li>for lumbar puncture or in contact with the meninges</li> <li>on open skin wounds or as a general skin cleanser</li> <li>under occlusive patch</li> </ul> When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

<b>Drug Facts</b> (continued) Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
<b>Directions</b> <ul style="list-style-type: none"> <li>use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)</li> <li>tear open packet and remove swab. Do not unfold swab prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.</li> <li>allow the prepped area to air dry for 30 seconds</li> <li>do not blot or wipe dry</li> <li>discard after a single use</li> </ul>
<b>Other information</b> <ul style="list-style-type: none"> <li>store between 68°-77° F (20-25°C)</li> <li>avoid freezing and excessive heat above 104°F (40°C)</li> <li>for hospital and professional use only</li> </ul>
<b>Inactive Ingredient</b> USP purified water
<b>Questions?</b> Call 1-800-999-6423, M-F 9:00 am-5:00 pm EST

Each Swab contains 0.034 fl. oz. (1 mL)  
 Professional Disposables International, Inc.  
 Orangeburg, NY 10962-1376 • USA • 1-800-999-6423  
 www.pdiinc.com 5KB114IS\_05\_20\_14Draft



5KB114IS\_05\_20\_14Draft\_Clean



SHS40750\_05\_20\_14Draft  
[Chlorhexidine Gluconate (3.15%) \* and Isopropyl Alcohol (70%) Swab]  
Patient preoperative skin preparation  
Non-sterile Solution  
\*equivalent to 268 mg of Chlorhexidine Gluconate per pouch  
Contains: 50 Individual Swabsticks,  
0.054 fl. oz. (1.6 mL) Each

## Prevantics® Swabstick

NDC 10819 4075 1

REORDER NO. S40750

REORDER NO. S40750

NDC 10819 4075 1

## Prevantics® Swabstick

### TEARALONG REFORMATION FOR DISPENSING

Drug Facts	
<b>Warnings</b>	<b>Directions</b>
<p>Do not use hypochlorite bleach on fabrics exposed to chlorhexidine gluconate.</p>	<p>Use for the preparation of the patient's skin prior to surgery or injection. Use for the preparation of the patient's skin prior to surgery or injection. Use for the preparation of the patient's skin prior to surgery or injection.</p>
<b>Directions</b>	<b>Other information</b>
<p>Use for the preparation of the patient's skin prior to surgery or injection. Use for the preparation of the patient's skin prior to surgery or injection. Use for the preparation of the patient's skin prior to surgery or injection.</p>	<p>Keep out of reach of children. See important information about prescription drugs on page 2 of this leaflet. See important information about prescription drugs on page 2 of this leaflet. See important information about prescription drugs on page 2 of this leaflet.</p>
<b>Other information</b>	<b>Contains</b>
<p>Contains: 50 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each</p>	<p>Contains: 50 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each</p>

## Prevantics® Swabstick

Do not use hypochlorite bleach on fabrics exposed to chlorhexidine gluconate.



Professional Disinfectants International, Inc.  
Cincinnati, OH 45222-1376 USA  
1-800-999-8428  
www.pdiinc.com



(01203106100028)



Contains: 50 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each

NDC 10819 4075 1

REORDER NO. S40750

## Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%) \* and Isopropyl Alcohol (70%) Swab]  
Patient preoperative skin preparation  
Non-sterile Solution  
\*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

- For external use only
- Single step, broad spectrum, persistent antiseptic that significantly reduces the number of microorganisms on intact skin
- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- Latex free
- For hospital and professional use only



Contains:  
50 Individual Swabsticks,  
0.054 fl.oz. (1.6 mL) Each

NDC 10819 4075 1

REORDER NO. S40750

## Prevantics® Swabstick

CONTENTS: 50 packets of swabsticks premoistened with 1.6 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)



LOT UNVARNISHED AREA  
EXP. UNVARNISHED AREA

SHS40750\_05\_20\_14Draft\_Clean

**NET CONTENTS: 10 BOXES OF 50 INDIVIDUAL SWABSTICKS**  
**REORDER NO. S40750**  
**NDC 10819-4075-1**

**Prevantics® Swabstick**  
 (Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab)  
 Patient preoperative skin preparation  
 Non-sterile Solution

\*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

<b>Drug Facts</b>	
<b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v..... Antiseptic Isopropyl alcohol 70% w/v..... Antiseptic	<b>Purpose</b> Antiseptic
<b>Use</b> For the preparation of the patient's skin prior to surgery or injection.	
<b>Warnings</b> <ul style="list-style-type: none"> <li>For external use only</li> <li>Flammable, keep away from fire or flame</li> <li>Do not use with electrocautery procedures</li> </ul>	
<b>Do not use</b> <ul style="list-style-type: none"> <li>on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol</li> <li>for lumbar puncture or in contact with the meninges</li> <li>on open skin wounds or as a general skin cleanser</li> <li>under occlusive patch</li> </ul>	
<b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
<b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> <ul style="list-style-type: none"> <li>use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> </ul>	
<b>Drug Facts (continued)</b>	
<ul style="list-style-type: none"> <li>maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)</li> <li>peel apart the pouch to expose the swabstick</li> <li>remove the swabstick from package. Do not touch the foam tip.</li> <li>prior to surgery or injection, apply swabstick to the procedure site</li> <li>Place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute</li> <li>turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>do not blot or wipe dry</li> <li>discard after a single use</li> </ul>	
<b>Other information</b> <ul style="list-style-type: none"> <li>store between 68-77°F (20-25°C)</li> <li>avoid freezing and excessive heat above 104°F (40°C)</li> <li>for hospital and professional use only</li> </ul>	
<b>Inactive ingredient</b> USP purified water	
<b>Questions?</b> Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	

**Each swabstick contains 0.054 fl.oz. (1.6 mL)**

**PDI** Professional Disposables International, Inc., Orangeburg, NY 10962-1376  
 USA • 1-800-999-6423 • [www.pdihc.com](http://www.pdihc.com)



(01)30310819000025  
 8GS40750\_05\_20\_14Draff



REORDER NO. S32450  
 NDC 10819-4075-2

# Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]  
 Patient preoperative skin preparation  
 Non-sterile Solution

Packaging is sterile when in sterile kit \*equivalent to 268 mg of Chlorhexidine Gluconate per pouch  
 Contains: 500 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each



Drug Facts (continued)	
<b>Drug Facts</b>	
<b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v ..... Antiseptic Isopropyl alcohol 70% v/v ..... Antiseptic	<b>Purpose</b> Patient preoperative skin preparation
<b>Use</b> For the preparation of the patient's skin prior to surgery or injection	
<b>Warnings</b>	
<ul style="list-style-type: none"> <li>For external use only</li> <li>Flammable, keep away from fire or flame</li> <li>Do not use with electrocautery procedures</li> </ul>	
<b>Do not use</b>	
<ul style="list-style-type: none"> <li>on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol</li> <li>for laceration or in contact with the meninges</li> <li>on open skin wounds or as a general skin cleanser</li> <li>under occlusive patch</li> </ul>	
<b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
<b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b>	<ul style="list-style-type: none"> <li>use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)</li> <li>tear open the pouch to expose the swabstick</li> <li>remove the swabstick from package. Do not touch the foam tip</li> <li>prior to surgery or injection, apply swabstick to the procedure site</li> <li>place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute</li> <li>turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>do not blot or wipe dry</li> <li>discard after a single use</li> </ul>
<b>Other information</b>	<ul style="list-style-type: none"> <li>store between 68-77°F (20-25°C)</li> <li>avoid freezing and excessive heat above 104°F (40°C)</li> <li>for hospital and professional use only</li> </ul>
<b>Inactive ingredient</b>	USP purified water
<b>Questions?</b>	Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)

**PDI** Professional Disposables International, Inc., Orangeburg, NY 10962-1376  
 USA • 1-800-999-6423 • [www.pdific.com](http://www.pdific.com)

8KS32450\_05\_20\_14draft\_Clean

REORDER NO. S32450  
 NDC 10819-4075-2

# Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]  
 Patient preoperative skin preparation

Non-sterile Solution

Packaging is sterile when in sterile kit

\*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

<p><b>Drug Facts</b></p> <p><b>Active Ingredients</b></p> <p>Chlorhexidine gluconate 3.15% w/v..... Antiseptic              Isopropyl alcohol 70% v/v..... Antiseptic</p> <p><b>Purpose</b></p> <p>Antiseptic</p> <p><b>Use</b> For the preparation of the patient's skin prior to surgery or injection</p> <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>■ For external use only</li> <li>■ Flammable, keep away from fire or flame</li> <li>■ Do not use with electrocautery procedures</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol</li> <li>■ for lumbar puncture or in contact with the meninges</li> <li>■ on open skin wounds or as a general skin cleanser</li> <li>■ under occlusive patch</li> </ul> <p><b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p>
--	--

<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>■ maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)</li> <li>■ tear open the pouch to expose the swabstick</li> <li>■ remove the swabstick from package. Do not touch the foam tip.</li> <li>■ prior to surgery or injection, apply swabstick to the procedure site</li> <li>■ place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>■ allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>■ do not blot or wipe dry</li> <li>■ discard after a single use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 68-77°F (20-25°C)</li> <li>■ avoid freezing and excessive heat above 104°F (40°C)</li> <li>■ for hospital and professional use only</li> </ul> <p><b>Inactive ingredient</b></p> <p>USP purified water</p> <p><b>Questions?</b> Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>
--	---

Each swabstick contains 0.054 fl. oz. (1.6 mL)

**PDI** Professional Disposables International, Inc.  
 Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • [www.pdihc.com](http://www.pdihc.com)



5KS324 S\_05\_20\_14Dra t

5KS324IS\_05\_20\_14Draft\_Clean

5HS41950 05 20 14Draft

**PDI**  
Contains: 30 Individual Maxi Swabsticks,  
0.172 fl. oz. (5.1 mL) Each

**Preventics® Maxi Swabstick**  
[Chlorhexidine Gluconate (3.15%)<sup>\*</sup> and Isopropyl Alcohol (70%) Swab]  
Patient/preoperative skin preparation  
Non-sterile Solution  
<sup>\*</sup>equivalent to 855 mg of Chlorhexidine Gluconate per pouch

**Preventics® Maxi Swabstick**  
NDC 10619 4076 4 REORDER NO. 541950

**Preventics® Maxi Swabstick**  
NDC 10619 4076 4 REORDER NO. 541950

TEAR ALONG PERFORATION FOR DISINTEGRATION

Usage Facts	
<b>Indications</b>	For the preparation of patient and preoperative skin preparation.
<b>Directions</b>	1. Remove the pouch from the box. 2. Tear along the perforation to open the pouch. 3. Remove the swabstick from the pouch. 4. Use the swabstick to prepare the skin. 5. Discard the swabstick after use.
<b>Warnings</b>	Do not use hypochlorite bleach on fabrics exposed to chlorhexidine gluconate.
<b>Contraindications</b>	None.
<b>Adverse Reactions</b>	None.
<b>How Supplied</b>	30 Individual Maxi Swabsticks, 0.172 fl. oz. (5.1 mL) Each.
<b>Other information</b>	Professional Disposables International, Inc. Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdiinc.com



**PDI**  
Contains: 30 Individual Maxi Swabsticks,  
0.172 fl. oz. (5.1 mL) Each

**Preventics® Maxi Swabstick**  
NDC 10619 4076 4 REORDER NO. 541950

Do not use hypochlorite bleach on fabrics exposed to chlorhexidine gluconate.



Professional Disposables International, Inc.  
Orangeburg, NY 10962-1376 USA  
1-800-999-6423  
www.pdiinc.com

**Preventics® Maxi Swabstick**  
NDC 10619 4076 4 REORDER NO. 541950

[Chlorhexidine Gluconate (3.15%)<sup>\*</sup> and Isopropyl Alcohol (70%) Swab]  
Patient/preoperative skin preparation  
Non-sterile Solution  
<sup>\*</sup>equivalent to 855 mg of Chlorhexidine Gluconate per pouch

- For external use only
- Single step, broad spectrum, persistent antiseptic that significantly reduces the number of microorganisms on intact skin
- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- Latex free
- For hospital and professional use only

**PDI**  
Contains: 30 Individual Maxi Swabsticks,  
0.172 fl. oz. (5.1 mL) Each

**Preventics® Maxi Swabstick**  
NDC 10619 4076 4 REORDER NO. 541950

**CONTENTS:** 30 packets of swabsticks premoistened with 5.1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)

**PDI**  
LOT: UNVARNISHED AREA  
EXP: UNVARNISHED AREA

5HS41950\_05\_20\_14Draft\_Clean

**NET CONTENTS: 10 BOXES OF 30 INDIVIDUAL SWABSTICKS**  
**REORDER NO. S41950**  
**NDC 10819-4075-4**

**Prevantics® Maxi Swabstick**

(Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab)  
Patient preoperative skin preparation  
Non-sterile Solution

\*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

<p><b>Drug Facts</b></p> <p><b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v .....Antiseptic Isopropyl alcohol 70% w/v .....Antiseptic</p> <p><b>Use</b> For the preparation of the patient's skin prior to surgery or injection</p> <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>■ For external use only</li> <li>■ Flammable, keep away from fire or flame</li> <li>■ Do not use with electrocautery procedures</li> </ul> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>■ on patients with known allergies to chlorhexidine</li> <li>■ for contact with mucous membranes</li> <li>■ for contact with the meninges</li> <li>■ on open skin wounds or as a general skin cleanser</li> <li>■ under occlusive patch</li> </ul> <p><b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p><b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>■ maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)</li> <li>■ peel apart the pouch to expose the swabstick</li> <li>■ remove the swabstick from package. Do not touch the foam tip.</li> <li>■ prior to surgery or injection, apply swabstick to the procedure site</li> <li>■ place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute</li> <li>■ turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>■ allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>■ do not blot or wipe dry</li> <li>■ discard after a single use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 68-77°F (20-25°C)</li> <li>■ avoid freezing and excessive heat above 104°F (40°C)</li> <li>■ for hospital and professional use only</li> </ul> <p><b>Inactive ingredient</b> USP purified water</p> <p><b>Questions?</b> Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>
---	---



Each swabstick contains 0.172 fl.oz. (5.1 mL)



Professional Disposables International, Inc., Orangeburg, NY 10962-1376  
USA • 1-800-999-6423 • [www.pdinc.com](http://www.pdinc.com)



86S41950\_05\_20\_14Draft\_Clean

**Prevantics® Maxi Swabstick** REORDER NO. S27350  
 [Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab] NDC 10819-4075-3  
 Patient preoperative skin preparation  
 Non-sterile Solution

Packaging is sterile when in sterile kit  
 \*equivalent to 855 mg of Chlorhexidine Gluconate per pouch  
**Contains: 300 Individual Swabsticks, 0.172 fl. oz. (5.1 mL) Each**

<b>Drug Facts</b>	
<b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v.....Antiseptic Isopropyl alcohol 70% w/v.....Antiseptic	<b>Purpose</b> Antiseptic
<b>Use</b> For the preparation of the patient's skin prior to surgery or injection	
<b>Warnings</b> ■ For external use only ■ Flammable, keep away from fire or flame ■ Do not use with electrocautery procedures  Do not use ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser ■ under occlusive patch  When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.  Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.  Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	<b>Directions</b> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm) ■ tear open the pouch to expose the swabstick ■ remove the swabstick from package. Do not touch the foam tip. ■ prior to surgical or injection, apply swabstick to the procedure site ■ place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute ■ turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute ■ allow the prepped area to air dry for one and one half minutes (1.5 minutes) ■ do not blot or wipe dry ■ discard after a single use
	<b>Other information</b> ■ store between 68-77°F (20-25°C) ■ avoid freezing and excessive heat above 104°F (40°C) ■ for hospital and professional use only
	<b>Inactive ingredient</b> USP purified water
	<b>Questions?</b> Call 1-800-999-6423, M-F (900 am-500 pm EST)



**PDI** Professional Disposables International, Inc., Orangeburg, NY 10962-1376  
 USA • 1-800-999-6423 • [www.pdihc.com](http://www.pdihc.com)

8127350\_05\_20\_14Draft\_Clean

REORDER NO. S27350  
 NDC 10819-4075-3

# Prevantics® Maxi Swabstick

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]  
 Patient preoperative skin preparation  
 Non-sterile Solution

**Packaging is sterile when in sterile kit**

\*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

Drug Facts	
<b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v..... Antiseptic Isopropyl alcohol 70% v/v..... Antiseptic	<b>Purpose</b> Antiseptic
<b>Use</b> For the preparation of the patient's skin prior to surgery or injection	
<b>Warnings</b>	
<ul style="list-style-type: none"> <li>■ For external use only</li> <li>■ Flammable, keep away from fire or flame</li> <li>■ Do not use with electrocautery procedures</li> </ul>	
<b>Do not use</b>	
<ul style="list-style-type: none"> <li>■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol</li> <li>■ for lumbar puncture or in contact with the meninges</li> <li>■ on open skin wounds or as a general skin cleanser</li> <li>■ under occlusive patch</li> </ul>	
<b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	

## Drug Facts (continued)

<b>Drug Facts (continued)</b>	<b>Drug Facts (continued)</b>
<b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	<ul style="list-style-type: none"> <li>■ place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>■ turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>■ allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>■ do not blot or wipe dry</li> <li>■ discard after a single use</li> </ul>
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	<b>Other information</b>
<b>Directions</b>	<ul style="list-style-type: none"> <li>■ store between 68-77°F (20-25°C)</li> <li>■ avoid freezing and excessive heat above 104°F (40°C)</li> <li>■ for hospital and professional use only</li> </ul>
<ul style="list-style-type: none"> <li>■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>■ maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)</li> <li>■ tear open the pouch to expose the swabstick</li> <li>■ remove the swabstick from package. Do not touch the foam tip.</li> <li>■ prior to surgery or injection, apply swabstick to the procedure site</li> </ul>	<b>Inactive ingredient</b> USP purified water
	<b>Questions?</b> Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)

Each Maxi Swabstick contains 0.172 fl. oz. (5.1 mL)

**PDI** Professional Disposables International, Inc.  
 Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • [www.pdihc.com](http://www.pdihc.com)



5K5273IS\_05\_20\_14Draft

**REORDER NO. S42850**  
**NDC 10819-4075-5**

**Prevantics® Swabstick**  
 [Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]  
 Patient preoperative skin preparation  
 Non-sterile Solution

Packaging is sterile when in sterile kit

\*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

Contains: 500 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each

<b>Drug Facts</b>	
<b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v..... Antiseptic Isopropyl alcohol 70% v/v..... Antiseptic	<b>Purpose</b> Antiseptic
<b>Use</b> For the preparation of the patient's skin prior to surgery or injection.	
<b>Warnings</b> <ul style="list-style-type: none"> <li>■ For external use only</li> <li>■ Flammable, keep away from fire or flame</li> <li>■ Do not use with electrocautery procedures</li> </ul> <b>Do not use</b> <ul style="list-style-type: none"> <li>■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol</li> <li>■ for lumbar puncture or in contact with the meninges</li> <li>■ on open skin wounds or as a general skin cleanser</li> <li>■ under occlusive patch</li> </ul> <b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
<b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> <ul style="list-style-type: none"> <li>■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>■ maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)</li> <li>■ tear open the pouch to expose the swabstick</li> <li>■ remove the swabstick from package. Do not touch the foam tip.</li> <li>■ prior to surgery or injection, apply swabstick to the procedure site</li> <li>■ place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>■ turn the swabstick over (unisex side of the foam tip) and repeat the procedure by prepping for one (1) minute (1.5 minutes)</li> <li>■ allow the prepped area to air dry for one and one half minutes</li> <li>■ do not blot or wipe dry</li> <li>■ discard after a single use</li> </ul>	
<b>Other information</b> <ul style="list-style-type: none"> <li>■ store at 68-77°F (20-25°C)</li> <li>■ avoid freezing and excessive heat above 104°F (40°C)</li> <li>■ for hospital and professional use only</li> </ul>	
<b>Inactive ingredient</b> USP purified water	
<b>Questions ?</b> Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	



**PDI** Professional Disposables International, Inc., Orangeburg, NY 10962-1376  
 USA • 1-800-999-6423 • [www.pdihc.com](http://www.pdihc.com)

8KS428CK\_05\_20\_14Draft\_Clean

REORDER NO. S42850  
 NDC 10819-4075-5

# Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]  
 patient preoperative skin preparation

Non-sterile Solution

Packaging is sterile when in sterile kit

\*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

Drug Facts (continued)	
<p><b>Drug Facts</b></p> <p><b>Active Ingredients</b>                      Chlorhexidine gluconate 3.15% w/v..... Antiseptic                      Isopropyl alcohol 70% w/v..... Antiseptic</p> <p><b>Use</b> For the preparation of the patient's skin prior to surgery or injection</p> <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>■ For external use only</li> <li>■ Flammable, keep away from fire or flame</li> <li>■ Do not use with electrocautery procedures</li> </ul>	<p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol</li> <li>■ for lumbar puncture or in contact with the meninges</li> <li>■ on open skin wounds or as a general skin cleanser</li> <li>■ under occlusive patch</li> </ul> <p><b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p>

<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>■ maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm).</li> <li>■ tear open the pouch to expose the swabstick</li> <li>■ remove the swabstick from package. Do not touch the foam tip.</li> <li>■ prior to surgery or injection, apply swabstick to the procedure site</li> <li>■ place one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes for one (1) minute</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>■ allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>■ do not blot or wipe dry</li> <li>■ discard after a single use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 68-77°F (20-25°C)</li> <li>■ avoid freezing and excessive heat above 104°F (40°C)</li> <li>■ for hospital and professional use only</li> </ul> <p><b>Inactive Ingredient</b>                      USP purified water</p> <p><b>Questions?</b> Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>
---	---

Each swabstick contains 0.054 fl. oz. (1.6 mL)

**PDI®** Professional Disposables International, Inc.  
 Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • [www.pdlinc.com](http://www.pdlinc.com)



5K5428IS\_05\_20\_14Draft