

**PDI**

Contains: 100 Individual Swabs, 0.034 fl. oz. (1 mL) Each

\*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

**ANTISEPTIC  
Non-sterile Solution**

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]

**Prevantics**

REORDER NO. B10800  
NDC 10819-1080-1

5NB10800

NDC 10819-1080-1

REORDER NO. B10800

**SWAB**

<p><b>Drug Facts</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="text-align: left;">Active ingredients</th> <th style="text-align: left;">Purpose</th> </tr> <tr> <td>Chlorhexidine gluconate 3.15% w/v</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v</td> <td>Antiseptic</td> </tr> </table> <p><b>Use</b> For the preparation of the patient's skin prior to an injection</p> <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>• For external use only. • Flammable, keep away from fire or flame.</li> <li>• Do not use with electrocautery procedures</li> </ul> <p><b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul>	Active ingredients	Purpose	Chlorhexidine gluconate 3.15% w/v	Antiseptic	Isopropyl alcohol 70% v/v	Antiseptic	<p><b>Drug Facts</b> (continued)</p> <p><b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p><b>Stop use and ask a doctor if</b> irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm).</li> <li>• tear open packet and remove swab. Do not unfold swab.</li> <li>• prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.</li> <li>• allow the prepped area to air dry for 30 seconds.</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul>	<p><b>Drug Facts</b> (continued)</p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>• store between 68–77°F (20–25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul> <p><b>Inactive ingredient</b> USP purified water</p> <p><b>Questions?</b> call 1-800-999-6423, M-F (9:00 am–5:00 pm EST)</p>
Active ingredients	Purpose							
Chlorhexidine gluconate 3.15% w/v	Antiseptic							
Isopropyl alcohol 70% v/v	Antiseptic							

**Prevantics**

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]

**ANTISEPTIC  
Non-sterile Solution**

\*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

**FOR EXTERNAL USE ONLY**

- Single step, broad spectrum, persistent antiseptic that significantly reduces the number of microorganisms on intact skin
- Use for the preparation of the patient's skin prior to an injection
- Helps reduce bacteria that potentially can cause skin infection
- Latex free
- For hospital and professional use only

Contains: 100 Individual Swabs, 0.034 fl.oz. (1 mL) Each

REORDER NO. B10800

**SWAB**

**STORAGE**


- store between 68–77°F (20–25°C)
- avoid freezing and excessive heat above 104°F (40°C)

**CONTENTS:**  
100 packets of swabs premoistened with 1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water


**Do not use hypochlorite bleach on fabrics exposed to chlorhexidine gluconate.**

LOT

EXP.



(01)10310819000014



Professional Disposables International, Inc.  
 Orangetown, NY  
 10962-1376 USA  
 1-800-999-6423  
 www.pdihc.com

5NB10800\_2017\_07\_27.PDF

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

**TEAR BELOW**

**EXP. LOT**

**CONTENTS:** One swab premoistened with 1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. **DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm). Tear open packet and remove swab. Do not unfold swab. Prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds. Allow the prepped area to air dry for 30 seconds. Do not blot or wipe dry. Discard after a single use. **WARNINGS:** For external use only. Flammable, keep away from fire or flame. **Allergy Alert** This product may cause a severe allergic reaction. Symptoms include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.  
Professional Disposables International, Inc., Orangeburg, NY 10962-1376 USA  
1-800-999-6423 www.pdihc.com

20B108

NDC 10819-1080-1 REORDER NO. B10800

**Prevantics®**  
[Chlorhexidine Gluconate (3.15%)\*  
and Isopropyl Alcohol (70%) Swab]


**ANTISEPTIC**  
**Non-sterile Solution**  
\*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

**FOR EXTERNAL USE ONLY**


- Use for the preparation of the patient's skin prior to an injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only • Latex free
- Do not use with electrocautery procedures

**IMPORTANT:** See carton for complete **Drug Facts** and safety information.


One Swab, 0.034 fl. oz. (1 mL) Each




(01)00310819000017



**SWAB**



2QB108\_2017\_03\_03.pdf




**NET CONTENTS: 10 BOXES OF 100 INDIVIDUAL SWABS**  
**REORDER NO. B10800**  
**NDC 10819-1080-1**


**[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]**  
**Antiseptic**  
**Non-sterile Solution**  
 \*equivalent to 167 mg of Chlorhexidine Gluconate per pouch


<p><b>Drug Facts</b></p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%;"><b>Active ingredients</b></td> <td style="width: 50%;"><b>Purpose</b></td> </tr> <tr> <td>Chlorhexidine gluconate 3.15% w/v</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v</td> <td>Antiseptic</td> </tr> </table> <p><b>Use</b> For the preparation of the patient's skin prior to an injection</p> <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>• For external use only</li> <li>• <b>Flammable, keep away from fire or flame</b></li> <li>• Do not use with electrocautery procedures</li> </ul> <p><b>Allergy Alert</b>              This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash              If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul> <p><b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p><b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p>	<b>Active ingredients</b>	<b>Purpose</b>	Chlorhexidine gluconate 3.15% w/v	Antiseptic	Isopropyl alcohol 70% v/v	Antiseptic	<p><b>Drug Facts</b> (continued)</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)</li> <li>• tear open packet and remove swab. Do not unfold swab.</li> <li>• prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.</li> <li>• allow the prepped area to air dry for 30 seconds</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>• store between 68-77°F (20-25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul> <p><b>Inactive ingredient</b>              USP purified water</p> <p><b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>
<b>Active ingredients</b>	<b>Purpose</b>						
Chlorhexidine gluconate 3.15% w/v	Antiseptic						
Isopropyl alcohol 70% v/v	Antiseptic						

**Each swab contains 0.034 fl. oz. (1 mL)**




Professional Disposables International, Inc.  
 Orangeburg, NY 10962-1376 USA  
 1-800-999-6423 www.pdihc.com





(01)20310819000011






8MB10800

LBL878v02

8MB10800\_2017\_07\_27.PDF

Prevantics Swab bulk outer carton 3000-count

		REORDER NO. B11400 NDC 10819-1080-2	
[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab] <b>Antiseptic</b> <b>Non-sterile Solution</b> <b>Packaging is sterile when in sterile kit</b> *equivalent to 167 mg of Chlorhexidine Gluconate per pouch <b>Contains: 3000 Individual Swabs, 0.034 fl. oz. (1 mL) Each</b>			
<b>Drug Facts</b> <b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v ..... Antiseptic Isopropyl alcohol 70% v/v ..... Antiseptic		<b>Drug Facts (continued)</b> <b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Use</b> For the preparation of the patient's skin prior to an injection		<b>Directions</b> <ul style="list-style-type: none"> <li>use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)</li> <li>tear open packet and remove swab. Do not unfold swab.</li> <li>prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.</li> <li>allow the prepped area to air dry for 30 seconds</li> <li>do not blot or wipe dry</li> <li>discard after a single use</li> </ul>	
<b>Warnings</b> <ul style="list-style-type: none"> <li>For external use only</li> <li>Flammable, keep away from fire or flame</li> <li>Do not use with electrocautery procedures</li> </ul>		<b>Other information</b> <ul style="list-style-type: none"> <li>store between 68-77°F (20-25°C)</li> <li>avoid freezing and excessive heat above 104°F (40°C)</li> <li>for hospital and professional use only</li> </ul>	
<b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash If an allergic reaction occurs, stop use and seek medical help right away.		<b>Inactive ingredient</b> USP purified water	
<b>Do not use</b> <ul style="list-style-type: none"> <li>on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>for lumbar puncture or in contact with the meninges</li> <li>on open skin wounds or as a general skin cleanser</li> <li>under occlusive patch</li> </ul>		<b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	
<b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.		8QB11400	
 Professional Disposables International, Inc. Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdihc.com		 (01)20310819000868	
		LBL878v02	

8QB11400\_2017\_07\_27.PDF

NDC 10819-1080-2 REORDER NO. B11400

**Prevantics®**  
[Chlorhexidine Gluconate (3.15%)\*  
and Isopropyl Alcohol (70%) Swab]

**ANTISEPTIC**  
**Non-sterile Solution**  
**Packaging is sterile when in sterile kit**  
\*equivalent to 1.67 mg of Chlorhexidine Gluconate per pouch

**FOR EXTERNAL USE ONLY**

- Use for the preparation of the patient's skin prior to an injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only • Latex free
- Do not use with electrocautery procedures

**IMPORTANT:** See insert for complete **Drug Facts** and safety information.  
**One Swab**, 0.034 fl. oz. (1 mL) Each

**SWAB**

**TEAR BELOW**


**CONTENTS:** One swab premoistened with 1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. **DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm). Tear open packet and remove swab. Do not unfold swab. Prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds. Allow the prepped area to air dry for 30 seconds. Do not blot or wipe dry. Discard after a single use. **WARNINGS: For external use only. Flammable, keep away from fire or flame.**


**Allergy Alert:** This product may cause a severe allergic reaction. Symptoms include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

Professional Disposables International, Inc.  
Orangeburg, NY 10962-1376 USA  
1-800-999-6423 [www.pdihc.com](http://www.pdihc.com)

LOT  
EXP.

01100310818000884






2PB114X

2PB114X\_2017\_03\_06.pdf

<b>Prevanitics<sup>®</sup> Swab</b> REORDER NO. B11400 NDC 10819-1080-2	
[Chlorhexidine Gluconate 3.15%]* and Isopropyl Alcohol [70%] Swab] Antiseptic Non-sterile Solution • Packaging is sterile when in sterile kit *equivalent to 167 mg of Chlorhexidine Gluconate per pouch	
<b>Drug Facts</b>	
<b>Active ingredients</b>	<b>Purpose</b>
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic
<b>Use</b> For the preparation of the patient's skin prior to an injection	
<b>Warnings</b>	
• For external use only	
• Flammable, keep away from fire or flame	
• Do not use with electrocautery procedures	
<b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include:	
• Wheezing/difficulty breathing • Shock • Facial swelling	
• Hives • Rash	
If an allergic reaction occurs, stop use and seek medical help right away.	
<b>Do not use</b>	
• on patients allergic to chlorhexidine gluconate or any other ingredient in this product	
• for lumbar puncture or in contact with the meninges	
• on open skin wounds or as a general skin cleanser	
• under occlusive patch	
<b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent	

MISC025v00

<b>Drug Facts (continued)</b>
injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.
<b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.
<b>Directions</b>
• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
• maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)
• tear open packet and remove swab. Do not unfold swab.
• prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.
• allow the prepped area to air dry for 30 seconds
• do not blot or wipe dry
• discard after a single use
<b>Other information</b>
• store between 68-77 °F (20-25 °C)
• avoid freezing and excessive heat above 104 °F (40 °C)
• for hospital and professional use only
<b>Inactive ingredient</b> USP purified water
<b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)
Each Swab contains 0.034 fl. oz. (1 mL)
<b>PDI</b> Professional Disposables International, Inc. Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdihc.com
 
5PB114IS

MISC025v00

5PB114IS\_2017\_07\_27.PDF

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

TEAR ALONG PERFORATION FOR DISPENSING

Drug Facts	Drug Facts (continued)
<p><b>Active ingredients</b></p> <p>Chlorhexidine gluconate 3.15% w/v ..... Antiseptic Isopropyl alcohol 70% v/v ..... Antiseptic</p> <p><b>Use</b> For the preparation of the patient's skin prior to surgery or injection</p> <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>• For external use only</li> <li>• <b>Flammable, keep away from fire or flame</b></li> <li>• Do not use with electrocautery procedures</li> </ul> <p><b>Allergy Alert</b></p> <p>This product may cause a severe allergic reaction. Symptoms may include:</p> <ul style="list-style-type: none"> <li>• Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash</li> </ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul> <p><b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p><b>Stop use and ask a doctor if</b> irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p>	<p><b>Directions</b></p> <ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swabstick is approximately: dry site 4 by 4 inches (10 by 10 cm) or moist site 3 by 5 inches (7.5 by 12.5 cm)</li> <li>• peel apart the pouch to expose the swabstick</li> <li>• remove the swabstick from package. Do not touch the foam tip.</li> <li>• prior to surgery or injection, apply swabstick to the procedure site</li> <li>• place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>• allow the prepped area to air-dry for one and one half minutes (1.5 minutes)</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>• store between 68-77°F (20-25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul> <p><b>Inactive ingredient</b> USP purified water</p> <p><b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>

5PS40750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]  
**PATIENT PREOPERATIVE SKIN PREPARATION**  
**Non-sterile Solution**  
\*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

**FOR EXTERNAL USE ONLY**

- Single step, broad spectrum, persistent antiseptic that significantly reduces the number of microorganisms on intact skin
- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- Latex free
- For hospital and professional use only

Professional Disposables International, Inc.  
Orangeburg, NY 10962-1376 USA  
1-800-999-6423 www.pdiinc.com



(01)1031081900021

Contains: 50 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 540750

**SWABSTICK**

NDC 10819-4075-1  
REORDER NO. 540750

**Prevantics**<sup>®</sup>

NDC 10819-4075-1  
REORDER NO. 54

NDC 10819-4075-1 REORDER NO. S40750

**Prevantics®**

[Chlorhexidine Gluconate (3.15%)\*  
and Isopropyl Alcohol (70%) Swab]

**PATIENT PREOPERATIVE SKIN PREPARATION**

**Non-sterile Solution**

\*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

**FOR EXTERNAL USE ONLY** • Use for the preparation of the patient's skin prior to surgery or injection

• Helps reduce bacteria that potentially can cause skin infection • For hospital and professional use only • Latex free

**IMPORTANT:** See carton for complete **Drug Facts** and safety information.

One Swabstick, 0.054 fl. oz. (1.6 mL) Each



SWABSTICK



**CONTENTS:** One swabstick premoistened with 1.6 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. **DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm). Peel apart the pouch to expose the swabstick. Remove the swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one and one half minutes (1.5 minutes). Do not blot or wipe dry. Discard after a single use. **WARNINGS: For external use only. Flammable, keep away from fire or flame.** Do not use with electrocautery procedures. **Allergy Alert** This product may cause severe allergic reaction. Symptoms include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away. Professional Disposables International, Inc. Orangeburg, NY 10962-1876 USA 1-800-999-6428 [www.pdihc.com](http://www.pdihc.com)



(01)00310819000024



LOT  
PK33

THIS SIDE UP  
PEEL APART HERE

2PS407

2PS407\_2017\_03\_03.pdf

**NET CONTENTS: 10 BOXES OF 50 INDIVIDUAL SWABSTICKS**  
**REORDER NO. S40750**  
**NDC 10819-4075-1**

**†Prevantics® Swabstick**  
**[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]**  
**Patient preoperative skin preparation**  
**Non-sterile Solution**  
 \*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

<p><b>Drug Facts</b></p> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%;"><b>Active ingredients</b></td> <td style="width: 50%;"><b>Purpose</b></td> </tr> <tr> <td>Chlorhexidine gluconate 3.15% w/v</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v</td> <td>Antiseptic</td> </tr> </table> <p><b>Use</b> For the preparation of the patient's skin prior to surgery or injection</p> <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>• For external use only</li> <li>• Flammable, keep away from fire or flame</li> <li>• Do not use with electrocautery procedures</li> </ul> <p><b>Allergy Alert</b>          This product may cause a severe allergic reaction. Symptoms may include:          • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash          If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul> <p><b>When using this product</b> Keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p><b>Stop use and ask a doctor if</b> irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p>	<b>Active ingredients</b>	<b>Purpose</b>	Chlorhexidine gluconate 3.15% w/v	Antiseptic	Isopropyl alcohol 70% v/v	Antiseptic	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swabstick is approximately:                      dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)</li> <li>• peel apart the pouch to expose the swabstick</li> <li>• remove the swabstick from package. Do not touch the foam tip.</li> <li>• prior to surgery or injection, apply swabstick to the procedure site</li> <li>• place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>• turn the swabstick over [unused side of the foam tip] and repeat the procedure by prepping for one (1) minute</li> <li>• allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>• store between 68-77°F (20-25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul> <p><b>Inactive ingredient</b>          USP purified water</p> <p><b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>
<b>Active ingredients</b>	<b>Purpose</b>						
Chlorhexidine gluconate 3.15% w/v	Antiseptic						
Isopropyl alcohol 70% v/v	Antiseptic						

**Each swabstick contains 0.054 fl. oz. (1.6 mL)**





Professional Disposables International, Inc.  
 Orangeburg, NY 10962-1376 USA  
 1-800-999-6423 www.pdihc.com

(01)20310819000028

8NS40750

LBL878V02

8NS40750\_2017\_07\_27.PDF

		REORDER NO. S32450 NDC 10819-4075-2	
[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab] Patient preoperative skin preparation Non-sterile Solution Packaging is sterile when in sterile kit			
			
		*equivalent to 268 mg of Chlorhexidine Gluconate per pouch (01)20310819000875	
<b>Contains: 500 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each</b>			
<b>Drug Facts</b>		<b>Drug Facts (continued)</b>	
<b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v ..... Antiseptic Isopropyl alcohol 70% v/v ..... Antiseptic		<b>Directions</b> <ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)</li> <li>• tear open the pouch to expose the swabstick</li> <li>• remove the swabstick from package. Do not touch the foam tip.</li> <li>• prior to surgery or injection, apply swabstick to the procedure site</li> <li>• place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>• allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul>	
<b>Use</b> For the preparation of the patient's skin prior to surgery or injection		<b>Other information</b> <ul style="list-style-type: none"> <li>• store between 68-77°F (20-25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul>	
<b>Warnings</b> <ul style="list-style-type: none"> <li>• For external use only</li> <li>• <b>Flammable, keep away from fire or flame</b></li> <li>• Do not use with electrocautery procedures</li> </ul>		<b>Inactive ingredient</b> USP purified water	
<b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash. If an allergic reaction occurs, stop use and seek medical help right away.		<b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	
<b>Do not use</b> <ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul>		<b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
<b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.		 	
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.		PDI Professional Disposables International, Inc. • Orangeburg, NY 10962-1376 USA • 1-800-999-6423 www.pdihc.com	

8RS32450

LBL878v02

8RS32450\_2017\_07\_27.PDF

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

NDC 10819-4075-2 REORDER NO. S32450

**Prevantics**<sup>®</sup>

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]

**PATIENT PREOPERATIVE SKIN PREPARATION**

**Non-sterile Solution**

**Packaging is sterile when in sterile kit**

\*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

**FOR EXTERNAL USE ONLY** • Use for the preparation of the patient's skin prior to surgery or injection

• Helps reduce bacteria that potentially can cause skin infection • For hospital and professional use only • Latex free **IMPORTANT:** See insert for complete **Drug Facts** and safety information.

**One Swabstick**, 0.054 fl.oz. (1.6 mL) Each



SWABSTICK



**CONTENTS:** One swabstick premoistened with 1.6 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. **DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm). Tear open the pouch to expose the swabstick. Remove the swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one and one half minutes (1.5 minutes). Do not blot or wipe dry. Discard after a single use. **WARNINGS: For external use only. Flammable, keep away from fire or flame.** Do not use with electrocautery procedures. **Allergy Alert** This product may cause a severe allergic reaction. Symptoms include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

Professional Disposables International, Inc.  
Orangeburg, NY 10962-1376 USA 1-800-999-6423 [www.pdiho.com](http://www.pdiho.com)



(01)00310819000871



2PS324

2PS324\_2017\_MAR\_01.ai

REORDER NO. S32450 NDC 10819-4075-2							
<h1 style="margin: 0;">+Prevantics® Swabstick</h1> <p style="margin: 0;"><b>[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]</b>                  Patient preoperative skin preparation                  Non-sterile Solution                  Packaging is sterile when in sterile kit</p> <p style="margin: 0; font-size: small;">*equivalent to 268 mg of Chlorhexidine Gluconate per pouch</p>							
<p><b>Drug Facts</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 2px;"><b>Active ingredients</b></td> <td style="width: 50%; padding: 2px;"><b>Purpose</b></td> </tr> <tr> <td style="padding: 2px;">Chlorhexidine gluconate 3.15% w/v</td> <td style="padding: 2px;">Antiseptic</td> </tr> <tr> <td style="padding: 2px;">Isopropyl alcohol 70% v/v</td> <td style="padding: 2px;">Antiseptic</td> </tr> </table> <p style="padding: 5px;"><b>Use</b> For the preparation of the patient's skin prior to surgery or injection</p> <p style="padding: 5px;"><b>Warnings</b></p> <ul style="list-style-type: none"> <li>• For external use only</li> <li>• Flammable, keep away from fire or flame</li> <li>• Do not use with electrocautery procedures</li> </ul>	<b>Active ingredients</b>	<b>Purpose</b>	Chlorhexidine gluconate 3.15% w/v	Antiseptic	Isopropyl alcohol 70% v/v	Antiseptic	<p><b>Drug Facts (continued)</b></p> <p style="padding: 5px;"><b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash. If an allergic reaction occurs, stop use and seek medical help right away.</p> <p style="padding: 5px;"><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul>
<b>Active ingredients</b>	<b>Purpose</b>						
Chlorhexidine gluconate 3.15% w/v	Antiseptic						
Isopropyl alcohol 70% v/v	Antiseptic						
MISC027v00							

<p><b>Drug Facts (continued)</b></p> <p style="padding: 5px;"><b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p style="padding: 5px;"><b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p style="padding: 5px;"><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p style="padding: 5px;"><b>Directions</b></p> <ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)</li> <li>• tear open the pouch to expose the swabstick</li> <li>• remove the swabstick from package. Do not touch the foam tip.</li> <li>• prior to surgery or injection, apply swabstick to the procedure site</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>• place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>• allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul> <p style="padding: 5px;"><b>Other information</b></p> <ul style="list-style-type: none"> <li>• store between 68-77°F (20-25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul> <p style="padding: 5px;"><b>Inactive ingredient</b></p> <p style="padding: 5px;">USP purified water</p> <p style="padding: 5px;"><b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>
Each swabstick contains 0.054 fl. oz. (1.6 mL)	
Professional Disposables International, Inc. Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdihc.com	5QS324IS
MISC027v00	

5QS324IS\_2017\_07\_27.PDF

**5PS41950**

**Prevantics**  
NDC 10819-4075-4  
REORDER NO. S41950

**MAXI SWABSTICK**

**5PS41950**

**Prevantics**  
NDC 10819-4075-4  
REORDER NO. S41950

**MAXI SWABSTICK**

**5PS41950**

**Prevantics**  
NDC 10819-4075-4  
REORDER NO. S41950

**MAXI SWABSTICK**

**5PS41950\_2017\_07\_27.PDF**

Contains: 30 Individual Maxi Swabsticks  
0.172 fl. oz. (5.1 mL) Each

**MAXI SWABSTICK**

**MAXI SWABSTICK**

Drug Facts		Drug Facts (continued)	
<b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v Isopropyl alcohol 70% w/v	<b>Purpose</b> Antiseptic Antibiotic	<b>Directions</b> • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swabstick is approximately dry size 7 1/2 inches (19 cm) or moist size 3 1/2 by 7 5/8 inches (9.5 by 19 cm). • peel apart the pouch to expose the swabstick. • remove the swabstick from package. Do not touch the foam tip. • prior to surgery or injection, apply swabstick to the procedure site. • place one flat side of foam tip to the proposed skin site and wipe the skin in vigorous back-and-forth repeated strokes for one (1) minute. • turn the swabstick over (tipped side of the foam tip) and repeat the procedure by prepping for one (1) minute. • allow the prepped area to air dry for one and one half minutes (1 1/2 minutes). • do not blot or wipe dry. • discard after a single use.	<b>Warnings</b> • For external use only. • Flammable, keep away from fire or flame. • Do not use with electrocautery procedures.
<b>Use</b> For the preparation of the patient's skin prior to surgery or injection.	<b>Warnings</b> • For external use only. • Flammable, keep away from fire or flame. • Do not use with electrocautery procedures.	<b>Other information</b> • Store between 68°F to 77°F (20°-25°C). • Avoid freezing and excessive heat above 104°F (40°C) for hospital and professional use only.	<b>Questions?</b> call 1-800-999-4423, 24-Hr (9:00 am-5:00 pm EST)
<b>Warnings</b> • For external use only. • Flammable, keep away from fire or flame. • Do not use with electrocautery procedures.	<b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Itching • Rash If an allergic reaction occurs, stop use and seek medical help right away.	<b>Inactive ingredient</b> USP purified water.	
<b>Do not use</b> • on patients allergic to chlorhexidine gluconate or any other ingredients in this product. • on lacerated puncture or in contact with the membranes. • on open skin wounds or as a general skin cleanser. • under occlusive dressings.	<b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause irritation or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.		
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.			

Contains: 30 Individual Maxi Swabsticks  
0.172 fl. oz. (5.1 mL) Each



Contains: 30 Individual Maxi Swabsticks  
0.172 fl. oz. (5.1 mL) Each



**MAXI SWABSTICK**

**Prevantics**  
NDC 10819-4075-4  
REORDER NO. S41950

**MAXI SWABSTICK**

**PDI**

**CONTENTS:**  
30 packets of swabsticks premoistened with 5.1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (w/v) Isopropyl alcohol, and USP purified water.

**STORAGE**  
• store between 68-77°F [20-25°C]  
• avoid freezing and excessive heat above 104°F [40°C]

Do not use hypochlorite bleach on fabrics exposed to chlorhexidine gluconate.

LOT  
EXP.

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]  
**PATIENT PREOPERATIVE SKIN PREPARATION**  
**Non-sterile Solution**  
\*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

- FOR EXTERNAL USE ONLY**
- Single step, broad spectrum, persistent antiseptic that significantly reduces the number of microorganisms on intact skin
  - Use for the preparation of the patient's skin prior to surgery or injection
  - Helps reduce bacteria that potentially can cause skin infection
  - Latex free
  - For hospital and professional use only

NDC 10819-4075-4 REORDER NO. S41950

**Prevantics**<sup>®</sup>

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]

**PATIENT PREOPERATIVE SKIN PREPARATION**

**Non-sterile Solution**

\*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

**FOR EXTERNAL USE ONLY**

- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only • Latex free

**IMPORTANT:** See carton for complete **Drug Facts** and safety information.

**One Maxi Swabstick, 0.172 fl. oz. (5.1 mL) Each**



**MAXI SWABSTICK**



**CONTENTS:** One Maxi Swabstick premoistened with 5.1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water.

**DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one maxi swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm). Peel apart the pouch to expose the swabstick. Remove swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one and one half minutes (1.5 minutes). Do not blot or wipe dry. Discard after a single use.

**WARNINGS**

**For external use only. Flammable, keep away from fire or flame.** Do not use with electrocautery procedures.

**Allergy Alert**

This product may cause a severe allergic reaction. Symptoms include: • Wheezing/difficulty breathing  
• Shock • Facial swelling • Hives • Rash

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

Professional Disposables International, Inc.  
Orangeburg, NY 10962-1376 USA  
1-800-999-6423 www.pdihc.com



(01)00310819008419



LOT →

THIS SIDE UP  
PEEL APART HERE

EXP. →

2PS419

2PS419\_2017\_03\_06.pdf

Prevanics Maxi Swabstick secondary carton Ten 30-count

**NET CONTENTS: 10 BOXES OF 30 INDIVIDUAL SWABSTICKS**  
**REORDER NO. S41950**  
**NDC 10819-4075-4**

**†Prevanics® Maxi Swabstick**  
**[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]**  
**Patient preoperative skin preparation**  
**Non-sterile Solution**  
 \*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

<p><b>Drug Facts</b></p> <p><b>Active ingredients</b></p> <p>Chlorhexidine gluconate 3.15% w/v ..... Antiseptic                  Isopropyl alcohol 70% v/v ..... Antiseptic</p> <p><b>Purpose</b></p> <p>Antiseptic</p> <p><b>Use</b></p> <p>For the preparation of the patient's skin prior to surgery or injection</p> <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>• For external use only</li> <li>• Flammable, keep away from fire or flame</li> <li>• Do not use with electrocautery procedures</li> </ul> <p><b>Allergy Alert</b></p> <p>This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash                  If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul> <p><b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p><b>Stop use and ask a doctor if</b> irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)</li> <li>• peel apart the pouch to expose the swabstick</li> <li>• remove the swabstick from package. Do not touch the foam tip.</li> <li>• prior to surgery or injection, apply swabstick to the procedure site</li> <li>• place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>• allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>• store between 68-77°F (20-25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul> <p><b>Inactive ingredient</b></p> <p>USP purified water</p> <p><b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>
---	--

**Each Maxi Swabstick contains 0.172 fl. oz. (5.1 mL)**

Professional Disposables International, Inc.  
 Orangeburg, NY 10962-1376 USA  
 1-800-999-6423 www.pdihc.com

(01)20310819008413

8NS41950

LBL878v02

8NS41950\_2017\_07\_27.PDF

Prevantics Maxi Swabstick bulk outer carton 300-count

<b>Prevantics® Maxi Swabstick</b> [Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab] Patient preoperative skin preparation Non-sterile Solution Packaging is sterile when in sterile kit		REORDER NO. S27350 NDC 10819-4075-3   (01)20310819000882																																									
*equivalent to 855 mg of Chlorhexidine Gluconate per pouch <b>Contains: 300 Individual Swabsticks, 0.172 fl. oz. (5.1 mL) Each</b>																																											
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td colspan="2"><b>Drug Facts</b></td> </tr> <tr> <td style="width: 50%;"><b>Active ingredients</b></td> <td style="width: 50%;"><b>Purpose</b></td> </tr> <tr> <td>Chlorhexidine gluconate 3.15% w/v.....</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v.....</td> <td>Antiseptic</td> </tr> <tr> <td colspan="2"><b>Use</b> For the preparation of the patient's skin prior to surgery or injection</td> </tr> <tr> <td colspan="2"><b>Warnings</b></td> </tr> <tr> <td colspan="2"> <ul style="list-style-type: none"> <li>• For external use only</li> <li>• <b>Flammable, keep away from fire or flame</b></li> <li>• Do not use with electrocautery procedures</li> </ul> </td> </tr> <tr> <td colspan="2"> <b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash                      If an allergic reaction occurs, stop use and seek medical help right away.                 </td> </tr> <tr> <td colspan="2"><b>Do not use</b></td> </tr> <tr> <td colspan="2"> <ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul> </td> </tr> <tr> <td colspan="2"> <b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.                 </td> </tr> <tr> <td colspan="2"> <b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.                 </td> </tr> <tr> <td colspan="2"> <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.                 </td> </tr> </table>	<b>Drug Facts</b>		<b>Active ingredients</b>	<b>Purpose</b>	Chlorhexidine gluconate 3.15% w/v.....	Antiseptic	Isopropyl alcohol 70% v/v.....	Antiseptic	<b>Use</b> For the preparation of the patient's skin prior to surgery or injection		<b>Warnings</b>		<ul style="list-style-type: none"> <li>• For external use only</li> <li>• <b>Flammable, keep away from fire or flame</b></li> <li>• Do not use with electrocautery procedures</li> </ul>		<b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash If an allergic reaction occurs, stop use and seek medical help right away.		<b>Do not use</b>		<ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul>		<b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.		<b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.		<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.		<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td colspan="2"><b>Drug Facts</b> (continued)</td> </tr> <tr> <td colspan="2"><b>Directions</b></td> </tr> <tr> <td colspan="2"> <ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)</li> <li>• tear open the pouch to expose the swabstick</li> <li>• remove the swabstick from package. Do not touch the foam tip.</li> <li>• prior to surgery or injection, apply swabstick to the procedure site</li> <li>• place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>• allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul> </td> </tr> <tr> <td colspan="2"><b>Other information</b></td> </tr> <tr> <td colspan="2"> <ul style="list-style-type: none"> <li>• store between 68 – 77°F (20 – 25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul> </td> </tr> <tr> <td colspan="2"><b>Inactive ingredient</b></td> </tr> <tr> <td colspan="2">USP purified water</td> </tr> <tr> <td colspan="2"><b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</td> </tr> </table>	<b>Drug Facts</b> (continued)		<b>Directions</b>		<ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)</li> <li>• tear open the pouch to expose the swabstick</li> <li>• remove the swabstick from package. Do not touch the foam tip.</li> <li>• prior to surgery or injection, apply swabstick to the procedure site</li> <li>• place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>• allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul>		<b>Other information</b>		<ul style="list-style-type: none"> <li>• store between 68 – 77°F (20 – 25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul>		<b>Inactive ingredient</b>		USP purified water		<b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	
<b>Drug Facts</b>																																											
<b>Active ingredients</b>	<b>Purpose</b>																																										
Chlorhexidine gluconate 3.15% w/v.....	Antiseptic																																										
Isopropyl alcohol 70% v/v.....	Antiseptic																																										
<b>Use</b> For the preparation of the patient's skin prior to surgery or injection																																											
<b>Warnings</b>																																											
<ul style="list-style-type: none"> <li>• For external use only</li> <li>• <b>Flammable, keep away from fire or flame</b></li> <li>• Do not use with electrocautery procedures</li> </ul>																																											
<b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash If an allergic reaction occurs, stop use and seek medical help right away.																																											
<b>Do not use</b>																																											
<ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul>																																											
<b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.																																											
<b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.																																											
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.																																											
<b>Drug Facts</b> (continued)																																											
<b>Directions</b>																																											
<ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)</li> <li>• tear open the pouch to expose the swabstick</li> <li>• remove the swabstick from package. Do not touch the foam tip.</li> <li>• prior to surgery or injection, apply swabstick to the procedure site</li> <li>• place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>• allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul>																																											
<b>Other information</b>																																											
<ul style="list-style-type: none"> <li>• store between 68 – 77°F (20 – 25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul>																																											
<b>Inactive ingredient</b>																																											
USP purified water																																											
<b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)																																											
Professional Disposables International, Inc., Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdihc.com		8SS27350  																																									

8QB11400\_2017\_07\_27.PDF

<b>REORDER NO. S27350</b> <b>NDC 10819-4075-3</b>							
<h1 style="margin: 0;">+Prevantics® Maxi Swabstick</h1> <p style="margin: 0;">[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]                  Patient preoperative skin preparation                  Non-sterile Solution                  Packaging is sterile when in sterile kit                  *equivalent to 855 mg of Chlorhexidine Gluconate per pouch</p>							
<p><b>Drug Facts</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="text-align: left; padding: 2px;">Active ingredients</th> <th style="text-align: left; padding: 2px;">Purpose</th> </tr> <tr> <td style="padding: 2px;">Chlorhexidine gluconate 3.15% w/v</td> <td style="padding: 2px;">Antiseptic</td> </tr> <tr> <td style="padding: 2px;">Isopropyl alcohol 70% v/v</td> <td style="padding: 2px;">Antiseptic</td> </tr> </table> <p><b>Use</b> For the preparation of the patient's skin prior to surgery or injection</p> <p><b>Warnings</b></p> <ul style="list-style-type: none"> <li>• For external use only</li> <li>• Flammable, keep away from fire or flame</li> <li>• Do not use with electrocautery procedures</li> </ul> <p><b>Allergy Alert</b>                  This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing ▶</p>	Active ingredients	Purpose	Chlorhexidine gluconate 3.15% w/v	Antiseptic	Isopropyl alcohol 70% v/v	Antiseptic	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>• Shock • Facial swelling • Hives • Rash</li> </ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>• on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>• for lumbar puncture or in contact with the meninges</li> <li>• on open skin wounds or as a general skin cleanser</li> <li>• under occlusive patch</li> </ul> <p><b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ▶</p>
Active ingredients	Purpose						
Chlorhexidine gluconate 3.15% w/v	Antiseptic						
Isopropyl alcohol 70% v/v	Antiseptic						

MISC027v00

<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>• use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>• maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)</li> <li>• tear open the pouch to expose the swabstick</li> <li>• remove the swabstick from package. Do not touch the foam tip.</li> <li>• prior to surgery or injection, apply swabstick to the procedure site ▶</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>• place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>• turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>• allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>• do not blot or wipe dry</li> <li>• discard after a single use</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>• store between 68-77°F (20-25°C)</li> <li>• avoid freezing and excessive heat above 104°F (40°C)</li> <li>• for hospital and professional use only</li> </ul> <p><b>Inactive ingredient</b>                  USP purified water</p> <p><b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>
--	--

**Each Maxi Swabstick contains 0.172 fl. oz. (5.1 mL)**

Professional Disposables International, Inc. Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdihc.com	5QS273IS			
---	----------	--	--	--

MISC027v00

5QS273IS\_2017\_07\_27.PDF

NDC 10819-4075-3

REORDER NO. S27350

**Prevantics**<sup>®</sup>

[Chlorhexidine Gluconate (3.15%)\* and Isopropyl Alcohol (70%) Swab]

**PATIENT PREOPERATIVE SKIN PREPARATION**

**Non-sterile Solution**

**Packaging is sterile when in sterile kit**

\*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

**FOR EXTERNAL USE ONLY**

- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only • Latex free

**IMPORTANT:** See insert for complete **Drug Facts** and safety information.

**One Maxi Swabstick, 0.172 fl.oz. (5.1 mL) Each**



**MAXI SWABSTICK**



**CONTENTS:** One Maxi Swabstick premoistened with 5.1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water.

**DIRECTIONS:** Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one maxi swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm). Tear open the pouch to expose the swabstick. Remove swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one and one half minutes (1.5 minutes). Do not blot or wipe dry. Discard after a single use.

**WARNINGS: For external use only. Flammable, keep away from fire or flame.** Do not use with electrocautery procedures. **Allergy Alert** This product may cause a severe allergic reaction. Symptoms include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

Professional Disposables International, Inc.  
Orangeburg, NY 10962-1376 USA  
1-800-999-6423 www.pdihc.com



(01)00310819000888








LOT

EXP  
THIS SIDE UP  
TEAR OPEN HERE

2QS273

2QS273\_2017\_MAR\_01.ai

		REORDER NO. S42850 NDC 10819-4075-5	
[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab] Patient preoperative skin preparation Non-sterile Solution Packaging is sterile when in sterile kit			
		 (01)20310819000998	
*equivalent to 268 mg of Chlorhexidine Gluconate per pouch <b>Contains: 500 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each</b>			
<b>Drug Facts</b> <b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v ..... Antiseptic Isopropyl alcohol 70% v/v ..... Antiseptic <b>Purpose</b> Antiseptic <b>Use</b> For the preparation of the patient's skin prior to surgery or injection <b>Warnings</b> <ul style="list-style-type: none"> <li>For external use only</li> <li>Flammable, keep away from fire or flame</li> <li>Do not use with electrocautery procedures</li> </ul> <b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing • Shock • Facial swelling • Hives • Rash. If an allergic reaction occurs, stop use and seek medical help right away. <b>Do not use</b> <ul style="list-style-type: none"> <li>on patients allergic to chlorhexidine gluconate or any other ingredient in this product</li> <li>for lumbar puncture or in contact with the meninges</li> <li>on open skin wounds or as a general skin cleanser</li> <li>under occlusive patch</li> </ul> <b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. <b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.		<b>Drug Facts (continued)</b> <b>Directions</b> <ul style="list-style-type: none"> <li>use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> <li>maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)</li> <li>tear open the pouch to expose the swabstick</li> <li>remove the swabstick from package. Do not touch the foam tip.</li> <li>prior to surgery or injection, apply swabstick to the procedure site</li> <li>place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute</li> <li>turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute</li> <li>allow the prepped area to air dry for one and one half minutes (1.5 minutes)</li> <li>do not blot or wipe dry</li> <li>discard after a single use</li> </ul> <b>Other information</b> <ul style="list-style-type: none"> <li>store between 68-77°F (20-25°C)</li> <li>avoid freezing and excessive heat above 104°F (40°C)</li> <li>for hospital and professional use only</li> </ul> <b>Inactive ingredient</b> USP purified water <b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)	
 Professional Disposables International, Inc., Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdihc.com		 	

8RS428CK

LBL878v02

8RS428CK\_2017\_07\_27.PDF

This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

<p>NDC 10819-4075-5    REORDER NO. S42850</p> <p><b>Prevantics</b><sup>®</sup></p> <p>[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]</p> <p>(01)00310819000994</p> <p><b>PATIENT PREOPERATIVE SKIN PREPARATION</b> Non-sterile Solution</p> <p><b>Packaging is sterile when in sterile kit</b> *equivalent to 268 mg of Chlorhexidine Gluconate per pouch</p> <p><b>FOR EXTERNAL USE ONLY</b> • Use for the preparation of the patient's skin prior to surgery or injection • Helps reduce bacteria that potentially can cause skin infection • For hospital and professional use only • Latex free</p> <p><b>IMPORTANT:</b> See insert for complete <i>Drug Facts</i> and safety information. <b>One Swabstick</b>, 0.054 fl. oz. (1.6 mL) Each</p>	 <p><b>SWABSTICK</b></p> 	<p><b>TEAR OPEN HERE</b></p> <p><b>CONTENTS:</b> One swabstick premoistened with 1.6 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water. <b>DIRECTIONS:</b> Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm). Tear open the pouch to expose the swabstick. Remove the swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply swabstick to the procedure site. Place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute. Turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute. Allow the prepped area to air dry for one and one half minutes (1.5 minutes). Do not blot or wipe dry. Discard after a single use. <b>WARNINGS: For external use only. Flammable, keep away from fire or flame.</b> Do not use with electrocautery procedures. <b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms include: • wheezing/difficulty breathing • shock • facial swelling • hives • rash. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Professional Disposables International, Inc. Orangeburg, NY 10962-1376 USA <b>PDI</b> 1-800-999-6423 • <a href="http://www.pdihc.com">www.pdihc.com</a></p>	<p>LOT</p> <p>EXP:</p> 
		<p>2QS428</p>	




2QS428\_2017\_MAR\_01.ai

<b>REORDER NO. S42850</b> <b>NDC 10819-4075-5</b>	
<b>†Prevantics® Swabstick</b>	
<b>[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]</b>	
<b>Patient preoperative skin preparation</b>	
<b>Non-sterile Solution</b>	
<b>Packaging is sterile when in sterile kit</b>	
*equivalent to 268 mg of Chlorhexidine Gluconate per pouch	
<b>Drug Facts</b> <hr/> <b>Active ingredients</b> Chlorhexidine gluconate 3.15% w/v ..... Antiseptic Isopropyl alcohol 70% v/v ..... Antiseptic <hr/> <b>Purpose</b> Antiseptic <hr/> <b>Use</b> For the preparation of the patient's skin prior to surgery or injection <hr/> <b>Warnings</b> • For external use only • Flammable, keep away from fire or flame • Do not use with electrocautery procedures <hr/> <b>Allergy Alert</b> This product may cause a severe allergic reaction. Symptoms may include: • Wheezing/difficulty breathing	<b>Drug Facts (continued)</b> <hr/> • Shock • Facial swelling • Hives • Rash If an allergic reaction occurs, stop use and seek medical help right away. <hr/> <b>Do not use</b> • on patients allergic to chlorhexidine gluconate or any other ingredient in this product • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch <hr/> <b>When using this product</b> keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

MISC027v00

<b>Drug Facts (continued)</b> <hr/> <b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. <hr/> <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away. <hr/> <b>Directions</b> • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm) • tear open the pouch to expose the swabstick • remove the swabstick from package. Do not touch the foam tip. • prior to surgery or injection, apply swabstick to the procedure site	<b>Drug Facts (continued)</b> <hr/> • place one flat side of foam tip to the proposed skin site and prep the skin in <b>vigorous</b> back-and-forth repeated strokes for one (1) minute • turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute • allow the prepped area to air dry for one and one half minutes (1.5 minutes) • do not blot or wipe dry • discard after a single use <hr/> <b>Other information</b> • store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only <hr/> <b>Inactive ingredient</b> USP purified water <hr/> <b>Questions?</b> call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)
--	---

**Each swabstick contains 0.054 fl. oz. (1.6 mL)**

 Professional Disposables International, Inc. Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdihc.com	5QS428IS  
---	--

MISC027v00