

NDC 10819-4076-4 REORDER NO. S41950

Prevantics[®]

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]

PATIENT PREOPERATIVE SKIN PREPARATION
Non-sterile Solution

*equivalent to 161 mg of Chlorhexidine Gluconate per pouch

FOR EXTERNAL USE ONLY

- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only • Latex free

IMPORTANT: See carton for complete **Drug Facts** and safety information.

One Maxi Swabstick, 0.172 fl. oz. (5.1 mL) Each **PDI**[®]

MAXI SWABSTICK

WARNING
FLAMMABLE



Keep away from fire or flame.

To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas.
- Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

CONTENTS: One Maxi Swabstick premoistened with 5.1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water.

DIRECTIONS: Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swabstick is approximately 8.4 in. x 8.4 in. (21.3 cm by 21.3 cm). Peel apart the pouch to expose the swabstick. Remove swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes, turning the swabstick over (unused side of the foam tip) halfway in the procedure.

dry site: Scrub repeatedly back and forth for 15 seconds with each side of the applicator for a total of 30 seconds. Allow to air dry for approximately 30 seconds. **moist site:** Scrub repeatedly for 1 minute with each side of applicator for a total of 2 minutes. Allow to air dry for approximately 1.5 minutes. • **if using an ignition source, allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. • discard after a single use

WARNINGS: For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire,

PREP CAREFULLY: • solution contains alcohol and gives off **flammable vapors** • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.** • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area

Allergy alert: This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing • shock • facial swelling • hives • rash

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Professional Disposables International, Inc.
Orangeburg, NY 10962-1376 USA
1-800-999-6423 www.pdihc.com



(01)00310819008419

2TS419



↑ PEEL ↑

↑ PEEL ↑

LOT

EXP.

0.172 fl. oz. (5.1 mL) Each
Contains: 30 Individual Maxi Swabsticks

PDI

MAXI SWABSTICK

PATIENT PREOPERATIVE SKIN PREPARATION
Non-sterile Solution

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]

Prevantics

NDC 10819-4076-4
REORDER NO. S41950

NDC 10819-4076-4
REORDER NO. S41950

TEAR ALONG PERFORATION FOR DISPENSING

Drug Facts	Drug Facts (continued)
<p>Active ingredients Chlorhexidine gluconate 3.15% w/v Isopropyl alcohol 70% v/v</p> <p>Purpose Antiseptic Antisepctic</p> <p>Use For the preparation of the patient's skin prior to surgery or injection</p> <p>Warnings For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY. • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area</p> <p>Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficulty breathing • shock • facial swelling • hives • rash If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Do not use • on patients allergic to chlorhexidine gluconate or any other ingredient in this product • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch</p> <p>When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause dizziness or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p>	<p>Step use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control center right away.</p> <p>Directions • Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • Maximum treatment area for one application is approximately 8.4 in. x 8.4 in. (21.3 cm by 21.3 cm). • Tilt head back to expose the swabstick. • Remove swabstick from package. Do not touch the foam tip. • Prior to surgery or injection, apply over the side of the head to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes, turning the swabstick over (smooth side of the foam tip) halfway in the procedure. • dry site. Repeat back and forth for 15 seconds with each side of the applicator for a total of 30 seconds. Allow hair dry for approximately 30 seconds. • wash site. Repeat for 1 minute with each side of applicator for a total of 2 minutes. Allow to air dry for approximately 15 minutes. • If sites on hairy areas, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not sit or sleep.</p> <p>Other information • Store between 68-77°F (20-25°C) • avoid freezing and excessive heat above 104°F (40°C) • see package insert for complete information.</p> <p>Inactive ingredient USP purified water</p> <p>Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)</p>

5SS41950

Prevantics

MAXI SWABSTICK

NDC 10819-4076-4
REORDER NO. S41950

Prevantics

NDC 10819-4076-4
REORDER NO. S41950

Prevantics

NDC 10819-4076-4
REORDER NO. S41950

Prevantics

NDC 10819-4076-4
REORDER NO. S41950

MAXI SWABSTICK

PDI

MAXI SWABSTICK

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
PATIENT PREOPERATIVE SKIN PREPARATION
Non-sterile Solution
*Equivalent to 161mg of Chlorhexidine Gluconate per pouch

FOR EXTERNAL USE ONLY

- Single step, broad spectrum, persistent antiseptic that significantly reduces the number of microorganisms on intact skin
- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- Latex free
- For hospital and professional use only

⚠ WARNING
FLAMMABLE

Keep away from fire or flame.
To reduce the risk of fire, PREP CAREFULLY:
• solution contains alcohol and gives off flammable vapors
• avoid getting solution into hairy areas.
• Hair may take up to 1 hour to dry. Wet hair is flammable.
• do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
• do not allow solution to pool
• remove wet materials from prep area

CONTENTS:
30 packets of swabsticks premoistened with 5.1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water

STORAGE
• store between 68-77°F (20-25°C)
• avoid freezing and excessive heat above 104°F (40°C)


Do not use hypochlorite bleach on fabrics exposed to chlorhexidine gluconate

LOT

EXP.

Contains: 30 Individual Maxi Swabsticks
0.172 fl. oz. (5.1 mL) Each

PDI



(01)10310819008416

Contains: 30 Individual Maxi Swabsticks
0.172 fl. oz. (5.1 mL) Each

PDI

NDC 10819-4076-3

REORDER NO. S27350

Prevantics[®]

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]

PATIENT PREOPERATIVE SKIN PREPARATION

Non-sterile Solution

Packaging is sterile when in sterile kit

*equivalent to 161 mg of Chlorhexidine Gluconate per pouch

FOR EXTERNAL USE ONLY

- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- For hospital and professional use only • Latex free

IMPORTANT: See insert for complete **Drug Facts** and safety information.

One Maxi Swabstick, 0.172 fl.oz. (5.1 mL) Each



MAXI SWABSTICK

**WARNING
FLAMMABLE**



Keep away from fire or flame.

To reduce the risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas.
- Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

CONTENTS: One Maxi Swabstick premoistened with 5.1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water.

DIRECTIONS: Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. Maximum treatment area for one swabstick is approximately 8.4 in. x 8.4 in. (21.3 cm by 21.3 cm). Tear open the pouch to expose the swabstick. Remove swabstick from package. Do not touch the foam tip. Prior to surgery or injection, apply one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes, turning the swabstick over (unused side of the foam tip) halfway in the procedure.

dry site: Scrub repeatedly back and forth for 15 seconds with each side of the applicator for a total of 30 seconds. Allow to air dry for approximately 30 seconds. **moist site:** Scrub repeatedly for 1 minute with each side of applicator for a total of 2 minutes. Allow to air dry for approximately 1.5 minutes. • **if using an ignition source, allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. • discard after a single use

WARNINGS: For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire,

PREP CAREFULLY: • solution contains alcohol and gives off **flammable vapors** • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.** • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area

Allergy alert: This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing • shock • facial swelling • hives • rash

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Professional Disposables International, Inc.
Orangeburg, NY 10962-1376 USA
1-800-999-6423 www.pdihc.com



(01)00310819000888



EXP.

THIS SIDE UP
TEAR OPEN HERE

LOT

2US273

REORDER NO. S27350 NDC 10819-4076-3							
<h1 style="margin: 0;">Prevantics®</h1> <h2 style="margin: 0;">Maxi Swabstick</h2> <p>[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab] Patient preoperative skin preparation Non-sterile Solution Packaging is sterile when in sterile kit *equivalent to 161mg of Chlorhexidine Gluconate per pouch</p>	<div style="background-color: #e91e63; color: white; text-align: center; padding: 5px; font-weight: bold;"> ⚠ WARNING FLAMMABLE </div> <div style="display: flex; align-items: center; margin-top: 10px;"> <div style="font-size: small;"> <p>Keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> solution contains alcohol and gives off flammable vapors avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) do not allow solution to pool remove wet materials from prep area </div> </div>						
Drug Facts	Drug Facts (continued)						
<table border="1" style="width: 100%; border-collapse: collapse; font-size: x-small;"> <tr> <td style="width: 50%;">Active ingredients</td> <td style="width: 50%;">Purpose</td> </tr> <tr> <td>Chlorhexidine gluconate 3.15% w/v.....</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v.....</td> <td>Antiseptic</td> </tr> </table>	Active ingredients	Purpose	Chlorhexidine gluconate 3.15% w/v.....	Antiseptic	Isopropyl alcohol 70% v/v.....	Antiseptic	solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet materials from prep area
Active ingredients	Purpose						
Chlorhexidine gluconate 3.15% w/v.....	Antiseptic						
Isopropyl alcohol 70% v/v.....	Antiseptic						
Use For the preparation of the patient's skin prior to surgery or injection	Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficulty breathing • shock • facial swelling • hives • rash If an allergic reaction occurs, stop use and seek medical help right away.						
Warnings For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY: • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until	Do not use • on patients allergic to chlorhexidine gluconate or any other ingredient in this product						

Drug Facts (continued) • for lumbar puncture or in contact with the meninges • on open skin wounds or as a general skin cleanser • under occlusive patch When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	Drug Facts (continued) • dry site: Scrub repeatedly back and forth for 15 seconds with each side of the applicator for a total of 30 seconds. Allow to air dry for approximately 30 seconds. • moist site: Scrub repeatedly for 1 minute with each side of applicator for a total of 2 minutes. Allow to air dry for approximately 1.5 minutes. • if using an ignition source, allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. • discard after a single use
Directions • use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. • maximum treatment area for one swabstick is approximately 8.4 in. x 8.4 in. (21.3 cm by 21.3 cm). • tear open the pouch to expose the swabstick. • remove swabstick from package. Do not touch the foam tip. • prior to surgery or injection, apply one flat side of foam tip to the proposed skin site and prep the skin in vigorous back-and-forth repeated strokes, turning the swabstick over (unused side of the foam tip) halfway in the procedure.	Other information • store between 68 – 77°F (20 – 25°C) • avoid freezing and excessive heat above 104°F (40°C) • for hospital and professional use only Inactive ingredient USP purified water Questions? call 1-800-999-6423 , M-F (9:00 am-5:00 pm EST)

Each Maxi Swabstick contains 0.172 fl. oz. (5.1 mL)

Professional Disposables International, Inc.
 Orangeburg, NY 10962-1376 USA
 1-800-999-6423 www.pdihc.com

5TS273IS

REORDER NO. S41950
NDC 10819-4076-4

Prevantics[®] Maxi Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution

*equivalent to 161mg of Chlorhexidine Gluconate per pouch

Contains: 10 Boxes of 30 Individual Swabsticks, 0.172 fl. oz. (5.1 mL) Each

STORE BETWEEN 68 – 77°F (20 – 25°C)
AVOID FREEZING AND EXCESSIVE HEAT ABOVE 104°F (40°C)



WARNING, FLAMMABLE



Keep away from fire or flame.



(01)20310819008413

8QS41950



Professional Disposables International, Inc., Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdihc.com



REORDER NO. S27350
NDC 10819-4076-3

Prevantics[®] Maxi Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]

Patient preoperative skin preparation

Non-sterile Solution

Packaging is sterile when in sterile kit

*equivalent to 161mg of Chlorhexidine Gluconate per pouch

Contains: 300 Individual Swabsticks, 0.172 fl.oz. (5.1 mL) Each

STORE BETWEEN 68 – 77°F (20 – 25°C)
AVOID FREEZING AND EXCESSIVE HEAT ABOVE 104°F (40°C)



WARNING, FLAMMABLE



Keep away from fire or flame.



(01)20310819000882

8US27350



Professional Disposables International, Inc., Orangeburg, NY 10962-1376 USA 1-800-999-6423 www.pdihc.com



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

FRANCIS E BECKER
09/20/2021 02:54:43 PM