

NDC 054365-400-15

1 ml APPLICATORS

Clear


Single Use
Latex Free
Applicator is STERILE
if package is intact

20 applicators
0.03 fl. oz. (1 ml) each

**CloraPrep[®]
One-Step**

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

Patient Preoperative
Skin Preparation



**WARNING
FLAMMABLE**

- Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:
- solution contains alcohol and gives off flammable vapors
 - avoid getting solution into hairy areas.
 - Hair may take up to 1 hour to dry. **Wet hair is flammable.**
 - do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hr in hair)
 - do not allow solution to pool
 - remove wet material from prep area

Store between 15-30 °C (59-86 °F)
Avoid freezing and excessive heat
above 40 °C (104 °F)

**CloraPrep[®]
One-Step**

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

1 ml APPLICATORS

Clear

LOT XXXXXX EXP DATE: XXXX
(printed in-line)



Cat. No. 260479
©2009 CareFusion. All rights reserved. CloraPrep is a registered trademark of CareFusion or one of its subsidiaries.

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients
Chlorhexidine gluconate 2% w/v.....
Isopropyl alcohol 70% v/v.....

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

- Warnings**
- For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:
 - solution contains alcohol and gives off flammable vapors
 - avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
 - do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
 - do not allow solution to pool
 - remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product, keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Purposes

Antiseptic

Antiseptic

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²).
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair).
- Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients • USP purified water

Questions? • www.chloraprep.com
• call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

(b) (4)

ChloroPrep 1 mL Cyrel

(b) (4)

(b) (4)



Trim Area

<div style="background-color: #f08080; text-align: center; padding: 2px;">⚠ WARNING</div> <div style="background-color: #f08080; text-align: center; padding: 2px;">FLAMMABLE</div> <p>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet material from prep area 	<h2 style="margin: 0;">ChloroPrep® One-Step</h2> <p style="margin: 0;">2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation</p> <p style="text-align: center; margin: 0;">1 ml APPLICATOR</p> <p style="margin: 0;">Single Use <input checked="" type="checkbox"/> Do Not Reuse <input checked="" type="checkbox"/> Latex Free</p> <p style="margin: 0;">External Use Only Professional Use Only</p> <p style="margin: 0;">Cat. No. 260479 NDC 054365-400-11</p>						
<h3 style="margin: 0;">Drug Facts</h3> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 70%;">Active ingredients</td> <td style="width: 30%;">Purposes</td> </tr> <tr> <td>Chlorhexidine gluconate 2% w/v.....</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v.....</td> <td>Antiseptic</td> </tr> </table> <p>Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. (11)</p> <p>Warnings</p> <p>For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area <p>Do not use</p> <ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser <p>When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ▶</p>		Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....	Antiseptic	Isopropyl alcohol 70% v/v.....	Antiseptic
Active ingredients	Purposes						
Chlorhexidine gluconate 2% w/v.....	Antiseptic						
Isopropyl alcohol 70% v/v.....	Antiseptic						


ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ use in a well ventilated area ■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic. ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes ■ allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. <p>Other information</p> <ul style="list-style-type: none"> ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) <p>Inactive ingredients</p> <ul style="list-style-type: none"> ■ USP purified water <p>Questions?</p> <p>■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)</p>

CareFusion, Leawood, KS 66211 • Artwork 6-775479• A0611 • Made in the USA of US and Imported parts

8
Chloraprep
 one-step



WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/vAntiseptic
 Isopropyl alcohol 70% v/vAntiseptic

Purposes

Antiseptic
 Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator horizontally with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- USP purified water

Questions?

- www.chloraprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

(b) (4)

72 mm

40 mm

Exp. Date xx/xx
Lot No. xxxxx

Cat. No. 260299 NDC 054365-400-02

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
1.5ml FREPP® APPLICATOR
Single Use • Latex Free

 **WARNING. FLAMMABLE.**
KEEP AWAY FROM FIRE OR FLAME.

**do not drape or use ignition source (e.g., cautery, laser)
until solution is completely dry (minimum of 3 minutes
on hairless skin; up to 1 hour in hair)**

APPLICATOR STERILE UNLESS SEAL IS BROKEN

CareFusion, Leewood, KS 66211
Artwork No. 6-000299 • K0611
Made in the USA of US and imported parts

Bend and Peel

(b) (4) ChloroPrep 2% Frepp Applicator, Non-Sterile - Folded Drug Facts Insert

(b) (4)

(b) (4)

7.5"

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
 1.5ml FREPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic

Use For the preparation of the patient's skin prior to surgery; Helps to reduce bacteria that potentially can cause skin infection.

Warnings
 For external use only. Flammable: Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator horizontally with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15–30 °C (59–86 °F)
 ■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloroprep.com
 ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

Reference ID: 3175064
 See other side for additional information

Front

Back

2.5"

ChloroPrep® One-Step
 2% w/v chlorhexidine gluconate (CHG)
 and 70% v/v isopropyl alcohol (IPA)
 Patient Preoperative Skin Preparation
 1.5ml FREPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

**⚠ WARNING
 FLAMMABLE**

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area


See Drug Facts for Full Labeling Information

Do Not Reuse Latex Free

External Use Only Professional Use Only


Single Use

CareFusion, Leawood, KS 66211
 Artwork 6-775403F - L0411
 Made in the USA of US and imported parts




(b) (4) ChloraPrep 3ml Applicator lidding

(b) (4)



<div style="background-color: #f08080; text-align: center; padding: 2px;">⚠ WARNING</div> <div style="background-color: #f08080; text-align: center; padding: 2px;">FLAMMABLE</div> <p>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet material from prep area 	<h2 style="margin: 0;">ChloraPrep[®] One-Step ChloraPrep[®] With Tint</h2> <p style="margin: 0;">2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation</p> <p style="text-align: center; font-weight: bold; font-size: 1.2em;">3 ml APPLICATOR</p> <p style="margin: 0;">Single Use <input checked="" type="checkbox"/> Do Not Reuse <input checked="" type="checkbox"/> Latex Free</p> <p style="margin: 0;">External Use Only Professional Use Only</p> <p style="margin: 0;">Cat. No. 260415 / 260415NS / 260415NSB NDC 054365-400-11 Cat. No. 260400 / 260400NS / 260500NS NDC 054365-400-01</p>						
<h3 style="margin: 0;">Drug Facts</h3> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 70%;">Active ingredients</td> <td style="width: 30%;">Purposes</td> </tr> <tr> <td>Chlorhexidine gluconate 2% w/v.....</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v.....</td> <td>Antiseptic</td> </tr> </table> <p style="margin: 5px 0;">Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. 11</p> <p>Warnings</p> <p>For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area <p>Do not use</p> <ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser <p>When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ▶</p>		Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....	Antiseptic	Isopropyl alcohol 70% v/v.....	Antiseptic
Active ingredients	Purposes						
Chlorhexidine gluconate 2% w/v.....	Antiseptic						
Isopropyl alcohol 70% v/v.....	Antiseptic						

<h2 style="margin: 0;">ChloraPrep[®] One-Step / ChloraPrep[®] With Tint</h2> <p style="margin: 0;">2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)</p>
<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ use in a well ventilated area ■ maximal treatment area for one applicator is approximately 4 in. x 5 in. (130 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings <u>only once</u> to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic. ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes ■ allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. <p>Other information</p> <ul style="list-style-type: none"> ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ The tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired. 9 <p>Inactive ingredients</p> <ul style="list-style-type: none"> ■ FD&C green #3 dye (Scrub-Teal applicators only) ■ FD&C yellow #6 dye (Hi-Lite Orange applicators only) ■ USP purified water <p>Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)</p>
<p style="font-size: 0.8em;">CareFusion, Leawood, KS 66211 • Artwork 6-775400- E0611 • Made in the USA of US and imported parts</p>



$1\frac{5}{32}$

$3\frac{1}{16}$

$\frac{1}{4}$

2

$5\frac{5}{8}$

$3\frac{7}{32}$

$12\frac{7}{32}$

$5\frac{9}{32}$

$2\frac{5}{64}$

$6\frac{1}{2}$

$3\frac{1}{4}$

$1\frac{7}{4}$

CloraPrep
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)

A G A B P WAY O O A

CloraPrep
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
3 ml
PP
S T *

CloraPrep
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
3 ml
PP
S b T *

3 ml
P
S T *

CloraPrep
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
P P v P
**WARNING
FLAMMABLE**


CloraPrep
With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
P P v P
3 ml
PP
S b T *




P W A d

2

30 B
v 40 4 v
2 0 3 m




(b) (4), 3-mL ChloroPrep With Tint (Teal) Applicator Lidding




(b) (4)


Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.






Lot/Exp Placeholder - for reference only:
 Lot # and Exp. Date are embossed on package.



(b) (4)

10.5-ml ChloroPrep One-Step
In-Line Mold Labels




(b) (4)

(b) (4)

(b) (4) 10.5-mL ChloraPrep Applicator Tyvek lidding for K-8

(b) (4)



Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.

<div style="background-color: #f08080; text-align: center; padding: 5px; font-weight: bold; color: white;"> ⚠ WARNING FLAMMABLE </div> <div style="padding: 5px;"> <p> Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet material from prep area </div>	<div style="text-align: center;"> <h2 style="margin: 0;">ChloraPrep® One-Step</h2> <h2 style="margin: 0;">ChloraPrep® With Tint</h2> <p style="margin: 0;">2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA) Patient Preoperative Skin Preparation</p> <h3 style="margin: 0;">10.5ml APPLICATOR</h3> </div> <div style="display: flex; justify-content: space-around; margin-top: 10px;"> <div style="text-align: center;"> Do Not Reuse </div> <div style="text-align: center;"> Latex Free </div> </div> <p style="text-align: center; margin-top: 5px;">External Use Only Professional Use Only</p> <p style="font-size: 0.8em; margin-top: 5px;">Cat. No. 260700 / 260700NS / 260700NSB NDC 054365-400-04 Cat. No. 260715 / 260715NS / 260715NSB NDC 054365-400-12 Cat. No. 260725 / 260725NS / 260725NSB NDC 054365-400-09</p>						
<div style="border: 1px solid black; padding: 5px;"> <h3 style="margin: 0;">Drug Facts</h3> <table style="width: 100%; border-collapse: collapse; font-size: 0.9em;"> <tr> <td style="width: 70%;">Active ingredients</td> <td style="width: 30%; text-align: right;">Purposes</td> </tr> <tr> <td>Chlorhexidine gluconate 2% w/v.....</td> <td style="text-align: right;">.....Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v.....</td> <td style="text-align: right;">.....Antiseptic</td> </tr> </table> <p style="font-size: 0.8em; margin-top: 5px;">Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. (11)</p> <p>Warnings</p> <p>For external use only. Flammable, keep away from fire or flame. To reduce risk of fire: PREP CAREFULLY</p> <ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area <p>Do not use</p> <ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser </div>		Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....Antiseptic	Isopropyl alcohol 70% v/v.....Antiseptic
Active ingredients	Purposes						
Chlorhexidine gluconate 2% w/v.....Antiseptic						
Isopropyl alcohol 70% v/v.....Antiseptic						

ChloraPrep® One-Step / ChloraPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm²)
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)
- The tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired. (9)

Inactive ingredients ■ FD&C green #3 dye (Scrub-Teal applicators only) ■ FD&C yellow #6 dye (High-Life Orange applicators only) ■ USP purified water

Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-640248 • I0411 • Made in the USA of US and imported parts

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.


Drug Facts	
Active ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Purposes Antiseptic Antiseptic
Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection. (11)	
Warnings For external use only. Flammable, keep away from fire or flame. To reduce the risk of fire, PREP CAREFULLY:	
<ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area 	
Do not use	
<ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser 	
When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.	
Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ use in a well ventilated area ■ maximal treatment area for one applicator is approximately 8.4 in. x 8.4 in. (457 cm²) ■ remove applicator from package; do not touch sponge ■ hold the applicator with the sponge down. Pinch wings <u>only once</u> to activate the ampule and release the antiseptic. ■ wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes ■ do not allow solution to pool; tuck prep towels to absorb solution, and then remove ■ allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available. 	
Other information	
<ul style="list-style-type: none"> ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) 	
Inactive ingredients	
<ul style="list-style-type: none"> ■ FD&C green #3 dye ■ USP purified water 	
Questions?	
<ul style="list-style-type: none"> ■ www.chloraprep.com ■ call 1-800-525-0502 (M-F 8 a.m.–5 p.m. CST) 	

Chloraprep
 W
 2% w/v chlorhexidine
 and 70% v/v isopropyl
 Patient Preparation

10
 APPLICATORS

Scrub Teal


25 applicators



(b) (4)

10.5-ml ChloroPrep With Tint
In-Line Mold Labels



(b) (4)



(b) (4)


(b) (4)

(b) (4) 10.5-mL ChloroPrep With Tint Applicator Tyvek lidding



Lot/Exp Placeholder - for reference only:
 Lot # and Exp. Date are embossed on package.

Lot # and Exp. Date are embossed on the blister film at this end of the package.



1 15/32

3 1/4

17 1/4

6 1/2

3 1/16

2 51/64

11

5 5/8

12 27/32


5 19/32

37 19/32


P g A d

(b) (4)


10.5-mL ChloraPrep With Tint In-Line Mold Labels



(b) (4)




(b) (4) 10.5-mL ChloroPrep With Tint (Orange) Applicator Tyvek
 lidding



ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
26ml APPLICATOR *Clear*

**⚠ WARNING
FLAMMABLE**

 **Keep away
from fire
or flame.**

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773800-L - F0811

ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
26ml APPLICATOR *Hi-Lite Orange®*

**⚠ WARNING
FLAMMABLE**

 **Keep away
from fire
or flame.**

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773815-L - F0811

ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
26ml APPLICATOR *Scrub Teal®*

**⚠ WARNING
FLAMMABLE**

 **Keep away
from fire
or flame.**

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Made in the USA of US and imported parts
CareFusion, Leawood, KS 66211
Artwork No. 6-773825-L - F0811


Made in the USA of US and imported parts
CareFusion, Leewood, KS 66211
Artwork No. 6-773800-R - F0611

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

26 ml APPLICATOR **Clear**

**⚠ WARNING
FLAMMABLE**

 Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

①


Made in the USA of US and imported parts
CareFusion, Leewood, KS 66211
Artwork No. 6-773815-R - F0611

ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

26 ml APPLICATOR **Hi-Life Orange®**

**⚠ WARNING
FLAMMABLE**

 Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

①


Made in the USA of US and imported parts
CareFusion, Leewood, KS 66211
Artwork No. 6-773825-R - F0811

ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

26 ml APPLICATOR **Scrub Teal®**

**⚠ WARNING
FLAMMABLE**

 Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in x 8.4 in.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

①

Cat. No. 260800 NDC 054365-400-14
Single Use • Applicator is STERILE if package is intact

ChloroPrep® One-Step

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
26ml APPLICATOR

Clear

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire; PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hr in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloroprep.com
■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-787800 • H0611
Made in the USA of US and imported parts

Cat. No. 260825 NDC 054365-400-05

Single Use • Applicator is STERILE if package is intact

ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

26ml APPLICATOR

Scrub Teal®

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire; PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- FD&C green #3 dye
- USP purified water

Questions?

- www.chloroprep.com
- call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 • Artwork No. 6-786825 - K0411
Made in the USA of US and imported parts

Cat. No. 260815 NDC 054365-400-13
Single Use • Applicator is STERILE if package is intact
ChloroPrep® With Tint
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
26ml APPLICATOR

Hi-Lite Orange®

WARNING, FLAMMABLE, KEEP AWAY FROM FIRE OR FLAME

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire; PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or Isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²). Do not use 26-ml applicator for area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead.
- do not use 26-ml applicator for head and neck surgery
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wing **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- FD&C yellow #6 dye
- USP purified water

Questions? ■ www.chloroprep.com
■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

CareFusion, Leawood, KS 66211 Artwork No. 6-786815 • J0411
Made in the USA of US and imported parts

⚠ WARNING
FLAMMABLE

Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery or on an area smaller than 8.4 in. x 8.4 in. Use a small applicator instead.
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

ChloroPrep® One-Step ChloroPrep® With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation

26ml APPLICATOR

L Latex Free D Do Not Reuse

External Use Only Professional Use Only

Cat. No. 260800 / 260800NS / 260800NSB NDC 054365-400-14
 Cat. No. 260815 / 260815NS / 260815NSB NDC 054365-400-13
 Cat. No. 260825 / 260825NS / 260825NSB NDC 054365-400-05

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- do not use 26-ml applicator for head and neck surgery
- do not use on an area smaller than 8.4 in. x 8.4 in. Use a smaller applicator instead
- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

ChloroPrep® One-Step/ChloroPrep With Tint

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

Drug Facts (continued)

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 13.2 in. x 13.2 in. (1126 cm²).
- remove applicator from package; do not touch sponge
- hold the applicator with the sponge down. Pinch wings only once to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- do not allow solution to pool; tuck prep towels to absorb solution, and then remove
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.


Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F) ■ The tint will slowly fade from the skin. Soap and water, or alcohol may be used to remove the tint if desired. ?

Inactive ingredients ■ FD&C green #3 dye (Scrub-Teal applicators only) ■ FD&C yellow #6 dye (Hi-Lite Orange applicators only) ■ USP purified water

Questions? ■ www.chloroprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)


CareFusion, Leawood, KS 66211 • Artwork No. 6-775815 • H0411 • Made in the USA of US and imported parts

(b) (4) - Intermediate Carton Dispenser - ChloroPrep Sepp Box artwork



(b) (4)

(b) (4)



(b) (4)

6-775449F ChloroPrep 2% Sepp Applicator, Non-Sterile - Folded Drug Facts Insert

2.5"

7.5"

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
0.67ml SEPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients	Purposes
Chlorhexidine gluconate 2% w/v.....	Antiseptic
Isopropyl alcohol 70% v/v.....	Antiseptic

Use for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.

Warnings
For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- remove applicator from package; do not touch applicator tip
- hold the sponge tip down and pinch the barrel of the applicator **only once** to activate the ampule and release the antiseptic
- wet the sponge tip by pressing and releasing the sponge against the treatment area until liquid is visible on the skin
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15–30 °C (59–86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloroprep.com
■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)

See other side for additional information

Front

2.5"

7.5"

ChloroPrep® One-Step
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
0.67ml SEPP® APPLICATOR

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.



⚠ WARNING
FLAMMABLE

Keep away from fire or flame.

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

See Drug Facts for Full Labeling Information

 Do Not Reuse
  Latex Free

External Use Only Professional Use Only

Single Use

CareFusion Leawood, KS 66211
Artwork 6-775449F • L0411
Made in the USA of US and imported parts

Back

(b) (4)

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v Antiseptic
Isopropyl alcohol 70% v/v Antiseptic

Purposes

Use for the preparation of the patient's skin prior to surgery or injection.
Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove all wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or Isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- tear pouch at side notch to reveal applicator handle. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information ■ store between 15-30 °C (59-86 °F)
■ avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients ■ USP purified water

Questions? ■ www.chloraprep.com
■ call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)


SING
SWABST
APPLICATOR

Clear

LOT XX





Cat. No. 260100
©2009 CoreFusion, All rights reserved.



(b) (4) Single Swabstick Foil Pouch

(b) (4)



.95" From left edge of live area to point of arrow for tear perforation

Lot/Exp Placeholder - for reference only:

Lot # and Exp. Date are embossed on package.

<div style="background-color: #f08080; text-align: center; padding: 2px;">⚠ WARNING</div> <div style="background-color: #f08080; text-align: center; padding: 2px;">FLAMMABLE</div> <p>Keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> • solution contains alcohol and gives off flammable vapors • avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. • do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) • do not allow solution to pool • remove wet material from prep area 	<h2 style="margin: 0;">ChloraPrep® Single Swabstick ChloraPrep® Triple Swabstick</h2> <p style="margin: 0;">2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)</p> <p style="margin: 0;">Patient Preoperative Skin Preparation</p> <p style="margin: 0;">1.75ml APPLICATOR 5.25ml APPLICATOR</p> <p style="margin: 0;"> L Latex Free X Do Not Reuse </p> <p style="margin: 0;">Single Use External Use Only Professional Use Only</p> <p style="margin: 0;">Cat. No. 260100 / 260100B NDC 054365-400-07 Cat. No. 260103 / 260103B NDC 054365-400-08</p>						
<h3 style="margin: 0;">Drug Facts</h3> <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 70%;">Active ingredients</td> <td style="width: 30%;">Purposes</td> </tr> <tr> <td>Chlorhexidine gluconate 2% w/v</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v</td> <td>Antiseptic</td> </tr> </table> <p>Use for the preparation of the patient's skin prior to surgery; Helps to reduce bacteria that potentially can cause skin infection.</p> <p>Warnings</p> <p>For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:</p> <ul style="list-style-type: none"> ■ solution contains alcohol and gives off flammable vapors ■ avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable. ■ do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair) ■ do not allow solution to pool ■ remove wet materials from prep area <p>Do not use</p> <ul style="list-style-type: none"> ■ on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol ■ for lumbar puncture or in contact with the meninges ■ on open skin wounds or as a general skin cleanser 		Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v	Antiseptic	Isopropyl alcohol 70% v/v	Antiseptic
Active ingredients	Purposes						
Chlorhexidine gluconate 2% w/v	Antiseptic						
Isopropyl alcohol 70% v/v	Antiseptic						

14

ChloraPrep® Single Swabstick/ChloraPrep® Triple Swabstick

2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

<p>Drug Facts (continued)</p> <p>When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p> <p>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p>Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p>
<p>Directions</p> <ul style="list-style-type: none"> ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ■ use in a well ventilated area ■ maximal treatment area using <ul style="list-style-type: none"> ■ one single swabstick applicator is approximately 2.5 in. x 2.5 in. (42 cm²) ■ three swabsticks sequentially is approximately 5 in. x 5 in. (160 cm²) ■ tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area. ■ completely wet the treatment area with antiseptic ■ dry surgical sites (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds ■ moist surgical sites (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes ■ allow the solution to completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away. ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.
<p>Other information ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)</p>
<p>Inactive ingredients ■ USP purified water</p>
<p>Questions? ■ www.chloraprep.com ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)</p>

CareFusion, Leawood, KS 66211 • Artwork No. 6-656900 • L0411 • Made in the USA of US and imported parts

WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Drug Facts

Active ingredients

Chlorhexidine gluconate 2% w/v Antiseptic
 Isopropyl alcohol 70% v/v Antiseptic

Purposes

Use for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove all wet materials from prep area

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- use in a well ventilated area
- maximal treatment area using three swabsticks sequentially is approximately 5 in x 5 in (160 cm²)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes
- **allow the solution to completely dry** (minimum of 3 minutes on hairless skin; up to 1 hour in hair). Do not blot or wipe away.
- discard the applicators after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredients

- USP purified water

Questions?

- www.chloraprep.com
- call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)

TRIP
SWABST

Clear

LOT



14

7

(b) (4) Triple Swabstick Foil Pouch

(b) (4)

5.5"

5"

2"

2.375"

ChloroPrep® Triple Swabstick
2% w/v chlorhexidine gluconate (CHG)
and 70% v/v isopropyl alcohol (IPA)
Patient Preoperative Skin Preparation
5.25ml APPLICATOR

See Drug Facts for full labeling information
Made in the USA of US and imported parts

Single Use
Latex Free
Cat. No. 260103 NDC 054365-400-08

Single use only
Manufactured by:
Carefusion
Leawood, KS 66211
1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)
www.chloroprep.com
Artwork No. 6-800104 60611

Tear Here

**WARNING
FLAMMABLE**

Keep away from fire or flame.
To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry.
Wet hair is flammable.
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair)
- do not allow solution to pool
- remove wet material from prep area

Directions:

- Tear pouch at notch.
- Pull across package to expose the ends of the swabsticks.
- Do not touch foam when removing swabsticks from pouch.
- Place foam flat side down on treatment area.

Lot # and Exp. Date are embossed on package film

.95" From left edge of live area to point of arrow for tear perforation

Lot/Exp Date placeholder - for reference only:

Lot # and Exp. Date are embossed on package film

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
08/16/2012