

Artwork No. 6-775449F, ChloroPrep 2% Sepp Applicator, Non-Sterile - Folded Drug Facts Insert

(b) (4)

(b) (4)

Dimensions: 7.5" x 2.5"

2.5"

**ChloroPrep® One-Step**  
 2% w/v chlorhexidine gluconate (CHG)  
 and 70% v/v isopropyl alcohol (IPA)  
**Patient Preoperative Skin Preparation**  
 Non-sterile Solution  
 Applicator is sterile if package is intact  
 0.67 ml SEPP® APPLICATOR

Drug Facts						
<table style="width: 100%; border-collapse: collapse;"> <tr> <th style="text-align: left;">Active ingredients</th> <th style="text-align: left;">Purposes</th> </tr> <tr> <td>Chlorhexidine gluconate 2% w/v.....</td> <td>Antiseptic</td> </tr> <tr> <td>Isopropyl alcohol 70% v/v.....</td> <td>Antiseptic</td> </tr> </table>	Active ingredients	Purposes	Chlorhexidine gluconate 2% w/v.....	Antiseptic	Isopropyl alcohol 70% v/v.....	Antiseptic
Active ingredients	Purposes					
Chlorhexidine gluconate 2% w/v.....	Antiseptic					
Isopropyl alcohol 70% v/v.....	Antiseptic					
<p><b>Use</b> for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.</p>						
<p><b>Warnings</b>                      For external use only  <b>Flammable, keep away from fire or flame</b>                      ■ do not use with electrocautery procedures</p>						
<p><b>Allergy Alert</b>                      This product may cause a severe allergic reaction. Symptoms may include:                      ■ wheezing/difficulty breathing ■ shock                      ■ facial swelling ■ hives ■ rash                      If an allergic reaction occurs, stop use and seek medical help right away.</p>						
<p><b>Do not use</b>                      ■ on patients allergic to chlorhexidine gluconate or any other ingredient in this product                      ■ for lumbar puncture or in contact with the meninges                      ■ on open skin wounds or as a general skin cleanser</p>						
<p><b>When using this product</b> keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.</p>						
<p><b>Stop use and ask a doctor if</b> irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p>						
<p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p>						
<p><b>Directions</b>                      ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.                      ■ maximal treatment area for one applicator is approximately 2.5 in. x 2.5 in. (42 cm<sup>2</sup>)                      ■ remove applicator from package; do not touch applicator tip                      ■ hold the sponge tip down and pinch the barrel of the applicator <u>only once</u> to activate the ampule and release the antiseptic                      ■ wet the sponge tip by pressing and releasing the sponge against the treatment area until liquid is visible on the skin                      ■ completely wet the treatment area with antiseptic                      ■ <b>dry surgical sites</b> (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for approximately 30 seconds. Allow the area to air dry for approximately 30 seconds.                      ■ <b>moist surgical sites</b> (e.g., inguinal fold): use gentle repeated back-and-forth strokes for approximately 2 minutes. Allow the area to air dry for approximately 1 minute.                      ■ do not blot or wipe away                      ■ discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.</p>						
<p><b>Other information</b> ■ store between 15–30 °C (59–86 °F)                      ■ avoid freezing and excessive heat above 40 °C (104 °F)</p>						
<p><b>Inactive ingredient</b> USP purified water</p>						
<p><b>Questions?</b> ■ <a href="http://www.chloroprep.com">www.chloroprep.com</a>                      ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)</p>						


See other side for additional information


Front

NDC 054365-400-03  
 Cat. No. 260441NSB / 260449NSB

**ChloroPrep® One-Step**  
 2% w/v chlorhexidine gluconate (CHG)  
 and 70% v/v isopropyl alcohol (IPA)  
**Patient Preoperative Skin Preparation**  
 Non-sterile Solution  
 Applicator is sterile if package is intact  
 0.67 ml SEPP® APPLICATOR

**See Drug Facts for Full Labeling Information**

 Do Not Reuse

 Not made with natural rubber latex

External Use Only

Professional Use Only

CareFusion, El Paso, TX 79912

Artwork 6-775449F · PXXXXXX  
 Made in the USA of US and imported parts

Back


(b) (4)

Artwork No. 6-000449, ChloroPrep One-Step Sepp (b) (4)


(b) (4)

(b) (4)

(b) (4)



Artwork No. 6-824449, Chloraprep One-Step SEPP Intermediate Carton



Artwork No. 6-656900, Chloraprep One-Step Drug Facts Insert for Single & Triple Swabsticks

(b) (4)

(b) (4)

4.75" width



Front

(b) (4)

Cat. No. 260100 / 260100B NDC 054365-400-07  
Cat. No. 260103 / 260103B NDC 054365-400-08

**Chloraprep® Single Swabstick / Chloraprep® Triple Swabstick**  
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation

**Non-sterile Solution**  
**Applicator is non-sterile**  
1.75 ml APPLICATOR / 5.25 ml APPLICATOR

 Not made with natural rubber latex       Do Not Reuse      Professional Use Only  
External Use Only

Drug Facts	
<b>Active ingredients</b>	<b>Purposes</b>
Chlorhexidine gluconate 2% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

**Use** for the preparation of the patient's skin prior to surgery or injection. Helps to reduce bacteria that potentially can cause skin infection.

**Warnings**  
For external use only. Flammable, keep away from fire or flame.  
■ do not use with electrocautery procedures

**Allergy Alert**  
This product may cause a severe allergic reaction. Symptoms may include:  
■ wheezing/difficulty breathing ■ shock ■ facial swelling ■ hives ■ rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**  
■ on patients allergic to chlorhexidine gluconate or any other ingredient in this product  
■ for lumbar puncture or in contact with the meninges  
■ on open skin wounds or as a general skin cleanser

3.8" length

**Chloraprep® Single Swabstick / Chloraprep® Triple Swabstick**  
2% w/v chlorhexidine gluconate (CHG) and 70% v/v isopropyl alcohol (IPA)

**Drug Facts (continued)**

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

**Stop use and ask a doctor** if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.  
**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area using
  - one single swabstick applicator is approximately 2.5 in. x 2.5 in. (42 cm<sup>2</sup>)
  - three swabsticks sequentially is approximately 5 in. x 5 in. (160 cm<sup>2</sup>)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes for 30 seconds (when using the triple swabstick applicators, use each swabstick sequentially within the 30 seconds). Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth strokes for 2 minutes (when using the triple swabstick applicators, use each swabstick sequentially within the 2 minutes). Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

**Other information** ■ store between 15–30 °C (59–86 °F) ■ avoid freezing and excessive heat above 40 °C (104 °F)

**Inactive ingredient** USP purified water

**Questions?** ■ [www.chloraprep.com](http://www.chloraprep.com) ■ call 1-800-523-0502 (M-F 8 a.m.–5 p.m. CST)



CareFusion, El Paso, TX 79912 · Artwork No. 6-656900 · OXXXXXX · Made in the USA of US and imported parts

Back

Artwork No. 6-800101, Chloraprep One-Step Single Swabstick Foil Pouch

(b) (4)


(b) (4)



.95" From left edge of live area to point of arrow for tear perforation

Lot/Exp Placeholder - for reference only.

Artwork No. 6-824100 ChloroPrep Single Swabstick Intermediate Carton - Non-sterile



Artwork No. 6-800104, Chloraprep One-Step Triple Swabstick Foil Pouch

(b) (4)

(b) (4)

5.5"

5"

2"

2.375"

**Chloraprep® Triple Swabstick**  
2% w/v chlorhexidine gluconate (CHG)  
and 70% v/v isopropyl alcohol (IPA)  
Patient Preoperative Skin Preparation  
**Non-sterile Solution**  
**Applicator is non-sterile**  
5.25 ml APPLICATOR  
See Drug Facts for full labeling information  
Made in the USA of US and imported parts

Do Not Reuse

Cat. No. 260103 NDC 054365-400-08

Image of three swabsticks.

Manufactured by:  
CareFusion  
El Paso, TX 79912  
1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)  
[www.chloraprep.com](http://www.chloraprep.com)  
Artwork No. 6-800104 1090314

Tear Here

**WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**

**Directions:**

- Tear pouch at notch.
- Pull across package to expose the ends of the swabsticks.
- Do not touch foam when removing swabsticks from pouch.
- Place foam flat side down on treatment area.

Lot # and Exp. Date are embossed on package film

Lot/Exp Date placeholder - for reference only.

95" From left edge of live area to point of arrow for tear perforation



---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/  
-----

VALERIE S PRATT  
09/12/2017