

NDC 63824-057-82

**Mucinex<sup>D</sup>**  
600 mg guaifenesin and  
60 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND  
NASAL DECONGESTANT**

2 Bi-layer Tablets

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

Contains FD&C Yellow Dye #6

**PHYSICIAN SAMPLE-  
NOT TO BE SOLD**

**Adams<sup>®</sup>**  
LABORATORIES, INC.  
Fort Worth, Texas 76155  
US Pat. 6,372,252 B1  
3105702 102003

3-6382405782-8


4 1/4"

15/16"

NDC 63824-057-20

**Mucinex<sup>D</sup>**  
600 mg guaifenesin and  
60 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND  
NASAL DECONGESTANT**



20 Bi-layer Tablets

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

See carton for complete Drug Facts.

US Pat. 6,372,252 B1

102003  
3005720

Contains FD&C Yellow Dye #6

*Please visit our Web site:  
[www.mucinex.com](http://www.mucinex.com)*

**Adams<sup>®</sup>**  
LABORATORIES, INC.  
Fort Worth, Texas 76155  
© 2003 Adams Laboratories, Inc.


5 1/4"

2"

NDC 63824-057-40

**Mucinex<sup>D</sup>**  
600 mg guaifenesin and  
60 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND  
NASAL DECONGESTANT**



40 Bi-layer Tablets

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

See carton for complete Drug Facts.

US Pat. 6,372,252 B1

102003  
3005740

Contains FD&C Yellow Dye #6

*Please visit our Web site:  
[www.mucinex.com](http://www.mucinex.com)*

**Adams<sup>®</sup>**  
LABORATORIES, INC.  
Fort Worth, Texas 76155  
© 2003 Adams Laboratories, Inc.

5 1/4"

2"

NDC 63824-034-82

**Mucinex<sup>D</sup>**  
1200 mg guaifenesin and  
120 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND  
NASAL DECONGESTANT**

2 Bi-layer Tablets

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

PHYSICIAN SAMPLE-  
NOT TO BE SOLD

**Adams<sup>®</sup>**  
LABORATORIES, INC.  
Fort Worth, Texas 76155  
US Pat. 6,372,252 B1  
3103402 102003

3-63824-03482-9


4 1/4"

15/16"

NDC 63824-034-01

**Mucinex<sup>®</sup>D**  
1200 mg guaifenesin and  
120 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND  
NASAL DECONGESTANT**



10 Bi-layer Tablets

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

See carton for complete Drug Facts.

US Pat. 6,372,252 B1

102003  
3003401

*Please visit our Web site:  
[www.mucinex.com](http://www.mucinex.com)*

**Adams<sup>®</sup>**  
LABORATORIES, INC.  
Fort Worth, Texas 76155  
© 2003 Adams Laboratories, Inc.


5 1/4"

2"

NDC 63824-034-20

**Mucinex<sup>®</sup>D**  
1200 mg guaifenesin and  
120 mg pseudoephedrine HCl per tablet

**EXPECTORANT AND  
NASAL DECONGESTANT**



20 Bi-layer Tablets

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

See carton for complete Drug Facts.

US Pat. 6,372,252 B1

102003  
3003420


**Adams<sup>®</sup>**  
LABORATORIES, INC.  
Fort Worth, Texas 76155  
© 2003 Adams Laboratories, Inc.

*Please visit our Web site:  
[www.mucinex.com](http://www.mucinex.com)*

5 1/4"

2"

FOR LAYOUT PURPOSES ONLY



# MUCINEX D 600/60 mg SAMPLE TRAY

Title:  
 14 pt. Helvetica Bold  
 Italic, left justified

Headings:  
 8 pt. Helvetica Bold  
 Italic, left justified

Bullets:  
 5 pt. solid square  
 Zapf Dingbats

Subheadings:  
 6 pt. Helvetica Bold  
 left justified

Graphic leading  
 to next panel

<b>Drug Facts</b>		<b>Drug Facts (continued)</b>	
<b>Active ingredients (in each extended-release tablet)</b>		<b>When using this product</b>	
Guaifenesin 600 mg	Expectorant	do not use more than directed	
Pseudoephedrine HCl 60 mg	Nasal Decongestant	<b>Stop use and ask a doctor if</b>	
<b>Uses</b>		<ul style="list-style-type: none"> <li>■ you get nervous, dizzy, or sleepless</li> <li>■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.</li> </ul>	
<ul style="list-style-type: none"> <li>■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive</li> <li>■ temporarily relieves nasal congestion due to:                             <ul style="list-style-type: none"> <li>■ common cold</li> <li>■ hay fever</li> <li>■ upper respiratory allergies</li> </ul> </li> <li>■ temporarily restores freer breathing through the nose</li> <li>■ promotes nasal and/or sinus drainage</li> <li>■ temporarily relieves sinus congestion and pressure</li> </ul>		<b>If pregnant or breast-feeding</b> , ask a health professional before use.	
<b>Warnings</b>		<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
<p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>■ heart disease</li> <li>■ high blood pressure</li> <li>■ thyroid disease</li> <li>■ diabetes</li> <li>■ trouble urinating due to an enlarged prostate gland</li> <li>■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema</li> <li>■ cough accompanied by too much phlegm (mucus)</li> </ul>		<b>Directions</b>	
		<ul style="list-style-type: none"> <li>■ do not crush, chew, or break tablet</li> <li>■ take with a full glass of water</li> <li>■ this product can be administered without regard for timing of meals</li> <li>■ adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours</li> <li>■ children under 12 years of age: do not use</li> </ul>	
		<b>Other information</b>	
		<ul style="list-style-type: none"> <li>■ tamper evident: do not use if seal on bottle printed "SEALED FOR YOUR PROTECTION" is broken or missing</li> <li>■ store at 20-25°C (68-77°F)</li> <li>■ see bottom of bottle for lot code and expiration date</li> </ul>	
		<b>Inactive ingredients</b>	
		carbomer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF	

Right justified

Title for continued panel:  
 8 pt. Helvetica Bold Italic


8 pt. Helvetica Regular

0.5 pt. hairline

2.5 pt. barline

Body text:  
 6 pt. Helvetica Regular  
 with 6.5 pt. leading,  
 left justified

2.5 pt. box barline



20 BI-LAYER TABLETS

**Mucinex<sup>D</sup>**  
NEW  
600 mg guaifenesin and  
60 mg pseudoephedrine HCl per tablet  
EXPECTORANT AND  
NASAL DECONGESTANT

**Drug Facts (continued)**

**When using this product**  
 ■ do not use more than directed

**Stop use and ask a doctor if**  
 ■ you get nervous, dizzy, or sleepless  
 ■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not crush, chew, or break tablet  
 ■ take with a full glass of water  
 ■ this product can be administered without regard for timing of meals  
 ■ adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours  
 ■ children under 12 years of age: do not use

**Other information**  
 ■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing  
 ■ store at 20-25°C (68-77°F)  
 ■ see bottom of bottle for lot code and expiration date

**Drug Facts (continued)**

**Inactive ingredients** carbomer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF


Please visit our Web site:  
[www.mucinex.com](http://www.mucinex.com)

Contains FD&C Yellow Dye #6

**Adams<sup>®</sup>**  
LABORATORIES, INC.  
Fort Worth, Texas 76155  
© 2003 Adams Laboratories, Inc.  
US Pat. 6,372,252 B1

102003  
2005720

NEW NDC 63824-057-20  
**Mucinex<sup>D</sup>**  
600 mg guaifenesin and  
60 mg pseudoephedrine HCl per tablet  
EXPECTORANT AND  
NASAL DECONGESTANT



20 BI-LAYER TABLETS

3-63824-057-20-0

**Drug Facts**

**Active ingredients (in each extended-release tablet)**

Active ingredients (in each extended-release tablet)	Purpose
Guaifenesin 600 mg	Expectorant
Pseudoephedrine HCl 60 mg	Nasal Decongestant

**Uses**  
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive  
 ■ temporarily relieves nasal congestion due to:  
 ■ common cold ■ hay fever  
 ■ upper respiratory allergies  
 ■ temporarily restores freer breathing through the nose  
 ■ promotes nasal and/or sinus drainage  
 ■ temporarily relieves sinus congestion and pressure

**Warnings**  
**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**  
 ■ heart disease ■ high blood pressure  
 ■ thyroid disease ■ diabetes  
 ■ trouble urinating due to an enlarged prostate gland  
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
 ■ cough accompanied by too much phlegm (mucus)

# MUCINEX D 600/60 mg (20 count) CARTON

Title:  
14 pt. Helvetica Bold  
Italic, left justified

Headings:  
8 pt. Helvetica Bold  
Italic, left justified

Bullets:  
5 pt. solid square  
Zapf Dingbats

Subheadings:  
6 pt. Helvetica Bold  
left justified

Graphic leading  
to next panel

----- Right justified

<b>Drug Facts</b>	
<b>Active ingredients (in each extended-release tablet)</b>	<b>Purpose</b>
Guafenesin 600 mg	Expectorant
Pseudoephedrine HCl 60 mg	Nasal Decongestant
<b>Uses</b>	
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive	
■ temporarily relieves nasal congestion due to:	
■ common cold	■ hay fever
■ upper respiratory allergies	
■ temporarily restores freer breathing through the nose	
■ promotes nasal and/or sinus drainage	
■ temporarily relieves sinus congestion and pressure	
<b>Warnings</b>	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have	
■ heart disease	■ high blood pressure
■ thyroid disease	■ diabetes
■ trouble urinating due to an enlarged prostate gland	
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema	
■ cough accompanied by too much phlegm (mucus)	

**Drug Facts (continued)**

**When using this product**

- do not use more than directed

**Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

**Other information**

- tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store at 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

**Drug Facts (continued)**

**Inactive ingredients** carbomer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

Title for continued panel:  
8 pt. Helvetica Bold Italic


----- 8 pt. Helvetica Regular

----- 2.5 pt. barline

Body text:  
6 pt. Helvetica Regular  
with 6.5 pt. leading,  
left justified

----- 0.5 pt. hairline

----- 2.5 pt. box barline



40 BI-LAYER TABLETS

**Mucinex<sup>D</sup>**  
 600 mg guaifenesin and  
 60 mg pseudoephedrine HCl per tablet  
 EXPECTORANT AND  
 NASAL DECONGESTANT

NEW

**Drug Facts (continued)**

**When using this product**  
 ■ do not use more than directed

**Stop use and ask a doctor if**  
 ■ you get nervous, dizzy, or sleepless  
 ■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not crush, chew, or break tablet  
 ■ take with a full glass of water  
 ■ this product can be administered without regard for timing of meals  
 ■ adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours  
 ■ children under 12 years of age: do not use

**Other information**  
 ■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing  
 ■ store at 20-25°C (68-77°F)  
 ■ see bottom of bottle for lot code and expiration date

**Drug Facts (continued)**

**Inactive ingredients** carboxer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF


Please visit our Web site:  
[www.mucinex.com](http://www.mucinex.com)

Contains FD&C Yellow Dye #6

**Adams<sup>®</sup>**  
 LABORATORIES, INC.  
 Fort Worth, Texas 76155  
 © 2003 Adams Laboratories, Inc.  
 US Pat. 6,372,252 B1

102003  
 2005740

NEW NDC 63824-057-40  
**Mucinex<sup>D</sup>**  
 600 mg guaifenesin and  
 60 mg pseudoephedrine HCl per tablet  
 EXPECTORANT AND  
 NASAL DECONGESTANT



40 BI-LAYER TABLETS

3-63824-057-40-8

**Drug Facts**

Active ingredients (in each extended-release tablet)	Purpose
Guaifenesin 600 mg	Expectorant
Pseudoephedrine HCl 60 mg	Nasal Decongestant

**Uses**  
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive  
 ■ temporarily relieves nasal congestion due to:  
 ■ common cold ■ hay fever  
 ■ upper respiratory allergies  
 ■ temporarily restores freer breathing through the nose  
 ■ promotes nasal and/or sinus drainage  
 ■ temporarily relieves sinus congestion and pressure

**Warnings**  
**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**  
 ■ heart disease ■ high blood pressure  
 ■ thyroid disease ■ diabetes  
 ■ trouble urinating due to an enlarged prostate gland  
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
 ■ cough accompanied by too much phlegm (mucus)

# MUCINEX D 600/60 mg (40 count) CARTON

Title:  
14 pt. Helvetica Bold  
Italic, left justified

Headings:  
8 pt. Helvetica Bold  
Italic, left justified

Bullets:  
5 pt. solid square  
Zapf Dingbats

Subheadings:  
6 pt. Helvetica Bold  
left justified

Graphic leading  
to next panel

----- Right justified

<b>Drug Facts</b>	
<b>Active ingredients (in each extended-release tablet)</b>	<b>Purpose</b>
Guafenesin 600 mg	Expectorant
Pseudoephedrine HCl 60 mg	Nasal Decongestant
<b>Uses</b>	
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive	
■ temporarily relieves nasal congestion due to:	
■ common cold	■ hay fever
■ upper respiratory allergies	
■ temporarily restores freer breathing through the nose	
■ promotes nasal and/or sinus drainage	
■ temporarily relieves sinus congestion and pressure	
<b>Warnings</b>	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have	
■ heart disease	■ high blood pressure
■ thyroid disease	■ diabetes
■ trouble urinating due to an enlarged prostate gland	
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema	
■ cough accompanied by too much phlegm (mucus)	

<b>Drug Facts (continued)</b>
<b>When using this product</b>
■ do not use more than directed
<b>Stop use and ask a doctor if</b>
■ you get nervous, dizzy, or sleepless
■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.
<b>If pregnant or breast-feeding</b> , ask a health professional before use.
<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.
<b>Directions</b>
■ do not crush, chew, or break tablet
■ take with a full glass of water
■ this product can be administered without regard for timing of meals
■ adults and children 12 years and older: two tablets every 12 hours; not more than 4 tablets in 24 hours
■ children under 12 years of age: do not use
<b>Other information</b>
■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
■ store at 20-25°C (68-77°F)
■ see bottom of bottle for lot code and expiration date

<b>Drug Facts (continued)</b>
<b>Inactive ingredients</b> carbomer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

Title for continued panel:  
8 pt. Helvetica Bold Italic

----- 8 pt. Helvetica Regular


----- 2.5 pt. barline

Body text:  
6 pt. Helvetica Regular  
with 6.5 pt. leading,  
left justified

----- 0.5 pt. hairline

----- 2.5 pt. box barline

FOR LAYOUT PURPOSES ONLY



# MUCINEX D 1200/120 mg SAMPLE TRAY

Title:  
 14 pt. Helvetica Bold  
 Italic, left justified

Headings:  
 8 pt. Helvetica Bold  
 Italic, left justified

Bullets:  
 5 pt. solid square  
 Zapf Dingbats

Subheadings:  
 6 pt. Helvetica Bold  
 left justified

Graphic leading  
 to next panel

<b>Drug Facts</b>		<b>Drug Facts (continued)</b>	
<b>Active ingredients (in each extended-release tablet)</b>		<b>When using this product</b>	
Guafenesin 1200 mg	Expectorant	■ do not use more than directed	
Pseudoephedrine HCl 120 mg	Nasal Decongestant	<b>Stop use and ask a doctor if</b> ■ you get nervous, dizzy, or sleepless	
<b>Uses</b> ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive		■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.	
■ temporarily relieves nasal congestion due to:		<b>If pregnant or breast-feeding</b> , ask a health professional before use.	
■ common cold ■ hay fever ■ upper respiratory allergies		<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	
■ temporarily restores freer breathing through the nose		<b>Directions</b>	
■ promotes nasal and/or sinus drainage		■ do not crush, chew, or break tablet ■ take with a full glass of water	
■ temporarily relieves sinus congestion and pressure		■ this product can be administered without regard for timing of meals	
<b>Warnings</b>		■ adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.		■ children under 12 years of age: do not use	
<b>Ask a doctor before use if you have</b>		<b>Other information</b>	
■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes		■ tamper evident: do not use if seal on bottle printed "SEALED FOR YOUR PROTECTION" is broken or missing	
■ trouble urinating due to an enlarged prostate gland		■ store at 20-25°C (68-77°F)	
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema		■ see bottom of bottle for lot code and expiration date	
■ cough accompanied by too much phlegm (mucus)		<b>Inactive ingredients</b> carbomer 934P, NF; FD&C red #40 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF	

Right justified

Title for continued panel:  
 8 pt. Helvetica Bold Italic


8 pt. Helvetica Regular

0.5 pt. hairline

2.5 pt. barline

Body text:  
 6 pt. Helvetica Regular  
 with 6.5 pt. leading,  
 left justified

2.5 pt. box barline



10 BI-LAYER TABLETS

**Mucinex<sup>D</sup>**  
1200 mg guaifenesin and  
120 mg pseudoephedrine HCl per tablet  
EXPECTORANT AND  
NASAL DECONGESTANT

**NEW**

**Drug Facts (continued)**

**When using this product**  
 ■ do not use more than directed

**Stop use and ask a doctor if**  
 ■ you get nervous, dizzy, or sleepless  
 ■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not crush, chew, or break tablet  
 ■ take with a full glass of water  
 ■ this product can be administered without regard for timing of meals  
 ■ adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours  
 ■ children under 12 years of age: do not use

**Other information**  
 ■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing  
 ■ store at 20-25°C (68-77°F)  
 ■ see bottom of bottle for lot code and expiration date

**Drug Facts (continued)**


**Inactive ingredients** carbomer 934P, NF; FD&C red #40 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

Please visit our Web site:  
[www.mucinex.com](http://www.mucinex.com)

**Adams<sup>®</sup>**  
LABORATORIES, INC.  
Fort Worth, Texas 76155  
© 2003 Adams Laboratories, Inc.  
US Pat. 6,372,252 B1

102003  
2003401

**NEW** NDC 63824-034-01  
**Mucinex<sup>D</sup>**  
1200 mg guaifenesin and  
120 mg pseudoephedrine HCl per tablet  
EXPECTORANT AND  
NASAL DECONGESTANT



10 BI-LAYER TABLETS

3-63824-034-01-0

**Drug Facts**

Active ingredients (in each extended-release tablet)	Purpose
Guaifenesin 1200 mg	Expectorant
Pseudoephedrine HCl 120 mg	Nasal Decongestant

**Uses**  
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive  
 ■ temporarily relieves nasal congestion due to:  
 ■ common cold ■ hay fever  
 ■ upper respiratory allergies  
 ■ temporarily restores freer breathing through the nose  
 ■ promotes nasal and/or sinus drainage  
 ■ temporarily relieves sinus congestion and pressure

**Warnings**  
**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**  
 ■ heart disease ■ high blood pressure  
 ■ thyroid disease ■ diabetes  
 ■ trouble urinating due to an enlarged prostate gland  
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
 ■ cough accompanied by too much phlegm (mucus)

# MUCINEX D 1200/120 mg (10 count) CARTON

Title:  
14 pt. Helvetica Bold  
Italic, left justified

Headings:  
8 pt. Helvetica Bold  
Italic, left justified

Bullets:  
5 pt. solid square  
Zapf Dingbats

Subheadings:  
6 pt. Helvetica Bold  
left justified

Graphic leading  
to next panel

----- Right justified

<b>Drug Facts</b>	
<b>Active ingredients (in each extended-release tablet)</b>	<b>Purpose</b>
Guafenesin 1200 mg.....Expectorant Pseudoephedrine HCl 120 mg.....Nasal Decongestant	
<b>Uses</b>	
■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive	
■ temporarily relieves nasal congestion due to:	
■ common cold	■ hay fever
■ upper respiratory allergies	
■ temporarily restores freer breathing through the nose	
■ promotes nasal and/or sinus drainage	
■ temporarily relieves sinus congestion and pressure	
<b>Warnings</b>	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.	
<b>Ask a doctor before use if you have</b>	
■ heart disease	■ high blood pressure
■ thyroid disease	■ diabetes
■ trouble urinating due to an enlarged prostate gland	
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema	
■ cough accompanied by too much phlegm (mucus)	

**Drug Facts (continued)**

**When using this product**  
■ do not use more than directed

**Stop use and ask a doctor if**  
■ you get nervous, dizzy, or sleepless  
■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not crush, chew, or break tablet  
■ take with a full glass of water  
■ this product can be administered without regard for timing of meals  
■ adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours  
■ children under 12 years of age: do not use

**Other information**  
■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing  
■ store at 20-25°C (68-77°F)  
■ see bottom of bottle for lot code and expiration date

**Drug Facts (continued)**

**Inactive ingredients** carbomer 934P, NF; FD&C red #40 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

Title for continued panel:  
8 pt. Helvetica Bold Italic


----- 8 pt. Helvetica Regular

-----2.5 pt. barline

Body text:  
6 pt. Helvetica Regular  
with 6.5 pt. leading,  
left justified

-----0.5 pt. hairline

-----2.5 pt. box barline



20 BI-LAYER TABLETS

**Mucinex<sup>®</sup>D**  
 NEW  
 1200 mg guaifenesin and  
 120 mg pseudoephedrine HCl per tablet  
 EXPECTORANT AND  
 NASAL DECONGESTANT

**Drug Facts (continued)**

**When using this product**  
 ■ do not use more than directed

**Stop use and ask a doctor if**  
 ■ you get nervous, dizzy, or sleepless  
 ■ symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not crush, chew, or break tablet  
 ■ take with a full glass of water  
 ■ this product can be administered without regard for timing of meals  
 ■ adults and children 12 years and older: 1 tablet every 12 hours; not more than 2 tablets in 24 hours  
 ■ children under 12 years of age: do not use

**Other information**  
 ■ tamper evident: do not use if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing  
 ■ store at 20-25°C (68-77°F)  
 ■ see bottom of bottle for lot code and expiration date

**Drug Facts (continued)**


**Inactive ingredients** carbomer 934P, NF; FD&C red #40 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

Please visit our Web site:  
[www.mucinex.com](http://www.mucinex.com)

**Adams<sup>®</sup>**  
 LABORATORIES, INC.  
 Fort Worth, Texas 76155  
 © 2003 Adams Laboratories, Inc.  
 US Pat. 6,372,252 B1

102003  
 2003420

**NEW** NDC 63824-034-20  
**Mucinex<sup>®</sup>D**  
 1200 mg guaifenesin and  
 120 mg pseudoephedrine HCl per tablet  
 EXPECTORANT AND  
 NASAL DECONGESTANT



20 BI-LAYER TABLETS

**Drug Facts**

Active ingredients (in each extended-release tablet)	Purpose
Guaifenesin 1200 mg	Expectorant
Pseudoephedrine HCl 120 mg	Nasal Decongestant

**Uses**  
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive  
 ■ temporarily relieves nasal congestion due to:  
 ■ common cold ■ hay fever  
 ■ upper respiratory allergies  
 ■ temporarily restores freer breathing through the nose  
 ■ promotes nasal and/or sinus drainage  
 ■ temporarily relieves sinus congestion and pressure

**Warnings**  
**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains a MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**  
 ■ heart disease ■ high blood pressure  
 ■ thyroid disease ■ diabetes  
 ■ trouble urinating due to an enlarged prostate gland  
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
 ■ cough accompanied by too much phlegm (mucus)

3-63824-034-20-1