

Drug Facts (continued)

Or

- using sterile scissors, cut off end seal of package
- transfer contents onto prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination
- use first cloth to prepare the skin area indicated for a moist or dry site, making certain to keep the second cloth where it will not be contaminated. Use second cloth to prepare larger areas.
- dry surgical sites** (such as abdomen or arm): use one cloth to cleanse each 161 cm² area (approximately 5 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- moist surgical sites** (such as inguinal fold): use one cloth to cleanse each 65 cm² area (approximately 2 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- discard each cloth after a single use
- after package has been opened discard any unused cloths

Other information

- store product flat
- store between 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

aloe vera, dimethicone, fragrance, glucono-delta-lactone, glycerin, Igepal®, polysorbate 20, propylene glycol, USP purified water
cloth: polyester

Questions or comments?
call toll-free 800-323-2220
(Monday to Friday 8 AM- 5 PM CST)
www.sageproducts.com

Reorder #9705



SAGE PRODUCTS

SAGE PRODUCTS LLC
3909 Three Oaks Road
Cary, Illinois 60013
www.sageproducts.com
800-323-2220

Patents:
www.sageproducts.com/patents
R5201J

NDC 053462-705

SAGE PRODUCTS

2% CHLORHEXIDINE GLUCONATE* CLOTH

PATIENT PREOPERATIVE SKIN PREPARATION

NON-STERILE

*equivalent to 500mg chlorhexidine gluconate per cloth

Rinse-Free — Alcohol-Free
For External Use Only — Single Use

- Provides rapid bactericidal action against a broad spectrum of microorganisms
- Significantly reduces the number of microorganisms on intact skin
- Demonstrates continued antimicrobial activity for up to 6 hours after application

2 disposable cloths
7.5in x 7.5in (19.1cm x 19.1cm)
DO NOT MICROWAVE

Drug Facts

Active ingredient Purpose
Chlorhexidine gluconate Antiseptic
2% solution

Uses

- helps reduce bacteria that can potentially cause skin infection
- for preparation of skin prior to surgery

Warnings
For external use only

Allergy Alert
This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar punctures or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product

- keep out of eyes, ears and mouth. May cause serious or permanent injury if chlorhexidine is permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children
If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns
- do not microwave**
- product and packaging are not sterile. Follow your hospital policy for skin preparation with non-sterile products.

To open package

- holding top of package in one hand, lift flap on backside of package with other hand
- grasp flap at top and pull down to tear flap away and expose foam
- hold outside of package to present foam and cloths to prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination

DO NOT FLUSH CLOTHS

NOT MADE WITH NATURAL RUBBER LATEX

NOTE: ARTWORK RUN-OUT TO BE ON TRAILING EDGE

Avoid excessive heat above 40°C (104°F)
Store between 20-25°C (68-77°F)

Drug Facts

Active ingredient Purpose
Chlorhexidine gluconate Antiseptic
2% solution

Uses

- helps reduce bacteria that can potentially cause skin infection
- for preparation of skin prior to surgery

Warnings
For external use only

Allergy Alert
This product may cause a severe allergic reaction. Symptoms may include:

- difficulty breathing
- swelling of the face, lips, tongue, or throat
- hives
- itching
- skin rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- on facial punctures or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product

- keep out of eyes, ears, and mouth. May cause serious or permanent injury if chlorhexidine is permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children
If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns
- do not microwave
- product and packaging are not sterile. Follow your hospital policy for skin preparation with non-sterile products.

To open package
holding top of package in one hand, lift flap on backside of package with other hand

- grasp flap at top and pull down to tear flap away and expose foam
- hold outside of package to present foam and cloths to prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination

Drug Facts (continued)

Or

- using sterile scissors, cut off end seal of package
- transfer contents onto prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination
- use first cloth to prepare the skin area indicated for a moist or dry site, making certain to keep the second cloth where it will not be contaminated. Use second cloth to prepare larger areas.
- dry surgical sites (such as abdomen or arm): use one cloth to cleanse each 65 cm² area (approximately 2 x 9 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- moist surgical sites (such as inguinal fold): use one cloth to cleanse each 65 cm² area (approximately 2 x 9 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- discard each cloth after a single use
- after package has been opened discard any unused cloths.


Other information

- store product flat
- store between 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients
aloe vera, dimethicone, fragrance, gluconic delta-lactone, glycerin, isopropyl alcohol, polysorbates 20, propylene glycol, USP purified water, cloth: polyester

Questions or comments?
call toll-free 800-323-2220 (Monday to Friday 9 AM-5 PM CST)
www.sageproducts.com

#9705



3 06 18029 85002 8

EXPIRATION DATE & LOT
PRINTED ON ATTACHED
LABEL

↑ THIS
END
UP ↑

SAGE
PRODUCTS

**2% CHLORHEXIDINE
GLUCONATE* CLOTH**
PATIENT PREOPERATIVE
SKIN PREPARATION

NON-STERILE

*equivalent to 500mg chlorhexidine gluconate per cloth

Rinse-Free — Alcohol-Free
For External Use Only — Single Use

- Provides rapid bactericidal action against a broad spectrum of microorganisms
- Significantly reduces the number of microorganisms on intact skin
- Demonstrates continued antimicrobial activity for up to 6 hours after application

DO NOT MICROWAVE

SAGE
PRODUCTS

**2% CHLORHEXIDINE
GLUCONATE* CLOTH**
PATIENT PREOPERATIVE
SKIN PREPARATION

NON-STERILE

*equivalent to 500mg chlorhexidine gluconate per cloth

Rinse-Free — Alcohol-Free
For External Use Only — Single Use

OPEN HERE

SAGE
PRODUCTS

**2% CHLORHEXIDINE
GLUCONATE* CLOTH**
PATIENT PREOPERATIVE
SKIN PREPARATION

NON-STERILE

*equivalent to 500mg chlorhexidine gluconate per cloth

Rinse-Free — Alcohol-Free
For External Use Only — Single Use

- Provides rapid bactericidal action against a broad spectrum of microorganisms
- Significantly reduces the number of microorganisms on intact skin
- Demonstrates continued antimicrobial activity for up to 6 hours after application


DO NOT MICROWAVE

Reorder #9705

Sage Products LLC
3909 Three Oaks Road
Cary, IL 60013 U.S.A.
800-323-2220 • 815-455-4700
www.sageproducts.com

MADE IN U.S.A.
R1768G

NET CONTENTS: #9705 - 24 PACKAGES (2 CLOTHS PER PACKAGE/48 CLOTHS PER BOX)



Drug Facts (continued)

Or

- using sterile scissors, cut off end seal of package
- transfer contents onto prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination
- use first cloth to prepare the skin area indicated for a moist or dry site, making certain to keep the second cloth where it will not be contaminated. Use second cloth to prepare larger areas.
- dry surgical sites** (such as abdomen or arm): use one cloth to cleanse each 161 cm² area (approximately 5 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- moist surgical sites** (such as inguinal fold): use one cloth to cleanse each 65 cm² area (approximately 2 x 5 inches) of skin to be prepared. Vigorously scrub skin back and forth for 3 minutes, completely wetting treatment area, then discard. Allow area to air dry for one (1) minute. Do not rinse.
- discard each cloth after a single use
- after package has been opened discard any unused cloths

Other information

- store product flat
- store between 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients
aloe vera, dimethicone, fragrance, glucono-delta-lactone, glycerin, Igepal®, polysorbate 20, propylene glycol, USP purified water
cloth: polyester

Questions or comments?
call toll-free 800-323-2220 (Monday to Friday 8 AM- 5 PM CST)
www.sageproducts.com

Reorder #9707



SAGE PRODUCTS
SAGE PRODUCTS LLC
3909 Three Oaks Road
Cary, Illinois 60013
www.sageproducts.com
800-323-2220
Patents:
www.sageproducts.com/patents
R5335F

NDC 053462-705-23

SAGE PRODUCTS

2% CHLORHEXIDINE GLUCONATE* CLOTH

PATIENT PREOPERATIVE SKIN PREPARATION

NON-STERILE

*equivalent to 500mg chlorhexidine gluconate per cloth

Rinse-Free — Alcohol-Free
For External Use Only — Single Use

- Provides rapid bactericidal action against a broad spectrum of microorganisms
- Significantly reduces the number of microorganisms on intact skin
- Demonstrates continued antimicrobial activity for up to 6 hours after application

2 disposable cloths
7.5in x 7.5in (19.1cm x 19.1cm)
DO NOT MICROWAVE

EXPIRATION DATE & LOT PLACEMENT HOLDER

Drug Facts

Active ingredient Purpose
Chlorhexidine gluconate Antiseptic
2% solution.....

Uses

- helps reduce bacteria that can potentially cause skin infection
- for preparation of skin prior to surgery

Warnings
For external use only

Allergy Alert
This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar punctures or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product

- keep out of eyes, ears and mouth. May cause serious or permanent injury if chlorhexidine is permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children
If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns
- do not microwave**
- product and packaging are not sterile. Follow your hospital policy for skin preparation with non-sterile products.

To open package

- holding top of package in one hand, lift flap on backside of package with other hand
- grasp flap at top and pull down to tear flap away and expose foam
- hold outside of package to present foam and cloths to prep table, avoiding contact between cloths and outside of package to reduce risk of cloth contamination

DO NOT FLUSH CLOTHS

NOT MADE WITH NATURAL RUBBER LATEX

2 X 0.37 EYESPOT

NOTE: ARTWORK RUN-OUT TO BE ON TRAILING EDGE

14 pt Helvetica Italic Bold

6/6.5 pt Helvetica Regular

2.5 pt Barline

2.5 pt Box Barline

6 pt Helvetica Bold Subheading

Added comma

.5 pt Hairline

8 pt Helvetica Italic Bold Heading

5 pt solid square

Added placeholder for Expiration date an

Avoid excessive heat above 40°C (104°F)
Store between 20-25°C (68-77°F)

Drug Facts

Active ingredient Purpose
Chlorhexidine gluconate Antiseptic
2% solution

Uses

- Helps reduce bacteria that can potentially cause skin infection
- For preparation of skin prior to surgery

Warnings

For external use only

Warnings

Do not use if you are allergic to chlorhexidine gluconate or any other ingredient in this product.

Directions

- Use with care to prevent irritation or chemical burns.
- Product and packaging are not sterile. Follow your hospital policy for use.
- Preparation and use instructions are on the inside of the package.
- Do not use if the top seal is broken or the top seal is damaged.
- Do not use if the top seal is broken or the top seal is damaged.
- Do not use if the top seal is broken or the top seal is damaged.

Drug Facts (continued)

Other information

- Keep out of reach of children.
- Use with care to prevent irritation or chemical burns.
- Product and packaging are not sterile. Follow your hospital policy for use.
- Preparation and use instructions are on the inside of the package.
- Do not use if the top seal is broken or the top seal is damaged.
- Do not use if the top seal is broken or the top seal is damaged.
- Do not use if the top seal is broken or the top seal is damaged.

#9707



3 08 16229 81055 9

EXPIRATION & LOT
PLACEMENT HOLDER

↑ THIS
END
UP ↑

Reorder #9707

Sage Products LLC
3909 Three Oaks Road
Cary, IL 60013 U.S.A.
800-323-2220 • 815-455-4700
www.sageproducts.com

MADE IN U.S.A.
R18530



OPEN HERE

SAGE PRODUCTS

2% CHLORHEXIDINE GLUCONATE* CLOTH

PATIENT PREOPERATIVE SKIN PREPARATION

NON-STERILE

*equivalent to 500mg chlorhexidine gluconate per cloth

Rinse-Free — Alcohol-Free
For External Use Only — Single Use

- Provides rapid bactericidal action against a broad spectrum of microorganisms
- Significantly reduces the number of microorganisms on intact skin
- Demonstrates continued antimicrobial activity for up to 6 hours after application

DO NOT MICROWAVE

2% CHLORHEXIDINE GLUCONATE* CLOTH

PATIENT PREOPERATIVE SKIN PREPARATION

NON-STERILE

*equivalent to 500mg chlorhexidine gluconate per cloth

Rinse-Free — Alcohol-Free
For External Use Only — Single Use

SAGE PRODUCTS

2% CHLORHEXIDINE GLUCONATE* CLOTH

PATIENT PREOPERATIVE SKIN PREPARATION

NON-STERILE

*equivalent to 500mg chlorhexidine gluconate per cloth

Rinse-Free — Alcohol-Free
For External Use Only — Single Use

- Provides rapid bactericidal action against a broad spectrum of microorganisms
- Significantly reduces the number of microorganisms on intact skin
- Demonstrates continued antimicrobial activity for up to 6 hours after application

DO NOT MICROWAVE

NET CONTENTS: #9707 - 16 PACKAGES (6 CLOTHS PER PACKAGE/96 CLOTHS PER BOX)