

Zantac 150 3ct Carton

06.13.19

ZAN_150_3ct_CTN_0140048-01_Mex_2019_FDA_DSN3



Sanofi Regulatory Specifications		Verified Date: 06/13/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9.0 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9.0 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%–102%
Leading (Minimum space in body of Drug Facts):	5.8 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.070 in
PDP dimensions (in millimeters)	H 60mm W 97mm	
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		V 8.81 pt. H 0.00 pt
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		V 61 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity to Logo/Largest Copy (pt.)		V 15% H 0%

Zantac 150 3ct Carton

06.13.19

ZAN_150_3ct_CTN_0140047-01_Spain_2019_FDA_DSN3

Tips for managing heartburn

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

MAXIMUM STRENGTH

Zantac

Ranitidine Tablets 150 mg/Acid Reducer **150**



PREVENTS & RELIEVES HEARTBURN associated with acid indigestion and sour stomach

3 TABLETS (3 DOSES)

Drug Facts (continued)

Inactive ingredients: hypromellose, magnesium stearate, microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

Questions? call 1-800-633-1610 or visit www.zantac.com

Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

LOT

EXP

Drug Facts

Active ingredient (in each tablet)
Ranitidine 150 mg (as ranitidine hydrochloride 168 mg)

Purpose
Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents or relieves heartburn, acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water
 - to prevent symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- do not use if individual blister unit is open or torn
- avoid excessive heat or humidity
- this product is sodium and sugar free

Distributed by: Otsuka, Inc., a Sanofi Company
P.O. Box 2219
Chattanooga, TN 37409-0219
© 2019 Origin Spain
0140047-01

6 81421 03101 1

Sanofi Regulatory Specifications		Verified Date: 06/13/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9.0 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9.0 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	5.8 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.070 in
PDP dimensions (in millimeters)	H 60mm W 97mm	
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")	V 8.81 pt. H 0.00 pt	
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")	V 61 pt. H 0.00 pt.	
Ratio of Statement of Identity to Logo/Largest Copy on PDP	V 15% H 0%	
Statement of Identity to Logo/Largest Copy (pt.)	V 15% H 0%	

Zantac 150 8ct Carton
03.28.19

ZAN_150_8ct_CTN_0140046-01_Mex_2019_FDA_DSN2



Sanofi Regulatory Specifications		Verified Date: 03/28/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 15% H 0%

Zantac 150 8ct Carton
03.28.19

ZAN_150_8ct_CTN_0140045-01_Spain_2019_FDA_DSN3



Sanofi Regulatory Specifications		Verified Date: 03/28/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 15% H 0%

Zantac 150 24ct Carton
03.18.19

ZAN_150_24ct_CTN_0140044-01_Mex_2019_FDA_DSN2



Sanofi Regulatory Specifications		Verified Date: 03/18/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	5.9 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 15% H 0%

Zantac 150 24ct Carton
03.14.19

ZAN_150_24ct_CTN_0140043-01_Spain_2019_FDA_DSN



Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	5.9 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 15% H 0%

Zantac 150 32ct Carton
03.14.19

ZAN_150_32ct_Bonus_CTN_0140074-01_Mex_2019_FDA_DSN



Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	5.9 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 15% H 0%

Zantac 150 32ct Carton
03.06.19

ZAN_150_32ct_Bonus_CTN_0140077-01_Spain_2019_FDA_DSN2



Sanofi Regulatory Specifications		Verified Date: 03/06/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	5.9 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 15% H 0%

Zantac 150 40ct Carton
03.14.19

ZAN_150_40ct_CTN_0140071-01_Mex_2019_FDA_DSN

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.) (Largest Logo Copy (pt.))		V 15% H 0%

Zantac 150 40ct Carton

03.14.19

ZAN_150_40ct_CTN_0140070-01_Spain_2019_FDA_DSN

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.) (Largest Logo Copy (pt.))		V 15% H 0%

Zantac 150 50ct Carton

03.14.19

ZAN_150_50ct_CTN_0140041-01_Spain_2019_FDA_DSN

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.) (Largest Logo Copy (pt.))		V 15% H 0%

Zantac 150 65ct Carton

02.28.19

ZAN_150_65ct_CTN_0140039-01_Mex_2019_FDA_DSN

MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

PREVENTS & RELIEVES
HEARTBURN associated with
acid indigestion and sour stomach

65 TABLETS
(65 DOSES)

Drug Facts (continued)

Active ingredient (in each tablet) Purpose
Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water
• to prevent symptoms, swallow 1 tablet with a glass of water
• 30 to 60 minutes before eating food or drinking beverages that cause heartburn
• can be used up to twice daily (do not take more than 2 tablets in 24 hours)
• children under 12 years: ask a doctor

Other information
• do not use if printed foil under bottle cap is open or torn
• store at 20°-25°C (68°-77°F)
• avoid excessive heat or humidity
• this product is sodium and sugar free

Inactive ingredients hypromellose, synthetic red iron oxide, titanium dioxide, microcrystalline cellulose, hypromellose, magnesium stearate, triacetin

Questions? call 1-800-833-1610 or visit www.zantac.com

Tips for managing heartburn
• Do not lie flat or bend over soon after eating
• Do not eat late at night, or just before bedtime
• Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

Stop use and ask a doctor if
• your heartburn continues or worsens
• you need to take this product for more than 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Ask a doctor before use if you have
• had heartburn over 3 months. This may be a sign of a more serious condition.
• heartburn with **lightheadedness, sweating or dizziness**
• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• frequent **chest pain**
• frequent wheezing, particularly with heartburn
• unexplained weight loss • nausea or vomiting
• stomach pain • kidney disease

Do not use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• with other acid reducers

Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Uses
• relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Tips for managing heartburn

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

0140039-01

©2019 Origin Mexico
Chattanooga, TN 37409-0219
Distributed by: Chatham, Inc., a Sanofi Company

SANOFI

Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

MOORE TIPS ON BOTTOM PANEL ▶

6 81421 03105 9

MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

65 TABLETS
(65 DOSES)

LOT

EXP

ZANTAC does not supply same brands

Sanofi Regulatory Specifications		Verified Date: 02/28/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.) (Largest Logo Copy (pt.))		V 15% H 0%

Zantac 150 65ct Carton

03.14.19

ZAN_150_65ct_CTN_0140040-01_Spain_2019_FDA_DSN

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.) (Largest Logo Copy (pt.))		V 15% H 0%

Zantac 150 78ct Carton

02.28.19

ZAN_150_78ct_Bonus_CTN_0140036-01_Mex_2019_FDA_DSN

MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

BONUS! 13 FREE TABLETS
BONUS! 13 FREE TABLETS

MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

PREVENTS & RELIEVES HEARTBURN associated with acid indigestion and sour stomach

78 ~~65~~ TABLETS (78 DOSES)

Drug Facts (continued)

Active ingredient (in each tablet) Purpose
Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer

Uses
• Relieves heartburn associated with acid indigestion and sour stomach
• Prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
Do not use
• If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• With other acid reducers

Ask a doctor before use if you have
• Had heartburn over 3 months. This may be a sign of a more serious condition.
• Heartburn with **lightheadedness, sweating or dizziness**, chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• Frequent **chest pain**
• Frequent wheezing, particularly with heartburn
• Unexplained weight loss • nausea or vomiting
• Stomach pain • kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
• Your heartburn continues or worsens
• You need to take this product for more than 14 days

Drug Facts (continued)

Other information
• Do not use if printed foil under bottle cap is open or torn
• Store at 20°-25°C (68°-77°F)
• Avoid excessive heat or humidity
• This product is sodium and sugar free

Inactive ingredients
microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

Directions
• Adults and children 12 years and over: swallow 1 tablet with a glass of water to relieve symptoms.
• To prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
• Can be used up to twice daily (do not take more than 2 tablets in 24 hours)
• Children under 12 years: ask a doctor

Tips for managing heartburn
• Do not eat late at night, or just before bedtime
• Do not lie flat or bend over soon after eating
• Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

Questions? call 1-800-833-1610 or visit www.zantac.com

Sanofi

Distributed by: Chatham, Inc., a Sanofi Company
P.O. Box 2219
Chattanooga, TN 37409-0219
©2019 Origin Mexico
0140036-01

Tips for managing heartburn

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

Drug Facts (continued)

Directions
• Adults and children 12 years and over: swallow 1 tablet with a glass of water to relieve symptoms.
• To prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
• Can be used up to twice daily (do not take more than 2 tablets in 24 hours)
• Children under 12 years: ask a doctor

Tips for managing heartburn
• Do not lie flat or bend over soon after eating
• Do not eat late at night, or just before bedtime
• Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

Questions? call 1-800-833-1610 or visit www.zantac.com

Drug Facts (continued)

Active ingredient (in each tablet) Purpose
Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer

Uses
• Relieves heartburn associated with acid indigestion and sour stomach
• Prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
Do not use
• If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• With other acid reducers

Ask a doctor before use if you have
• Had heartburn over 3 months. This may be a sign of a more serious condition.
• Heartburn with **lightheadedness, sweating or dizziness**, chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• Frequent **chest pain**
• Frequent wheezing, particularly with heartburn
• Unexplained weight loss • nausea or vomiting
• Stomach pain • kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
• Your heartburn continues or worsens
• You need to take this product for more than 14 days

Sanofi

Distributed by: Chatham, Inc., a Sanofi Company
P.O. Box 2219
Chattanooga, TN 37409-0219
©2019 Origin Mexico
0140036-01

Tips for managing heartburn

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

Drug Facts (continued)

Directions
• Adults and children 12 years and over: swallow 1 tablet with a glass of water to relieve symptoms.
• To prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
• Can be used up to twice daily (do not take more than 2 tablets in 24 hours)
• Children under 12 years: ask a doctor

Tips for managing heartburn
• Do not lie flat or bend over soon after eating
• Do not eat late at night, or just before bedtime
• Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

Questions? call 1-800-833-1610 or visit www.zantac.com

Drug Facts (continued)

Active ingredient (in each tablet) Purpose
Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer

Uses
• Relieves heartburn associated with acid indigestion and sour stomach
• Prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
Do not use
• If you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• With other acid reducers

Ask a doctor before use if you have
• Had heartburn over 3 months. This may be a sign of a more serious condition.
• Heartburn with **lightheadedness, sweating or dizziness**, chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• Frequent **chest pain**
• Frequent wheezing, particularly with heartburn
• Unexplained weight loss • nausea or vomiting
• Stomach pain • kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
• Your heartburn continues or worsens
• You need to take this product for more than 14 days

Sanofi

Distributed by: Chatham, Inc., a Sanofi Company
P.O. Box 2219
Chattanooga, TN 37409-0219
©2019 Origin Mexico
0140036-01

Tips for managing heartburn

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

Sanofi Regulatory Specifications		Verified Date: 02/28/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 15% H 0%

Zantac 150 78ct Carton

03.14.19

ZAN_150_78ct_Bonus_CTN_0140072-01_Spain_2019_FDA_DSN

MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

BONUS! 13 FREE TABLETS
BONUS! 13 FREE TABLETS

MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

PREVENTS & RELIEVES HEARTBURN associated with acid indigestion and sour stomach

78 TABLETS (78 DOSES)

Drug Facts (continued)

Active ingredient (in each tablet) Purpose
Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer

Uses
• relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
Do not use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• with other acid reducers

Ask a doctor before use if you have
• had heartburn over 3 months. This may be a sign of a more serious condition.
• heartburn with **lightheadedness, sweating or dizziness**, chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• frequent chest pain
• frequent wheezing, particularly with heartburn
• unexplained weight loss • nausea or vomiting
• stomach pain • kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
• your heartburn continues or worsens
• you need to take this product for more than 14 days

Drug Facts (continued)

Other information
• do not use if printed foil under bottle cap is open or torn
• store at 20°-25°C (68°-77°F)
• avoid excessive heat or humidity
• this product is sodium and sugar free

Inactive ingredients
microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

Directions
• adults and children 12 years and over: swallow 1 tablet with a glass of water to relieve symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages
• to prevent symptoms, swallow 1 tablet with a glass of water that cause heartburn
• can be used up to twice daily (do not take more than 2 tablets in 24 hours)
• children under 12 years: ask a doctor

Tips for managing heartburn
• Do not eat late at night, or just before bedtime
• Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
• You need to take this product for more than 14 days

Tips for managing heartburn

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

0140072-01

©2019 Origin Spain
Chattanooga, TN 37409-0219
P.O. Box 2219
Distributed by: Chatterm, Inc., a Sanofi Company

SANOFI

Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

More tips on bottom panel.

Drug Facts (continued)

Other information
• do not use if printed foil under bottle cap is open or torn
• store at 20°-25°C (68°-77°F)
• avoid excessive heat or humidity
• this product is sodium and sugar free

Inactive ingredients
microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

Directions
• adults and children 12 years and over: swallow 1 tablet with a glass of water to relieve symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages
• to prevent symptoms, swallow 1 tablet with a glass of water that cause heartburn
• can be used up to twice daily (do not take more than 2 tablets in 24 hours)
• children under 12 years: ask a doctor

Tips for managing heartburn
• Do not eat late at night, or just before bedtime
• Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
• You need to take this product for more than 14 days

Drug Facts (continued)

Active ingredient (in each tablet) Purpose
Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer

Uses
• relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
Do not use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• with other acid reducers

Ask a doctor before use if you have
• had heartburn over 3 months. This may be a sign of a more serious condition.
• heartburn with **lightheadedness, sweating or dizziness**, chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
• frequent chest pain
• frequent wheezing, particularly with heartburn
• unexplained weight loss • nausea or vomiting
• stomach pain • kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
• your heartburn continues or worsens
• you need to take this product for more than 14 days

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 15% H 0%

Zantac 150 1ct Sachet/Pouch Dispenser (80ct)
03.18.19
ZAN_150_80ct_Dispenser_0140081-01_Mex_2019_FDA_DSN

Sanofi Regulatory Specifications		Verified Date: 03/18/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	11.5 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	9.25 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	7 point type
Body Text	Font name: Helvetica Neue 57 Condensed	7 point type
Bullets	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets on same line and of statement separated from bulleted statement by two ems		
Spacing of the hairline from edge of box - i.e. Minimum of 1 space either side of Drug Facts Box		
YES		
Tracing	0 to 8 pt.	Horizontal Scale: 97%-100%
Leading (Minimum space in body of Drug Facts)	6.5 pt.	Maximum Characters/Inch: 30
Bursters	2.5 pt.	Vertical: 5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		
.0903 in		
PDP dimensions (in inches)		
H 4.00 in W 4.75 in		
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		
V 13 pt. H 0.00 pt.		
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		
V 94 pt. H 0.00 pt.		
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
V 15% H 0%		
Statement of Identity (pt.) (Divided by Largest Logo Copy pt.)		



Zantac 150 1ct Sachet/Pouch Dispenser (80ct)
03.12.19
ZAN_150_80ct_Dispenser_0140080-01_Spain_2019_FDA_DSN3

Sanofi Regulatory Specifications		Verified Date: 03/12/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	11.5 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	9.25 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	7 point type
Body Text	Font name: Helvetica Neue 57 Condensed	7 point type
Bullets	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets on same line and of statement separated from bulleted statement by two ems		
Spacing of the hairline from edge of box - i.e. Minimum of 1 space either side of Drug Facts Box		
YES		
Tracing	0 to 8 pt.	Horizontal Scale: 97%-100%
Leading (Minimum space in body of Drug Facts)	6.5 pt.	Maximum Characters/Inch: 30
Bursters	2.5 pt.	Vertical: 5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		
.0903 in		
PDP dimensions (in inches)		
H 4.00 in W 4.75 in		
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		
V 13 pt. H 0.00 pt.		
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		
V 94 pt. H 0.00 pt.		
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
V 15% H 0%		



Zantac 150 90ct Carton

03.14.19

ZAN_150_90ct_Bonus_CTN_0140038-01_Mex_2019_FDA_DSN

MAXIMUM STRENGTH
Zantac
 Ranitidine Tablets 150 mg/Acid Reducer
150

VALUE SIZE 90 TABLETS

MAXIMUM STRENGTH
Zantac
 Ranitidine Tablets 150 mg/Acid Reducer
150

90 TABLETS
(90 DOSES)



PREVENTS & RELIEVES
HEARTBURN associated with
 acid indigestion and sour stomach

90 TABLETS
(90 DOSES)



Drug Facts (continued)

Active ingredient (in each tablet) Purpose
 Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer

Uses
 • relieves heartburn associated with acid indigestion and sour stomach
 • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
 • **Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers
 • Do not use if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 • Ask a doctor before use if you have heartburn with heartburn, frequent wheezing, particularly with heartburn, unexplained weight loss, • nausea or vomiting, stomach pain, • kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
 • your heartburn continues or worsens
 • you need to take this product for more than 14 days

Drug Facts (continued)

Other information
 • do not use if printed foil under bottle cap is open or torn
 • store at 20°-25°C (68°-77°F)
 • avoid excessive heat or humidity
 • this product is sodium and sugar free

Inactive ingredients
 microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

Questions? call 1-800-633-1610 or visit www.zantac.com

Tips for managing heartburn
 • Do not eat late at night, or just before bedtime
 • Do not lie flat or bend over soon after eating
 • Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

MOORE TIPS ON BOTTOM PANEL

Sanofi

Distributed by: Chatham, Inc., a Sanofi Company
 P.O. Box 2219
 Chattanooga, TN 37409-0219
 ©2019 Origin Mexico
 0140038-01

Tips for managing heartburn

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

MAXIMUM STRENGTH
Zantac
 Ranitidine Tablets 150 mg/Acid Reducer
150

90 TABLETS
(90 DOSES)

6 81421 03143 1

LOT

EXP

ZANTAC does not supply store brands

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.) / Logo/Largest Copy (pt.)		V 15% H 0%

Zantac 150 90ct Carton

03.14.19

ZAN_150_90ct_Bonus_CTN_0140037-01_Spain_2019_FDA_DSN

Read the directions and warnings before use. Keep the carton.

It contains important information including tips for managing heartburn.

Tips for managing heartburn

- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

SANOFI

Distributed by: Chatham, Inc., a Sanofi Company
P.O. Box 2219
Chattanooga, TN 37409-0219
©2019 Origin Spain
0140037-01

0140037-01

Read the directions and warnings before use. Keep the carton.

It contains important information including tips for managing heartburn.

ZANTAC does not supply store brands

LOT

EXP

MORE TIPS ON BOTTOM PANEL

Tips for managing heartburn

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

Questions? call 1-800-633-1610 or visit www.zantac.com

Inactive ingredients
microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

Other information

- this product is sodium and sugar free
- avoid excessive heat or humidity
- store at 20°-25°C (68°-77°F)
- do not use if printed foil under bottle cap is open or torn

Directions
adults and children 12 years and over:
• to relieve symptoms, swallow 1 tablet with a glass of water
• to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Warnings
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- Ask a doctor before use if you have had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness; chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness; frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting
- stomach pain • kidney disease

Do not use
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have
• had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating or dizziness; chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness; frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting
- stomach pain • kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.
- your heartburn continues or worsens
- you need to take this product for more than 14 days

6 81421 03143 1

90 TABLETS (90 DOSES)

Zantac 150[®]
Ranitidine Tablets 150 mg/Acid Reducer

MAXIMUM STRENGTH

Zantac

Ranitidine Tablets 150 mg/Acid Reducer **150[®]**

VALUE SIZE 90 TABLETS

VALUE SIZE 90 TABLETS

MAXIMUM STRENGTH

Zantac

Ranitidine Tablets 150 mg/Acid Reducer **150[®]**

PREVENTS & RELIEVES

HEARTBURN associated with acid indigestion and sour stomach

90

TABLETS
(90 DOSES)

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.)	Logo/Largest Copy (pt.)	V 15% H 0%

Zantac 150 Twin Card (140ct)

03.25.19

ZAN_150_TWIN_140CT_Card_0140082-01_2019_FDA_DSN5

Twin Pack 2 Bottles x 70ct = 140ct

MAXIMUM STRENGTH
Zantac
 Ranitidine Tablets 150 mg / Acid Reducer **150**[®]



140
 TABLETS
 (140 DOSES)

**PREVENTS &
 RELIEVES
 HEARTBURN**
 associated with
 acid indigestion
 and sour stomach



KO Varnish
 Lot Code

Tips for managing heartburn

- Do not lie flat or bend over soon after eating
- Do not eat late at night, or just before bedtime
- Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
- Eat slowly and do not eat big meals
- If you are overweight, lose weight
- If you smoke, quit smoking
- Raise the head of your bed
- Wear loose fitting clothing around your stomach

Read the directions and warnings before use. Keep the card. It contains important information including tips for managing heartburn.

SANOFI

Distributed by: Chatham, Inc., a Sanofi Company
 P.O. Box 2219
 Chattanooga, TN 37409-0219
 ©2019
 0140082-01

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other information

- do not use if printed foil under bottle cap is open or torn
- store at 20-25°C (68-77°F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

Inactive ingredients hypromellose, magnesium stearate, microcrystalline cellulose, synthetic red iron oxide, titanium dioxide, triacetin

Questions? call 1-800-633-1610 or visit www.zantac.com

Drug Facts

Active ingredient (in each tablet) Purpose
 Ranitidine hydrochloride 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with high blood pressure, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss • nausea or vomiting
- stomach pain • kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Sanofi Regulatory Specifications		Verified Date: 03/25/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	11.5 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	9.25 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	7 point type
Body Text	Font name: Helvetica Neue 57 Condensed	7 point type
Bullets	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.5 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		.0903 in
PDP dimensions (in inches)		H 6.75 in W 7.75 in
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 13 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 94 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (as measured by) Largest Logo Copy (pt.)		V 15% H 0%

Zantac 150 Cool Mint 8ct Carton

03.28.19

ZAN_150_CoolMint_8ct_CTN_0140069-01_Mex_2019_FDA_DSN3



Sanofi Regulatory Specifications		Verified Date: 03/28/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 15% H 0%

Zantac 150 Cool Mint 24ct Carton

03.14.19

ZAN_150_CoolMint_24ct_CTN_0140068-01_Mex_2019_FDA_DSN



Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	5.9 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 15% H 0%

Zantac 150 Cool Mint 32ct Carton

03.06.19

ZAN_150_CoolMint_32ct_Bonus_CTN_0140075-01_Mex_2019_FDA_DSN2



Sanofi Regulatory Specifications		Verified Date: 03/06/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	5.9 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (if not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (if not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		V 15% H 0%

Zantac 150 Coolmint 40ct Carton

02.28.19

ZAN_150_Coolmint_40ct_CTN_0140063-01_MEX_2019_FDA_DSN.ai

Sanofi Regulatory Specifications		Verified Date: 02/28/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.)	Ratio of Statement of Identity to Logo/Largest Copy (pt.)	V 15% H 0%

Zantac 150 Coolmint 50ct Carton

03.14.19

ZAN_150_CoolMint_50ct_CTN_0140067-01_Mex_2019_FDA_DSN

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150
50 TABLETS (50 DOSES)

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150
50 TABLETS (50 DOSES)

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

Releases a Cooling Sensation in Mouth & Throat

PREVENTS & RELIEVES HEARTBURN associated with acid indigestion and sour stomach

SUGAR FREE **50** TABLETS (50 DOSES)

Drug Facts

Active ingredient (in each tablet) Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer

Purpose If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water • to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn

Warnings • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers

Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers

Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with **lightheadedness, sweating or dizziness**; chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days

Drug Facts (continued)

Other information • do not use if printed foil under bottle cap is open or torn • store at 20-25°C (68-77°F) • avoid excessive heat or humidity • this product is sodium and sugar free

Inactive ingredients FD&C Blue No. 1, flavors, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polypropylene glycol, titanium dioxide

Questions? call 1-800-833-1610 or visit www.zantac.com

Tips for managing heartburn • Do not lie flat or bend over soon after eating • Do not eat late at night, or just before bedtime • Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

Tips for managing heartburn • Eat slowly and do not eat big meals • If you are overweight, lose weight • If you smoke, quit smoking • Raise the head of your bed • Wear loose fitting clothing around your stomach

0140067-01
©2019 Origin Mexico
Chattanooga, TN 37409-0219
Distributed by: Chatterm, Inc., a Sanofi Company

Read the directions and warnings before use. Keep the carton. It contains important information including tips for managing heartburn.

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150
50 TABLETS (50 DOSES)

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150
50 TABLETS (50 DOSES)

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.) (Largest Logo Copy (pt.))		V 15% H 0%

Zantac 150 Coolmint 65ct Carton

03.14.19

ZAN_150_Coolmint_65ct_CTN_0140066-01_Mex_2019_FDA_DSN

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.) / Logo/Largest Copy (pt.)		V 15% H 0%

Zantac 150 Coolmint 78ct Carton

03.14.19

ZAN_150_Coolmint_78ct_Bonus_CTN_0140065-01_Mex_2019_FDA_DSN

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

BONUS! 13 FREE TABLETS
BONUS! 13 FREE TABLETS

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

Releases a Cooling Sensation in Mouth & Throat™

PREVENTS & RELIEVES HEARTBURN associated with acid indigestion and sour stomach

78 TABLETS (78 DOSES)

SUGAR FREE

Drug Facts (continued)

Active ingredient (in each tablet) Purpose
Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer

Uses
• relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• with other acid reducers

Ask a doctor before use if you have
• had heartburn over 3 months. This may be a sign of a more serious condition.
• heartburn with **lightheadedness, sweating or dizziness**; chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Do not use
• frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain • kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if
• your heartburn continues or worsens
• you need to take this product for more than 14 days

Drug Facts (continued)

Directions
• adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water
• to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
• can be used up to twice daily (do not take more than 2 tablets in 24 hours)
• do not chew tablet
• children under 12 years: ask a doctor

Other information
• do not use if printed foil under bottle cap is open or torn
• store at 20-25°C (68-77°F) • avoid excessive heat or humidity
• this product is sodium and sugar free

Inactive ingredients FD&C Blue No. 1, flavors, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polypropylene glycol, sucralose, titanium dioxide

Questions? call 1-800-833-1610 or visit www.zantac.com

Tips for managing heartburn
• Do not lie flat or bend over soon after eating
• Do not eat late at night, or just before bedtime
• Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables

Tips for managing heartburn
• Eat slowly and do not eat big meals
• If you smoke, quit smoking
• Raise the head of your bed
• Wear loose fitting clothing around your stomach

0140065-01
©2019 Origin Mexico
Chattanooga, TN 37409-0219
P.O. Box 2219
Distributed by: Chatterm, Inc., a Sanofi Company

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150

78 TABLETS (78 DOSES)

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.

Principal Display Panel Info	
Font size of Net Quantity of Contents (Smallest character height in inches)	0.088 in
PDP dimensions (in millimeters)	H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")	V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")	V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP	
Statement of Identity (pt.)	Ratio of Statement of Identity to Logo/Largest Copy (pt.)
Reference ID: 4493200	V 15% H 0%

Zantac 150 Coolmint 90ct Carton

03.14.19

ZAN_150_Coolmint_90ct_Value_CTN_0140064-01_Mex_2019_FDA_DSN

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150
90 TABLETS (90 DOSES)

VALUE SIZE 90 TABLETS
VALUE SIZE 90 TABLETS

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150
90 TABLETS (90 DOSES)

Releases a Cooling Sensation in Mouth & Throat

PREVENTS & RELIEVES HEARTBURN associated with acid indigestion and sour stomach

SUGAR FREE

90 TABLETS (90 DOSES)

Drug Facts
Ranitidine 150 mg (as ranitidine hydrochloride 168 mg) Acid reducer
Active ingredient (in each tablet) Purpose
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
adults and children 12 years and over:
• to relieve symptoms, swallow 1 tablet with a glass of water
• to prevent symptoms, swallow 1 tablet with a glass of water
30 to 60 minutes before eating food or drinking beverages that cause heartburn
• can be used up to twice daily (do not take more than 2 tablets in 24 hours)
• do not chew tablet
• children under 12 years: ask a doctor

Other information
• store at 20-25°C (68-77°F) • avoid excessive heat or humidity
• do not use if printed foil under bottle cap is open or torn
• this product is sodium and sugar free

Inactive ingredients FD&C Blue No. 1, flavors, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polypropylene glycol, titanium dioxide

Questions? call 1-800-633-1610 or visit www.zantac.com

Tips for managing heartburn
• Do not eat late at night, or just before bedtime
• Do not lie flat or bend over soon after eating
• Certain foods or drinks are more likely to cause heartburn, such as rich, spicy, fatty, and fried foods, chocolate, caffeine, alcohol, even some fruits and vegetables
• You need to take this product for more than 14 days
• your heartburn continues or worsens
• Stop use and ask a doctor if prescription drugs. Acid reducers may interact with certain prescription drugs.
• stomach pain • kidney disease
• frequent wheezing, particularly with heartburn
• unexplained weight loss • nausea or vomiting
• frequent chest pain
• pain spreading to arms, neck or shoulders; or lightheadedness
• chest pain or shoulder pain with shortness of breath; sweating; heartburn with lightheadedness, sweating or dizziness; serious condition.
• had heartburn over 3 months. This may be a sign of a more serious condition.
• Ask a doctor before use if you have
• with other acid reducers
• serious condition. See your doctor.
• blood, or bloody or black stools. These may be signs of a serious condition.
• if you have trouble or pain swallowing food, vomiting with acid reducers
• Do not use
• if you have trouble or pain swallowing food, vomiting with acid reducers
• Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
Warnings
• stomach brought on by eating or drinking certain foods and prevents heartburn associated with acid indigestion and sour stomach
• relieves heartburn associated with acid indigestion and sour stomach
Uses
• relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and

Read the directions and warnings before use. Keep the carton.
It contains important information including tips for managing heartburn.

Tips for managing heartburn
• Eat slowly and do not eat big meals
• If you are overweight, lose weight
• If you smoke, quit smoking
• Raise the head of your bed
• Wear loose fitting clothing around your stomach

Distributed by: Chatham, Inc., a Sanofi Company
P.O. Box 2219
Chattanooga, TN 37409-0219
©2019 Origin Mexico
0140064-01

6 81421 03230 8

Drug Facts (continued)

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150
90 TABLETS (90 DOSES)

COOL MINT TABLETS
MAXIMUM STRENGTH
Zantac
Ranitidine Tablets 150 mg/Acid Reducer
150
90 TABLETS (90 DOSES)

Sanofi Regulatory Specifications		Verified Date: 03/14/19
Drug Facts Info		
Drug Facts (Title)	Font name: Helvetica Neue 77 Bold Condensed Oblique	10 point type
Drug Facts (continued)	Font name: Helvetica Neue 77 Bold Condensed Oblique	9 point type
Headings	Font name: Helvetica Neue 77 Bold Condensed Oblique	8 point type
Subheadings	Font name: Helvetica Neue 77 Bold Condensed	6 point type
Body Text	Font name: Helvetica Neue 57 Condensed	6 point type
Bullets	Font name: Helvetica Neue 57 Condensed	5 point type
Bullets on same lines: end of statement separated from bulleted statement by two ems		YES
Spacing of the hairlines from edge of box – i.e. Minimum of 1 space either side of Drug Facts Box		YES
Tracking:	0 to 8 pt. Horizontal Scale:	97%-102%
Leading (Minimum space in body of Drug Facts):	6.0 pt. Maximum Characters/Inch:	39
Barlines:	2.5 pt. Hairlines:	0.5 pt.
Principal Display Panel Info		
Font size of Net Quantity of Contents (Smallest character height in inches)		0.088 in
PDP dimensions (in millimeters)		H 89mm W 97mm
Font size of Statement of Identity (If not live text, to be measured in Helvetica capital "M")		V 11.5 pt. H 0.00 pt.
Font size of Logo/Largest Copy on PDP (If not live text, to be measured in Helvetica capital "M")		V 82 pt. H 0.00 pt.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		V 15% H 0%
Statement of Identity (pt.) / Logo/Largest Copy (pt.)		V 15% H 0%

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
09/18/2019 02:41:47 PM