


Allegra-D 12 Hour Allergy and Congestion

04.30.14


ALG-D_12hr_30ct_CTN_NI_2014



Allegra-D 12 Hour Allergy and Congestion


04.30.14

ALG-D_12hr_20ct_CTN_NI_2014





04.30.14

Allegra-D 12 Hour Allergy and
04.30.14
ALG-D_12hr_10oct_CTN_NI_201









04.30.14

ALG-D 24hr 15ct CTN NI 2014







Allegra-D®

Allergy & Congestion

180 mg / antihistamine
240 mg / nasal decongestant

Extended Release Tablets

Indoor and Outdoor Allergies

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away.

Directions
 Adults and children 12 years of age and older:
 Take 1 tablet with a glass of water every 24 hours on a regular schedule, do not take more than 1 tablet in 24 hours.

Other Information
 Each tablet contains **sodium 33 mg** (about 0.5% of the daily sodium allowance for adults).

Warnings
 • Do not take more than 1 tablet in 24 hours.
 • Do not take with other products containing pseudoephedrine or other nasal decongestants.
 • Do not take with other products containing antihistamines or other allergy medications.
 • Do not take with other products containing blood pressure medications.
 • Do not take with other products containing MAO inhibitors (e.g., phenelzamine, tranylcypromine, or others).
 • Do not take with other products containing alcohol.
 • Do not take with other products containing grapefruit juice.
 • Do not take with other products containing St. John's Wort.
 • Do not take with other products containing warfarin or other blood thinners.
 • Do not take with other products containing lithium.
 • Do not take with other products containing digoxin.
 • Do not take with other products containing theophylline.
 • Do not take with other products containing insulin.
 • Do not take with other products containing thyroid medicine.
 • Do not take with other products containing birth control pills.
 • Do not take with other products containing antidepressants.
 • Do not take with other products containing sedatives.
 • Do not take with other products containing muscle relaxants.
 • Do not take with other products containing anti-anxiety drugs.
 • Do not take with other products containing painkillers.
 • Do not take with other products containing antibiotics.
 • Do not take with other products containing antifungals.
 • Do not take with other products containing antivirals.
 • Do not take with other products containing chemotherapy drugs.
 • Do not take with other products containing immunosuppressants.
 • Do not take with other products containing hormone therapy.
 • Do not take with other products containing cancer drugs.
 • Do not take with other products containing heart disease medicines.
 • Do not take with other products containing diabetes medicines.
 • Do not take with other products containing thyroid medicine.
 • Do not take with other products containing Parkinson's disease medicine.
 • Do not take with other products containing glaucoma medicine.
 • Do not take with other products containing eye drops.
 • Do not take with other products containing ear drops.
 • Do not take with other products containing inhalers.
 • Do not take with other products containing nasal sprays.
 • Do not take with other products containing skin creams.
 • Do not take with other products containing eye ointments.
 • Do not take with other products containing ear ointments.
 • Do not take with other products containing rectal suppositories.
 • Do not take with other products containing vaginal suppositories.
 • Do not take with other products containing birth control pills.
 • Do not take with other products containing antidepressants.
 • Do not take with other products containing sedatives.
 • Do not take with other products containing muscle relaxants.
 • Do not take with other products containing anti-anxiety drugs.
 • Do not take with other products containing painkillers.
 • Do not take with other products containing antibiotics.
 • Do not take with other products containing antifungals.
 • Do not take with other products containing antivirals.
 • Do not take with other products containing chemotherapy drugs.
 • Do not take with other products containing immunosuppressants.
 • Do not take with other products containing hormone therapy.
 • Do not take with other products containing cancer drugs.
 • Do not take with other products containing heart disease medicines.
 • Do not take with other products containing diabetes medicines.
 • Do not take with other products containing thyroid medicine.
 • Do not take with other products containing Parkinson's disease medicine.
 • Do not take with other products containing glaucoma medicine.
 • Do not take with other products containing eye drops.
 • Do not take with other products containing ear drops.
 • Do not take with other products containing inhalers.
 • Do not take with other products containing nasal sprays.
 • Do not take with other products containing skin creams.
 • Do not take with other products containing eye ointments.
 • Do not take with other products containing ear ointments.
 • Do not take with other products containing rectal suppositories.
 • Do not take with other products containing vaginal suppositories.

Drug Facts

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away.

Directions
 Adults and children 12 years of age and older:
 Take 1 tablet with a glass of water every 24 hours on a regular schedule, do not take more than 1 tablet in 24 hours.

Other Information
 Each tablet contains **sodium 33 mg** (about 0.5% of the daily sodium allowance for adults).

Warnings
 • Do not take more than 1 tablet in 24 hours.
 • Do not take with other products containing pseudoephedrine or other nasal decongestants.
 • Do not take with other products containing antihistamines or other allergy medications.
 • Do not take with other products containing blood pressure medications.
 • Do not take with other products containing MAO inhibitors (e.g., phenelzamine, tranylcypromine, or others).
 • Do not take with other products containing alcohol.
 • Do not take with other products containing grapefruit juice.
 • Do not take with other products containing St. John's Wort.
 • Do not take with other products containing warfarin or other blood thinners.
 • Do not take with other products containing lithium.
 • Do not take with other products containing digoxin.
 • Do not take with other products containing theophylline.
 • Do not take with other products containing insulin.
 • Do not take with other products containing thyroid medicine.
 • Do not take with other products containing Parkinson's disease medicine.
 • Do not take with other products containing glaucoma medicine.
 • Do not take with other products containing eye drops.
 • Do not take with other products containing ear drops.
 • Do not take with other products containing inhalers.
 • Do not take with other products containing nasal sprays.
 • Do not take with other products containing skin creams.
 • Do not take with other products containing eye ointments.
 • Do not take with other products containing ear ointments.
 • Do not take with other products containing rectal suppositories.
 • Do not take with other products containing vaginal suppositories.

Drug Facts

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away.

Directions
 Adults and children 12 years of age and older:
 Take 1 tablet with a glass of water every 24 hours on a regular schedule, do not take more than 1 tablet in 24 hours.

Other Information
 Each tablet contains **sodium 33 mg** (about 0.5% of the daily sodium allowance for adults).

Warnings
 • Do not take more than 1 tablet in 24 hours.
 • Do not take with other products containing pseudoephedrine or other nasal decongestants.
 • Do not take with other products containing antihistamines or other allergy medications.
 • Do not take with other products containing blood pressure medications.
 • Do not take with other products containing MAO inhibitors (e.g., phenelzamine, tranylcypromine, or others).
 • Do not take with other products containing alcohol.
 • Do not take with other products containing grapefruit juice.
 • Do not take with other products containing St. John's Wort.
 • Do not take with other products containing warfarin or other blood thinners.
 • Do not take with other products containing lithium.
 • Do not take with other products containing digoxin.
 • Do not take with other products containing theophylline.
 • Do not take with other products containing insulin.
 • Do not take with other products containing thyroid medicine.
 • Do not take with other products containing Parkinson's disease medicine.
 • Do not take with other products containing glaucoma medicine.
 • Do not take with other products containing eye drops.
 • Do not take with other products containing ear drops.
 • Do not take with other products containing inhalers.
 • Do not take with other products containing nasal sprays.
 • Do not take with other products containing skin creams.
 • Do not take with other products containing eye ointments.
 • Do not take with other products containing ear ointments.
 • Do not take with other products containing rectal suppositories.
 • Do not take with other products containing vaginal suppositories.



(b) (4)