

Drug Facts
 Active ingredient (as hydrochloride): Loperamide HCl
 Compare to the active ingredient in Imodium® A-D
 NDC 0602-015-07



USA
 100% Recycled Paper

Loperamide HCl
Soft Gelatin Capsules, 2 mg
 Anti-Diarrheal
 Suitable for adults and children 12 years and over
 48 Softgels*
 *each Liquid-filled capsule contains 2 mg loperamide HCl
 NDC 0602-015-07

Compare to the active ingredient in Imodium® A-D


Loperamide
2 mg Softgels*

Loperamide HCl
Soft Gelatin Capsules, 2 mg
 Anti-Diarrheal
 Suitable for adults and children 12 years and over
 48 Softgels*
 *each Liquid-filled capsule contains 2 mg loperamide HCl
 NDC 0602-015-07

Compare to the active ingredient in Imodium® A-D

Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	14pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	8pt	Headlines:	1.5pt
Body Text:	8pt	Maxlines:	.5pt
Drug Facts (continued):	N/A		
Horizontal Scale: 100%		Average Kerning: 0%	

Drug Facts
 Contains 48 capsules in 1 blister pack.
 NDC 0602-015-07



Drug Facts
 Active ingredient in each capsule:
 Loperamide

Compare to the active ingredient in Includium® A-D

Loperamide HCl
Soft Gelatin Capsules, 2 mg
Anti-Diarrheal

Suitable for adults and children 12 years and over

60 Softgels*
 *each Liquid-filled capsule contains 2 mg loperamide HCl

Labeling Format Information:
 Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique
 Drug Facts: 14pt Leading: 6.5pt
 Header: 8pt Bullets: 5pt
 Subheader: 6pt Barlines: 1.5pt
 Body Text: 6pt Headlines: 5pt
 Drug Facts (continued): N/A
 Horizontal Scale: 100% Average Kerning: 0%

Loperamide HCl
Soft Gelatin Capsules, 2 mg
Anti-Diarrheal

Suitable for adults and children 12 years and over

60 Softgels*
 *each Liquid-filled capsule contains 2 mg loperamide HCl

Loperamide HCl
Soft Gelatin Capsules, 2 mg
Anti-Diarrheal

Suitable for adults and children 12 years and over

60 Softgels*
 *each Liquid-filled capsule contains 2 mg loperamide HCl

Drug Facts
 Active ingredient in each capsule:
 Loperamide


Compare to the active ingredient in Includium® A-D

Loperamide HCl
Soft Gelatin Capsules, 2 mg
Anti-Diarrheal

Suitable for adults and children 12 years and over

60 Softgels*
 *each Liquid-filled capsule contains 2 mg loperamide HCl

Reference ID: 4066942



Drug Facts
 Active ingredient (see ingredients)
 Compare to the active ingredient in Imodium A-D



Loperamide HCl
Soft Gelatin Capsules, 2 mg
 Anti-Diarrheal
 Suitable for adults and children 12 years and over
 72 Softgels*
 *each Liquid-filled capsule contains 2 mg loperamide HCl

Loperamide HCl
Soft Gelatin Capsules, 2 mg
 Anti-Diarrheal
 Suitable for adults and children 12 years and over
 72 Softgels*
 *each Liquid-filled capsule contains 2 mg loperamide HCl

Labeling Format Information:
 Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	1.5pt	Leadings:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	1.5pt
Body Text:	6pt	Headlines:	.5pt
Drug Facts (continued):	N/A		

Horizontal Scale: 100% Average Kerning: 0%

Drug Facts
 (continued)

NOT OFFICIAL SIZE/DIELINE

Drug Facts (continued)

Active ingredient (in each capsule)
Loperamide HCl 1 mg..... Anti-diarrheal

Use controls symptoms of diarrhea, including Traveler's Diarrhea

Warnings Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
 Heart alert: Taking more than directed can cause serious heart problems or death
 Do not use if you have bloody or black stool
 Ask a doctor before use if you have fever, mucus in the stool, or a history of liver disease
 Ask a doctor or pharmacist before use if you are taking antibiotics

When using this product, dizziness, drowsiness, or lightheadedness may occur. Be careful when driving or operating machinery.
 Stop use and ask a doctor if symptoms get worse, diarrhea lasts for more than 2 days
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.
 Find right dose on chart. If possible, use weight to dose; otherwise, use age.

Adults and children 12 years and over: 4 softgels after the first loose stool; 2 softgels after each subsequent loose stool, but no more than 8 softgels in 24 hours
 Children 9-11 years (60-95 lbs): 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool, but no more than 6 softgels in 24 hours
 Children 6-8 years (48-80 lbs): 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool, but no more than 4 softgels in 24 hours
 Children under 6 years (up to 47 lbs): ask a doctor

Other information
 Store at 20°-25°C (68°-77°F). Protect from light
 Avoid excessive heat above 40°C (104°F)
 Do not use if carton or blister unit is open or torn
 See side panel for lot number and expiration date

Manufactured for: Bionpharma Inc.
 600 Alexander Road
 Princeton, NJ 08540

Lot #
 Expiration Date
 Ref # 10/16

Loperamide 1 mg Softgels*

NDC 69452-177-04

Compare to the active ingredient in Imodium® A-D

Loperamide HCl

Soft Gelatin Capsules, 1 mg

Anti-Diarrheal

Suitable for adults and children 6 years and over

6 Softgels*
 *each Liquid-filled capsule contains 1 mg loperamide HCl

3 69452 17704 4

Drug Facts (continued)

Inactive ingredients butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxy 40 hydrogenated castor oil, purified water

Questions or comments? call toll free 1-888-235-2466

®

Carton Size : 0.75" x 3.375" x 4.375"
 Track : 18/10/2016, 24/10/2016, 26/10/2016

Colors Used



Labeling Format Information:			
Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique			
Drug Facts:	14pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	1.5pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	N/A		
Horizontal Scale: 100%		Average Kerning: 0%	

NOT OFFICIAL SIZE/DIELINE

Drug Facts (continued)

Other information

- store at 20°-25°C (68°-77°F); protect from light
- avoid excessive heat above 40°C (104°F)
- see side panel for lot number and expiration date
- do not use if carton or blister unit is open or torn

Inactive ingredients butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxy 40 hydrogenated castor oil, purified water

Questions or comments? call toll free 1-888-235-2466

Manufactured for: Biopharma Inc.
600 Alexander Road,
Princeton, NJ 08540

Drug Facts

Active ingredient (in each capsule)
Loperamide HCl 1 mg.....Anti-diarrhea1

Use controls symptoms of diarrhea, including travelers' diarrhea

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Heart alert: Taking more than directed can cause serious heart problems or death
Do not use ■ if you have blood or black stool
Ask a doctor before use if you have ■ fever ■ hives ■ a history of liver disease
When using this product tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.
Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days
If pregnant or breast-feeding, ask a health professional for use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea.
■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over
4 softgels after the first loose stool; 2 softgels after each subsequent loose stool;

children 9 - 11 years (60 - 95 lbs)
2 softgels after the first loose stool; 1 softgel after each subsequent loose stool;

children 6 - 8 years (45 - 59 lbs)
2 softgels after the first loose stool; 1 softgel after each subsequent loose stool;

children under 6 years (up to 47 lbs)
ask a doctor

Labeling Format Information:

Fonts: Helvetica Condensed Bold, Helvetica Condensed, Helvetica Condensed Bold Oblique

Drug Facts:	14pt	Leading:	6.5pt
Header:	8pt	Bullets:	5pt
Subheader:	6pt	Barlines:	1.5pt
Body Text:	6pt	Hairlines:	.5pt
Drug Facts (continued):	N/A		

Horizontal Scale: 100% Average Kerning: 0%

Compare to the active ingredient in Imodium® A-D

Loperamide HCl Soft Gelatin Capsules, 1 mg Anti-Diarrheal

Suitable for adults and children 6 years and over

72 Softgels*
*each Liquid-filled capsule contains 1 mg loperamide HCl

NDC 69452-177-76

Compare to the active ingredient in Imodium® A-D

Loperamide HCl Soft Gelatin Capsules, 1 mg Anti-Diarrheal

Suitable for adults and children 6 years and over

72 Softgels*
*each Liquid-filled capsule contains 1 mg loperamide HCl

NDC 69452-177-76

Loperamide 1 mg Softgels*

Loperamide 1 mg Softgels*

Loperamide 1 mg Softgels*

Loperamide 1 mg Softgels*

Loperamide 1 mg Softgels*

Loperamide 1 mg Softgels*

Loperamide 1 mg Softgels*

Carton Size : 3" x 3.375" x 4.375"
Track : 18/10/2016, 24/10/2016,
26/10/2016

Reference ID: 4066942

Colors Used

(b) (4)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
03/09/2017