


alli[®]
Orlistat 60mg Capsules
Weight Loss Aid
60 CAPSULES

TAMPER-EVIDENT features for your protection. Do not use if bottle was not in a carton with security seals on both ends that are printed with GSK logo OR foil seal under cap imprinted with "sealed for your protection" OR dark blue band around center of capsule are missing, torn or broken.

Warnings

Organ transplant alert: • do not use if you have had an organ transplant. Orlistat interferes with the medicines used to prevent transplant rejection. **Allergy alert:** do not use if you are allergic to any of the ingredients in orlistat capsules. Do not use • If you are taking cyclosporine • if you have been diagnosed with problems absorbing food • if you are not overweight **Ask a doctor before use if you have ever had:** • gallbladder problems • kidney stones • pancreatitis **Ask a doctor or pharmacist if you are:** • taking any of the following prescription medications. Some prescription doses may need to be adjusted, or some medications may not work as well when used with orlistat. • anticoagulants (blood thinning medicine) • amiodarone (heart medicine) • diabetes medicine • thyroid disease medicine • seizure medicine • antiretrovirals (HIV medicine) • taking other weight loss products **When using this product:** • take a multivitamin once a day, at bedtime. Orlistat can reduce the absorption of some vitamins. • bowel changes may occur if you eat too much fat. You may get: • gas with oily spotting • loose stools • more frequent stools that may be hard to control **Stop use and ask a doctor if:** • you develop itching, yellow eyes or skin, dark urine or loss of appetite. There have been rare reports of liver injury in people taking orlistat. • severe or continuous abdominal pain occurs. This may be a sign of a serious medical condition. • you are taking medicine for seizures and your seizures happen more often or get worse. **If pregnant or breast feeding, do not use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • For overweight adults 18 years and older • take 1 capsule with each meal containing fat • do not take more than 3 capsules daily

Other information • store at 20-25°C (68-77°F) • protect drug from excessive light, humidity and temperatures over 30°C (86°F)

Questions or comments?
Call toll-free 1-800-671-2554
Distributed by: GSK Consumer Healthcare
Trade dress are owned by or licensed to: Warren, IN 07055
© 2015 GSK group of companies
All rights reserved. All logo reserved.

LOT/EXPIRATION: 17-008CHFS



alli[®]
Orlistat 60mg Capsules
Weight Loss Aid
120 CAPSULES

TAMPER-EVIDENT features for your protection. Do not use if bottle was not in a carton with security seals on both ends that are printed with GSK logo OR foil seal under cap imprinted with "sealed for your protection" OR dark blue band around center of capsule are missing, torn or broken.

Warnings

Organ transplant alert: • do not use if you have had an organ transplant. Orlistat interferes with the medicines used to prevent transplant rejection. **Allergy alert:** do not use if you are allergic to any of the ingredients in orlistat capsules. **Do not use** • if you are taking cyclosporine • if you have been diagnosed with problems absorbing food • if you are not overweight **Ask a doctor before use if you have ever had:** • gallbladder problems • kidney stones • pancreatitis **Ask a doctor or pharmacist if you are** • taking any of the following prescription medications: Some prescription doses may need to be adjusted, or some medications may not work as well when used with orlistat. • anticoagulants (blood thinning medicine) • amiodarone (heart medicine) • diabetes medicine • thyroid disease medicine • seizure medicine • antiretrovirals (HIV medicine) • taking other weight loss products **When using this product:** • **Take a multivitamin once a day, at bedtime.** Orlistat can reduce the absorption of some vitamins. • bowel changes may occur if you eat too much fat. You may get: • gas with oily spotting • loose stools • more frequent stools that may be hard to control **Stop use and ask a doctor if** • you develop itching, yellow eyes or skin, dark urine or loss of appetite. There have been rare reports of liver injury in people taking orlistat. • severe or continuous abdominal pain occurs. This may be a sign of a serious medical condition. • you are taking medicine for seizures and your seizures happen more often or get worse. **If pregnant or breast feeding, do not use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • for overweight adults 18 years and older • take 1 capsule with each meal containing fat • do not take more than 3 capsules daily.

Other information • store at 20-25°C (68-77°F) • protect drug from excessive light, humidity and temperatures over 30°C (86°F)

Questions or comments? call toll-free 1-800-671-2554
Trademarks are owned by or licensed to GSK Consumer Healthcare, Warren, NJ 07059
Distributed by GSK Consumer Healthcare to the GSK group of companies.

© 2015 GSK group of companies or its licensor. All rights reserved.

LOT/EXPIRATION:

17-010CHFS



alli[®]
Orlistat 60mg Capsules
Weight Loss Aid
170 CAPSULES


TAMPER-EVIDENT features for your protection. Do not use if bottle was not in a carton with security seals on both ends that are printed with GSK logo OR foil seal under cap imprinted with "sealed for your protection" OR dark blue band around center of capsule are missing, torn or broken.

Warnings

Organ transplant alert: • do not use if you have had an organ transplant. Orlistat interferes with the medicines used to prevent transplant rejection. **Allergy alert:** do not use if you are allergic to any of the ingredients in orlistat capsules. Do not use • if you are taking cyclosporine • if you have been diagnosed with problems absorbing food. • if you are not overweight. **Ask a doctor before use if you have ever had:** • gallbladder problems • kidney stones • pancreatitis **Ask a doctor or pharmacist if you are** • taking any of the following prescription medications. Some prescription doses may need to be adjusted, or some medications may not work as well when used with orlistat. • anticoagulants (blood thinning medicine) • amiodarone (heart medicine) • diabetes medicine • thyroid disease medicine • seizure medicine • antiretrovirals (HIV medicine) • taking other weight loss products. **When using this product:** • **Take a multivitamin once a day, at bedtime.** Orlistat can reduce the absorption of some vitamins. • bowel changes may occur if you eat too much fat. You may get: • gas with oily spotting • loose stools • more frequent stools that may be hard to control **Stop use and ask a doctor if** • you develop itching, yellow eyes or skin, dark urine or loss of appetite. There have been rare reports of liver injury in people taking orlistat. • severe or continuous abdominal pain occurs. This may be a sign of a serious medical condition. • you are taking medicine for seizures and your seizures happen more often or get worse. **If pregnant or breast feeding, do not use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions • for overweight adults 18 years and older •** take 1 capsule with each meal containing fat • do not take more than 3 capsules daily. **Other information •** store at 20-25°C (68-77°F) • protect drug from excessive light, humidity and temperatures over 30°C (86°F) **Questions or comments? call toll-free 1-800-671-2554** Trademarks are owned by or licensed to GSK Consumer Healthcare, a division of the GSK group of companies. © 2015 GSK group of companies or its licensor. All rights reserved.

LOT/EXPIRATION:

17-013CHF5



NDC 0135-0461-06

alli[®]

The Refill Pack includes:

- Read Me First brochure
- Up to 56 day supply (170 capsules)

For more information, visit www.myalli.com

How is this different than the Starter Pack?

The Starter Pack contains a convenient carrying case and additional materials designed to support healthy lifestyle changes.

Drug Facts	Purpose
Active ingredient (in each sealed capsule) Orlistat (60mg)	Weight loss aid
Use	
• for weight loss in overweight adults, 18 years and older, when used along with a reduced-calorie and low-fat diet	
Warnings	
Organ transplant alert: • do not use if you have had an organ transplant. Orlistat interferes with the medicines used to prevent transplant rejection.	
Allergy alert: • do not use if you are allergic to any of the ingredients in orlistat capsules	
Do not use	
• if you are taking cyclosporine	
• if you have been diagnosed with problems absorbing food	
• if you are not overweight	
Ask a doctor before use if you have ever had	
• gallbladder problems • kidney stones • pancreatitis	
Ask a doctor or pharmacist if you are	
• taking any of the following prescription medications. Some prescription doses may need to be adjusted, or some medications may not work as well when used with orlistat.	
• anticoagulants (blood thinning medicine)	
• amiodarone (heart medicine)	
• diabetes medicine	
• thyroid disease medicine	
• seizure medicine	
• antiretroviral (HIV medicine)	
• taking other weight loss products	
When using this product	
• take a multivitamin once a day, at bedtime. Orlistat can reduce the absorption of some vitamins.	
• follow a well-balanced, reduced-calorie, low-fat diet. Try starting this diet before taking orlistat.	
• orlistat works by preventing the absorption of some of the fat you eat. The fat passes out of your body, so you may have bowel changes. You may get:	
• gas with oily spotting	
• loose stools	
• more frequent stools that may be hard to control	
• eating a low-fat diet lowers the chance of having these bowel changes	
• for every 5 pounds you lose from diet alone, orlistat can help you lose 2-3 pounds more. In studies, most people lost 6-10 pounds over 6 months.	
Stop use and ask a doctor if	
• you develop itching, yellow eyes or skin, dark urine or loss of appetite.	
• there have been rare reports of liver injury in people taking orlistat.	
• severe or continuous abdominal pain occurs. This may be a sign of a serious medical condition.	
• you are taking medicine for seizures and your seizures happen more often or get worse	

Drug Facts (continued)	HL/WL
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	4' 10" 129 lbs.
Directions	4' 11" 133 lbs.
• read the enclosed brochure for other important information	5' 0" 138 lbs.
• diet and exercise are the starting points for any weight loss program. Try these first before adding orlistat. Check with your doctor before starting any exercise program.	5' 1" 143 lbs.
• to see if orlistat capsules are right for you, find your height on the chart to the right. You may consider starting a weight loss program with orlistat if your weight is the same or more than the weight shown for your height.	5' 2" 147 lbs.
• for overweight adults 18 years and older:	5' 3" 152 lbs.
• take 1 capsule with each meal containing fat	5' 4" 157 lbs.
• do not take more than 3 capsules daily	5' 5" 162 lbs.
• use with a reduced-calorie, low-fat diet and exercise program until you reach your weight loss goal. Most weight loss occurs in the first 6 months.	5' 6" 167 lbs.
• if you stop taking orlistat, continue with your diet and exercise program	5' 7" 172 lbs.
• if you start to regain weight after you stop taking orlistat, you may need to start taking orlistat again along with your diet and exercise program	5' 8" 177 lbs.
• take a multivitamin once a day, at bedtime, when using orlistat	5' 9" 182 lbs.
Other information	5' 10" 188 lbs.
• store at 20°-25°C (68°-77°F)	5' 11" 193 lbs.
• protect drug from excessive light, humidity and temperatures over 30°C (86°F)	5' 0" 189 lbs.
Inactive ingredients	5' 1" 194 lbs.
FD&C Blue No. 2, croscellose, gelatin, iron oxide, microcrystalline cellulose, polydioxane, sodium lauryl sulfate, sodium starch glycolate, talc, titanium dioxide	5' 2" 200 lbs.
Questions or comments? call toll-free 1-800-471-2954	5' 3" 206 lbs.
	5' 4" 211 lbs.
	5' 5" 217 lbs.

For more information and to learn more about alli, visit us at www.myalli.com.

Trademarks are owned by or licensed to the GSK group of companies.

Distributed by:
GSK Consumer Healthcare
Warren, NJ 07059

D551440

© 2015 GSK group of companies or its licensee. All rights reserved.

TAMPER-EVIDENT features for your protection. This package has a security seal on both ends of the carton printed with the GSK logo. The bottle of capsules has a foil seal under the cap imprinted with the words "sealed for your protection". Each individual capsule is sealed with a dark blue band around the center of the capsule. **DO NOT USE THIS PRODUCT IF ANY OF THESE TAMPER-EVIDENT FEATURES ARE MISSING, TORN OR BROKEN.**

