


FOR FDA SUBMISSION ONLY



FOR FDA SUBMISSION ONLY



FOR FDA SUBMISSION ONLY



FOR FDA SUBMISSION ONLY

Children's Allegra Hives
Ages 6 Years and Older

Children's Allegra Hives
Ages 6 Years and Older

The makers of Allegra® do not make store brand products.
The trade dress of this Allegra® package is subject to trademark protection.

12 HOUR

Children's Allegra Hives
Ages 6 Years and Older

LOT
EXP

**ORIGINAL PRESCRIPTION STRENGTH
NON-DROWSY**

Children's Allegra Hives

Ages 6 Years and Older

RELIEF OF ITCHING DUE TO HIVES

Dissolves without water

12 Hour

12 Orally Disintegrating Tablets 30 mg each

Orange Cream Flavored

Drug Facts (continued)

Inactive ingredients (continued)
sorbitate, mannitol, methacrylic acid copolymer, microcrystalline cellulose, povidone, sodium bicarbonate, sodium starch glycolate

Questions or comments?
call toll-free 1-800-633-1610 or www.allegra.com

Drug Facts

Active ingredient (in each tablet)
Fexofenadine HCl 30 mg..... Antihistamine

Purpose
Antihistamine

Warnings
Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms:

- trouble swallowing
- trouble breathing
- dizziness or loss of consciousness
- swelling of tongue
- swelling in or around mouth

Uses
Reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- place 1 tablet on tongue; tablet disintegrates, with or without water
- adults and children take 2 tablets every 12 hours on an empty stomach; do not take more than 4 tablets in 24 hours
- children 6 to under 12 years of age take 1 tablet every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
- children under 6 years of age do not use
- ask a doctor
- adults 65 years of age and older ask a doctor
- consumers with kidney disease ask a doctor

Other information

- each tablet contains sodium 5 mg
- phenylephrine/acetaminophen combination products: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)
- use tablet immediately after opening individual blister

5009XXXX
Chattanooga, TN 37409-0219 ©2010
Dist. By: Chatham, Inc. (part of the same-vents Group).

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

ANDREA LEONARD SEGAL
01/24/2011