

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ **do not take more than directed** ■ **the smallest effective dose should be used** ■ **do not take longer than 10 days**, unless directed by a doctor (see Warnings) ■ **drink a full glass of water with each dose**


■ if taken with food, this product may take longer to work

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12- hour period
- do not exceed 3 capsules in a 24 hour period

children under 12 years: ■ ask a doctor

Other information ■ **each capsule contains:** sodium 20 mg ■ contains color additives including FD&C Yellow No. 5 (tartrazine) ■ store at 20°-25°C (68°-77°F) avoid high humidity and excessive heat above 40°C (104°F) ■ do not use if carton is open or if Tamper-Evident seal under bottle cap imprinted with "Sealed for Your Protection" is broken or missing. ■ read all directions and warnings before use. Keep carton.



NDC 00000-000-00

See new warnings information.

Naproxen Sodium Capsules, 220 mg

Pain Reliever/Fever Reducer (NSAID)


15 Capsules*

*Each Liquid-filled capsule contains 200 mg naproxen

Drug Facts (continued)

Inactive Ingredients: FD&C Blue #1, FD&C Yellow #5, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments?
 call toll free 1-800-447-1140



Drug Facts

Active ingredient (in each capsule) Naproxen Sodium 220 mg (naproxen 200 mg) (NSAID)*Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Purposes


Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Warnings


Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed



Manufactured by: Benmar Pharmaceuticals Inc.
 4125 Perimeter Drive
 High Point, NC 27265

Lot #
 Expiration Date
 Rev. 7/2015



Drug Facts (continued)

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

■ right before or after heart surgery

Ask a doctor before use if you have ■ problems or serious side effects from taking pain relievers or fever reducers

■ stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain ■ ulcers ■ bleeding problems ■ high blood pressure ■ heart or kidney disease ■ taken a diuretic ■ reached age 60 or older

Ask a doctor or pharmacist before use if you are ■ taking any other drug containing an NSAID (prescription or nonprescription) ■ taking a blood thinning (anticoagulant) or steroid drug

■ under a doctor's care for any serious condition ■ taking any other drug


When using this product ■ take with food or milk if stomach upset occurs

■ long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding. ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days

■ stomach pain or upset gets worse or lasts ■ redness or swelling is present in the painful area ■ any new symptoms appear ■ you have difficulty swallowing ■ it feels like the capsule is stuck in your throat

■ you develop heartburn



#pp
12015

(b) (4)

Pharmalabel

(b) (4)

□

□

□

□

□

□

NDC 00000-000-00

See new warnings information.

Naproxen Sodium Capsules, 220 mg

Pain Reliever/Fever Reducer (NSAID)

15 Capsules*

*Each Liquid-filled capsule contains 200 mg naproxen

Active Ingredient Purposes
 (in each capsule)
 Naproxen Sodium 220 mg Pain reliever/
 (naproxen 200 mg) fever reducer
 (NSAID)* nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to: ■ headache
 ■ muscular aches ■ minor pain of
 arthritis ■ toothache ■ backache
 ■ the common cold ■ menstrual
 cramps ■ temporarily reduces fever

Warnings
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
 Allergy alert
 Stomach bleeding warning

Directions ■ do not take more than directed ■ the smallest effective dose should be used
 ■ do not take longer than 10 days, unless directed by a doctor (see Warnings) ■ drink a full glass of water with each dose ■ if taken with food, this product may take longer to work

adults and children 12 years and older:
 ■ take 1 capsule every 8 to 12 hours while symptoms last
 ■ for the first dose you may take 2 capsules within the first hour
 ■ do not exceed 2 capsules in any 8- to 12-hour period
 ■ do not exceed 3 capsules in a 24 hour period
 ■ ask a doctor

children under 12 years:

Manufactured by: Banner Pharmacaps Inc.
 4125 Premier Drive
 High Point, NC 27265

Lot #
 Expiration Date
 Rev # 2/05

(b) (4)

Pharmalabel

(b) (4)

FRONT

NDC 00000-000-00

See new warnings information.

Naproxen Sodium Capsules, 220 mg

Pain Reliever/Fever Reducer (NSAID)
200 Capsules*

*Each Liquid-filled capsule contains 200 mg naproxen

Drug Facts

Active ingredient (in each capsule) Naproxen Sodium 220 mg (naproxen 200 mg) (NSAID) Pain reliever/fever reducer - nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - minor pain of arthritis
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Very Important: Naproxen Sodium may cause a severe allergic reaction, especially in people who are allergic to aspirin. Symptoms of a severe allergic reaction include hives, rash, difficulty breathing, swelling of the face, lips, tongue, or throat, or dizziness. If you experience any of these symptoms, stop taking this product and seek medical help right away. If you are allergic to aspirin, you should not take this product. Naproxen Sodium may cause stomach bleeding. The chance is higher if you are taking aspirin or other drugs that increase the risk of bleeding. Do not take Naproxen Sodium if you are taking aspirin or other drugs that increase the risk of bleeding. Do not take Naproxen Sodium if you are taking a blood thinner (anticoagulant) or steroid drug (corticosteroid). Do not take Naproxen Sodium if you are taking a diuretic. Do not take Naproxen Sodium if you are taking any other drug containing an NSAID (nonsteroidal anti-inflammatory drug). Do not take Naproxen Sodium if you are taking a blood thinner (anticoagulant) or steroid drug (corticosteroid). Do not take Naproxen Sodium if you are taking a diuretic. Do not take Naproxen Sodium if you are taking any other drug containing an NSAID (nonsteroidal anti-inflammatory drug).

Do not use if you are allergic to any other pain reliever/fever reducer.

Ask a doctor before use if you have:

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- bleeding problems
- high blood pressure
- heart or kidney disease
- kidney problems
- stomach ulcers
- are taking a diuretic
- are taking a blood thinner (anticoagulant) or steroid drug (corticosteroid)
- are taking any other drug containing an NSAID (nonsteroidal anti-inflammatory drug)
- are taking a diuretic
- are taking any other drug containing an NSAID (nonsteroidal anti-inflammatory drug)

When using this product:

- take with food or milk if stomach upset occurs
- take any other drug
- long term continuous use may increase the risk of heart attack or stroke

Manufactured by: Banner Pharmaceps Inc., 725 Premier Drive, High Point, NC 27266

Lot # _____
Expiration Date _____
Rev # 2/06

BACK

Drug Facts (Continued)

Stop use and ask a doctor if:

- you feel faint, vomit blood, or have bloody or black stools
- these are signs of stomach bleeding
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat
- you develop hives

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions **do not take more than directed**

- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

Other information **each capsule contains:** sodium 20 mg

- contains color additives including FD&C Yellow No. 5 (tartrazine)
- store at 20°-25°C (68°-77°F) avoid high humidity and excessive heat above 40°C (104°F)
- do not use if tamper-evident seal under bottle cap is broken or missing
- read all directions and warnings before use.

Inactive ingredients FD&C Blue #1, FD&C Yellow #5, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? Call toll free 1-800-441-1140

(b) (4)

(b) (4) **Pharmalabel**

■

□

□

□

□

■

□

□

□

□

□

□

□

□

□

□

□

□

□

□

**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Andrea Segal
2/17/2006 02:39:32 PM