

Do not use if printed safety seal under cap is torn or missing.

Manufactured by:
Barnier Pharmaceuticals, Inc.
4126 Premier Drive
High Point, NC 27265

FPO 90% UPC
TRUNCATED
000000100000

10 FREE

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

Strength to last 12 hours

30 LIQUID GELS
(LIQUID FILLED CAPSULES)

LOT & EXP

LOT _____

EXP _____

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

Drug Facts

Active Ingredient (in each capsule)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) Pain reliever/fever reducer

Nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if:

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a history of stomach bleeding
- you are taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - belching
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - you have difficulty swallowing
 - it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important.

NO COATING

MODIFIED DFL

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>7.0 pt	7.1 pt
Drug Facts (continued):	N/A	7 pt	7 pt
Headings:	N/A	7 pt	7 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

Compare to the active ingredient of ALEVE® Liquid Gels***

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

30 LIQUID GELS (LIQUID FILLED CAPSULES)

Strength to last 12 hours

Do not use if printed safety seal under cap is torn or missing. **KEEP OUT OF REACH OF CHILDREN. WARNING: ALLERGY ALERT: Naproxen sodium may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing) or shock. Stomach bleeding (especially risk-inflammatory drug (NSAID)) which may cause severe stomach bleeding or other • have had stomach ulcers or bleeding problems • use blood thinning (anticoagulant) drugs • use other drugs containing prescription or naproxen, or others) • have 3 or more alcoholic drinks every day while using this medicine • have had an allergic reaction to any other pain reliever/fever reducer. • pregnant or breast-feeding, ask a health professional before use. Directions: Adults and children 12 years of age and older: Take with water every 8 to 12 hours while symptoms last. For the first dose you may take 2 capsules within the first hour. Do not exceed 3 capsules in a 24-hour period. Children under 12 years: ask a doctor. **Call your doctor at 1-888-858-2722 or 1-888-858-2725** if you have any questions. ***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels. Manufactured by Banner Pharmaceuticals, Inc. 10000 Banner Blvd., Suite 1000, Charlotte, NC 27225**

LOT/EXP

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	4.5 pt
Subheadings:	N/A	6 pt	4.5 pt
Text:	N/A	6 pt	4.5 pt
Bullets:	N/A	5 pt (no larger)	4.5 pt
Leading:	N/A	+0.5 pt	4.9 pt
Characters Per Inch:	N/A	No more than 39	45

NO COATING

Compare to the active ingredient
of ALEVE® Liquid Gels***

**Naproxen Sodium
Capsules, 220 mg (NSAID)**
Pain Reliever/Fever Reducer

20 + 10 = 30 LIQUID GELS
(LIQUID FILLED CAPSULES)

Strength to last 12 hours

NO COATING

--- NO COATING ---

Do not use if printed safety seal under cap is torn or missing.

***This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Barringer Pharmaceuticals, Inc.
4125 Premier Drive
High Point, NC 27265

FPO, 90% UPC
TRUNCATED
000000000000

Drug Facts (continued)

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer.

Ask a doctor before use if:

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver disease, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are:

- under a doctor's care for any serious condition
- taking any other drug

When using this product:

- take with food or milk
- if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important to

Drug Facts (continued)

Inactive ingredients: FD&C Blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitol, sorbitol

Questions or comments? 1-336-812-8700 (Mon-Fri 9AM - 5PM EST)

LOT

EXP

Drug Facts (continued)

not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period


ask a doctor

Other information

- each capsule contains sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use.
- Keep carton.

MODIFIED DFL

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>7.0 pt	7.1 pt
Drug Facts	N/A	7 pt	7 pt
(continued):	N/A	7 pt	7 pt
Headings:	N/A	7 pt	7 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39



MODIFIED DFL

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>7.0 pt	7.1 pt
Drug Facts	N/A	7 pt	7 pt
(continued):	N/A	7 pt	7 pt
Headings:	N/A	7 pt	7 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

NO COATING

Compare to the active ingredient of ALEVE® Liquid Gels***

10 FREE

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

Strength to last 12 hours

50 LIQUID GELS
(LIQUID FILLED CAPSULES)

NO COATING

LIQUID GELS WITH THIS PURCHASE

--- NO-COATING ---

Do not use if printed safety seal under cap is torn or missing.

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)Pain reliever/fever reducer

Uses
*nonsteroidal anti-inflammatory drug

Temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- backache
- headache
- the common cold
- temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.
 Symptoms may include:

- hives
- facial swelling
- shock
- skin redness
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use
 if you have ever had an allergic reaction to any other pain reliever/fever reducer
 right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as

Drug Facts (continued)

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
 you experience any of the following signs of stomach bleeding:

- heartburn
- vomit blood
- have bloody or black stools
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

Drug Facts (continued)

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C Blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-866-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
 Banner Pharmaceuticals, Inc.
 4725 Premier Drive
 High Point, NC 27265

LOT

EXP

LOT & EXP

F.P.O. 100% UPC

0 100000 100000 0

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

Compare to the active ingredient of ALEVE® Liquid Gels***

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

50 LIQUID GELS
(LIQUID FILLED CAPSULES)

Strength to last 12 hours

Do not use if printed safety seal under cap is torn or missing. KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION. Warnings Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include rash, hives, itching, dizziness, trouble breathing, facial swelling, asthma (wheezing), or difficulty swallowing. If these symptoms occur, stop taking this product and seek medical help right away. **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription (aspirin, ibuprofen, naproxen, diclofenac, celecoxib, valdecoxib, or rofecoxib) or over-the-counter (aspirin, ibuprofen, naproxen, or celecoxib) pain relievers or for a longer time than directed. **Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. Ask a doctor or pharmacist before use if you are taking any other drug. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:** heart burn, vomit blood, have bloody or black stools, have stomach pain, and gain weight or lose weight. **Keep out of reach of children or children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** Adults and children 12 years and older: take 1 capsule with water every 8 to 12 hours while symptoms last. For the first dose you may take 2 capsules within the first hour. Do not exceed 2 capsules in a 24-hour period. Children under 12 years: ask a doctor. **Questions or comments?** 1-800-912-8100 (M-F 9 AM - 5 PM EST)

***The product is not manufactured or distributed by Bayer Healthcare LLC, the manufacturer of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4125 Premier Drive
High Point, NC 27265

LOT/EXP

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	5 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	43

NO COATING

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer **40 + 10 = 50 LIQUID GELS**
(LIQUID FILLED CAPSULES)
Strength to last 12 hours

NO COATING

--- NO-COATING ---

Do not use if printed safety seal under cap is torn or missing.

Pain Reliever/Fever Reducer (NSAID)
Capsules, 220 mg

Drug Facts (continued)

hearing loss

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - heartburn
 - vomit blood
 - have bloody or black stools
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

LOT

EXP

LOT & EXP

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID) Pain reliever/fever reducer

Purposes
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - backache
 - headache
 - the common cold
 - temporarily reduces fever
- temporarily relieves minor aches and pains due to:
 - muscular aches
 - menstrual cramps
 - toothache

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- in the stomach bleeding warning applies to you
- you have a history of stomach problems, such as

Drug Facts (continued)

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C Blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-866-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Banner Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

Drug Facts (continued)

heartburn

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - heartburn
 - vomit blood
 - have bloody or black stools
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

F.P.O. 100% UPC

0 100000 100000 0

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

NO COATING

**BONUS SIZE
40 + 10 FREE**

Compare to the active ingredient
of ALEVE® Liquid Gels***

**Naproxen
Sodium
Capsules,
220 mg (NSAID)**

Pain Reliever/Fever Reducer

Strength to last 12 hours

50 ~~LIQUID FILLED CAPSULES~~
(LIQUID FILLED CAPSULES)

NO COATING

Do not use if printed safety seal under cap is torn or missing.

**Naproxen Sodium
Capsules, 220 mg
(NSAID)**

Pain Reliever/Fever Reducer

Drug Facts

Active Ingredient
(in each capsule)
Naproxen sodium 220 mg
(naproxen 200 mg) (NSAID) *Pain reliever/
fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- skin redness
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

Do not use

- take more or for a longer time than directed
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as

Drug Facts (continued)

heartburn

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - heart burn
 - vomit blood
 - have bloody or black stools
- stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

Drug Facts (continued)

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

Drug Facts (continued)

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

Drug Facts (continued)

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

Drug Facts (continued)

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

Drug Facts (continued)

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

Drug Facts (continued)

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

Drug Facts (continued)

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

Drug Facts (continued)

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

Drug Facts (continued)

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

Drug Facts (continued)

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period
- ask a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F); avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use. Keep carton.


Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-812-8700 (Mon-Fri 9AM-5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels.

Manufactured by:
Bayer Pharmaceuticals, Inc.
4725 Premier Drive
High Point, NC 27265

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts (continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39



DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	8.1 pt
Drug Facts	N/A	8 pt	8 pt
(continued):	N/A	8 pt	8 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

**Compare to the active ingredient
of ALEVE® Liquid Gels*****

**Naproxen Sodium
Capsules, 220 mg (NSAID)**
Pain Reliever/Fever Reducer

EASY OPEN ARTHRITIS CAP
Strength to last 12 hours

100 LIQUID GELS
(LIQUID FILLED CAPSULES)

Do not use if printed safety seal under cap is torn or missing. KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION Warnings: Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, or blisters. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ● are age 60 or older ● have had stomach ulcers or bleeding problems ● take a blood thinner (anticoagulant) or steroid drug ● take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● take more or for a longer time than directed. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. Ask a doctor before use if the stomach bleeding warning applies to you, you have a history of stomach problems (such as heartburn), you have high blood pressure, heart disease, liver cirrhosis, or kidney disease, you are taking a diuretic, you have problems or serious side effects from taking pain relievers or fever reducers, or you have asthma. Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition or taking any other drug. When using this product take with food or milk if stomach upset occurs. The risk of heart attack or stroke may increase if you use more than directed or for longer than directed. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear, you have difficulty swallowing, or if feels like the capsule is stuck in your throat. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Directions: Do not take more than directed. The smallest effective dose should be used. Drink a full glass of water with each dose. If taken with food, this product may take longer to work. Adults and children 12 years and older: take 1 capsule every 8 to 12 hours while symptoms last. For the first dose you may take 2 capsules within the first hour. Do not exceed 2 capsules in any 8- to 12-hour period. Do not exceed 3 capsules in 24-hour period. Children under 12 years: ask a doctor. Store at 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F). Questions or comments? 1-336-812-8700 (Mon-Fri 9AM-5PM EST) This product may be manufactured or distributed by Bayer Healthcare LLC, a subsidiary of Bayer AG, at the Bayer facility in Deerfield, IL. ALEVE Liquid Gels. Manufactured by: Bayer Pharmaceuticals, Inc. 4125 Premier Drive High Point, NC 27265 L01EXP

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts (continued):	N/A	8 pt	-
Headings:	N/A	8 pt	6 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

Factory Applied Aleve 200ct Caplets \$2.00 IRC Instruction Sheet



**PLACE
IRC
HERE**



**Place IRC in upper right corner on the front of the Aleve 200ct. Caplets carton.
Wrap clear adhesive on right side of IRC around the right side of the carton. Do not cover any text.**

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER

04/01/2013