

NOT OFFICIAL SIZE/DIELINE



Label Size : 4.625" x 1"
 Track : 14/09/2016, 23/02/2017-II

Colors

(b) (4)

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts (continued):	N/A	8 pt	-
Headings:	N/A	8 pt	4.5 pt
Subheadings:	N/A	6 pt	4.5 pt
Text:	N/A	6 pt	4.5 pt
Bullets:	N/A	5 pt (no larger)	4.5 pt
Leading:	N/A	+0.5 pt	4.9 pt
Characters Per Inch:	N/A	No more than 39	45

NOT OFFICIAL SIZE/DIELINE



Carton Size : 1.65" x 1.6" x 3"
 Track : 14/09/2016, 08/02/2017,
 23/02/2017-II

MODIFIED DFL

DRUG FACTS INFO.		
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED
Drug Facts Title:	N/A	>7.0 pt
Drug Facts (continued):	N/A	7 pt
Headings:	N/A	7 pt
Subheadings:	N/A	6 pt
Text:	N/A	6 pt
Bullets:	N/A	5 pt (no larger)
Leading:	N/A	+0.5 pt
Characters Per Inch:	N/A	No more than 39
		ACTUAL SIZE IN FILE



NOT OFFICIAL SIZE/DIELINE




Carton Size : 1.875" x 1.812" x 4.18"
Track : 14/09/2016, 08/02/2017,
23/02/2017-11

Colors



CATEGORY	DRUG FACTS INFO.	
	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED
Drug Facts Title:	N/A	>8.0 pt
Drug Facts (continued):	N/A	8 pt
Headings:	N/A	8 pt
Subheadings:	N/A	6 pt
Text:	N/A	6 pt
Bullets:	N/A	5 pt (no target)
Leading:	N/A	+0.5 pt
Characters Per Inch:	N/A	No more than 39
		ACTUAL SIZE IN FILE

NOT OFFICIAL SIZE/DIELINE



Label Size : 5.25" x 1.875"
Track : 14/09/2016, 23/02/2017-II

Colors



DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	5 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	43

NOT OFFICIAL SIZE/DIELINE

Carton Size : 1.875" x 1.812" x 4.18"
 Track : 15/09/2016, 08/02/2017,
 23/02/2017-11
 Colors



CATEGORY	DRUG FACTS INFO.	
	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED
Drug Facts Title:	N/A	>8.0 pt
Drug Facts (continued):	N/A	8 pt
Headings:	N/A	8 pt
Subheadings:	N/A	6 pt
Text:	N/A	6 pt
Bullets:	N/A	5 pt (no target)
Leading:	N/A	+0.5 pt
Characters Per Inch:	N/A	No more than 39
		ACTUAL SIZE IN FILE
		10 pt
		9 pt
		9 pt
		8 pt
		8 pt
		6 pt
		6 pt
		5 pt
		6.5 pt
		39

NOT OFFICIAL SIZE/DIELINE

NDC 69452-179-16 Compare to the active ingredient
of ALEVE® Liquid Gels***

**Naproxen Sodium
Capsules, 220 mg (NSAID)**
Pain Reliever/Fever Reducer

50 CAPSULES

Strength to last 12 hours

Do not use if printed safety seal under cap is torn or missing.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Warning: Allergic reaction to naproxen sodium may cause severe allergic reaction, especially when taking aspirin or other NSAIDs. Discontinue use immediately if you experience hives, rash, or difficulty breathing. In patients taking aspirin or other NSAIDs, serious skin reactions like Stevens-Johnson syndrome and toxic epidermal necrolysis may occur without warning. This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinner (anticoagulant) or steroid drug • take other drugs containing naproxen or naproxol sodium (ibuprofen, naproxen, naproxol, or ibuprofen) • take aspirin, aspirin-like drugs, or other NSAIDs • use alcohol, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risks are higher if you use more than directed or for longer than indicated. Do not use if you have ever had a legic reaction to any other pain reliever/fever reducer. Ask a doctor or pharmacist before use if you are taking any other drug. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: (stomach pain, vomit blood, have bloody or black stools, have stomach pain that does not go away), or (pain or fever in the throat, nose, or mouth, or pain in the chest, stomach, or back). **Mean and all death of children. In case of overdose, get medical help or contact a Poison Control Center right away. If medicine kills and children 12 years and older take 1 capsule with water every 6 to 12 hours while symptoms last. For their first dose you may take 2 capsules with the first hour. Do not exceed 3 capsules in a 24-hour period. Do not use under 12 years, as a child.**

Questions are answered? 1-800-234-2466 (Mon-Fri 9AM - 5PM EST)

*This product is not manufactured or distributed by Bayer Pharmaceuticals Corporation or Aleve® Liquid Gels. Manufactured for Bayer Pharmaceuticals Inc., Princeton, NJ 08540. Lot/EXP

02/17

Label Size : 5.25" x 1.875"
 Track : 14/09/2016, 23/02/2017-II

Colors



DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	5 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	43

NOT OFFICIAL SIZE/DIELINE



Carton Size : 1.875" x 1.812" x 4.18"
Track : 15/09/2016, 08/02/2017,
23/02/2017-11

Colors



CATEGORY	DRUG FACTS INFO.		
	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	10 pt
Drug Facts (continued):	N/A	8 pt	9 pt
Headings:	N/A	8 pt	9 pt
Subheadings:	N/A	6 pt	8 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (or larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

NOT OFFICIAL SIZE/DIELINE

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

NDC 69452-179-78 Compare to the active ingredient
of ALEVE® Liquid Gels***

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

EASY OPEN ARTHRITIS CAP

80 CAPSULES

Strength to last 12 hours

Do not use if printed safety seal under cap is torn or missing.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Warnings Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, or blisters. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed. Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. • Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. Ask a doctor before use if the stomach bleeding warning applies to you, you, have a history of stomach problems (such as heartburn), you have high blood pressure, heart disease, liver or kidney disease, kidney disease, asthma, or had a stroke. You are taking a diuretic, you have problems or serious side effects from taking pain relievers or fever reducers. Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition or taking any other drug. When using this product take with food or milk if stomach upset occurs. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding (fever, faint, vomit blood, have bloody or black stools, have stomach pain that does not get better), you have symptoms of heart problems or stroke (chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling), pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present in the painful area, any new symptoms appear, you have difficulty swallowing, or if these like the capsule is stuck in your throat. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. **Directions** Do not take more than directed. The smallest effective dose should be used. Drink a full glass of water with each dose. If taken with food, this product may be able to get to work. Adults and children 12 years and older: Take 1 capsule every 8 to 12 hours while symptoms last. For the first 2 days you may take 2 capsules within the first hour. Do not exceed 2 capsules in any 8- to 12-hour period. Do not exceed 3 capsules in a 24-hour period. Children under 12 years: Ask a doctor before use. Store at 20°-25° C (68°-77° F) and protect from moisture. See USP Controlled Room Temperature. Excipients: See USP Controlled Room Temperature. **Questions or comments?** 1-888-265-2466 (Mon-Fri 9AM - 5PM EST)

***This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of ALEVE® Liquid Gels. Manufactured for: Biopharma Inc., 600 Alexander Road, Princeton, NJ 08540

03/17
LOT/EXP

Label Size : 7.5" x 2.5"
Track : 14/09/2016, 12/02/2017,
23/02/2017-II, 01/03/2017

Colors



DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	6 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

NOT OFFICIAL SIZE/DIELINE

NO COATING

NO COATING

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

NO COATING

NO COATING

EASY OPEN ARTHRITIS CAP
Strength to last 12 hours

80 CAPSULES

NDC 69452-179-78

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Compare to the active ingredient of ALEVE® Liquid Gels***

EASY OPEN ARTHRITIS CAP
Strength to last 12 hours

80 CAPSULES

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*.....Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Purposes
Pain reliever/fever reducer

Uses
temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- backache
- headache
- the common cold
- temporarily reduces fever
- muscular aches
- menstrual cramps
- toothache

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- shock
- facial swelling
- skin reddening
- asthma (wheezing)
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
 • if you have ever had an allergic reaction to any other pain reliever/fever reducer
 • right before or after heart surgery

Ask a doctor before use if
 • stomach bleeding warning applies to you
 • you have a history of stomach problems, such as heartburn
 • you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 • you are taking a diuretic
 • you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are
 • under a doctor's care for any serious condition
 • taking any other drug

When using this product
 • take with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if
 • you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 • do not take more than directed
 • the smallest effective dose should be used
 • drink a full glass of water with each dose
 • if taken with food, this product may take longer to work

adults and children 12 years and older:	<ul style="list-style-type: none"> • take 1 capsule every 8 to 12 hours while symptoms last • for the first dose you may take 2 capsules within the first hour • do not exceed 2 capsules in any 8- to 12-hour period • do not exceed 3 capsules in a 24-hour period
children under 12 years:	<ul style="list-style-type: none"> • ask a doctor

Other information
 • each capsule contains: sodium 20 mg
 • store at 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F)
 • read all directions and warnings before use. Keep carton.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon-Fri 9AM - 5PM EST)

***This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of ALEVE® Liquid Gels.

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

EASY OPEN ARTHRITIS CAP

80 CAPSULES

Strength to last 12 hours

Manufactured for:
BioPharma Inc.
600 Alexander Road,
Princeton, NJ 08540



3 69452 17978 9

Do not use if printed safety seal under cap is torn or missing.

Rev # 02/17
Lot No.:
Exp. Date:

LOT & EXP

Colors

(b) (4)

Carton Size : 2.625" x 2.625" x 4.875"
Track : 14/09/2016, 08/02/2017, 23/02/2017-II

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	10 pt
Drug Facts (continued):	N/A	8 pt	9 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

NOT OFFICIAL SIZE/DIELINE



Carton Size : 1.875" x 1.812" x 4.18"
Track : 15/09/2016, 08/02/2017,
23/02/2017-11

Colors

(b) (4)


CATEGORY	DRUG FACTS INFO.		ACTUAL SIZE IN FILE
	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	
Drug Facts Title:	N/A	>8.0 pt	10 pt
Drug Facts (continued):	N/A	8 pt	9 pt
Headings:	N/A	8 pt	9 pt
Subheadings:	N/A	6 pt	8 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (or larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

NOT OFFICIAL SIZE/DIELINE



Carton Size : 1.875" x 1.812" x 4.18"
 Track : 15/09/2016, 08/02/2017,
 23/02/2017-11

Colors



CATEGORY	DRUG FACTS INFO.		
	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	10 pt
Drug Facts (continued):	N/A	8 pt	9 pt
Headings:	N/A	8 pt	9 pt
Subheadings:	N/A	6 pt	8 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (or larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

NOT OFFICIAL SIZE/DIELINE

NDC 69452-170-79 Compare to the active ingredient
of ALEVE® Liquid Gels***

**Naproxen Sodium
Capsules, 220 mg (NSAID)**
Pain Reliever/Fever Reducer

160 CAPSULES

Strength to last 12 hours

Do not use if printed safety seal under cap is torn or missing.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Warning: Allergic reaction to naproxen sodium may cause severe allergic reaction, especially if you have had allergic reactions to aspirin, salicylates, or other NSAIDs. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinner (anticoagulant) or steroid drug • take other drugs containing naproxen or naproxol sodium (ibuprofen, naproxen, naproxen sodium) • have a history of ulcers or stomach ulcers • are taking aspirin • are taking other NSAIDs • are taking aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risks are higher if you use more than directed or for longer than indicated. Do not use if you have ever had a life-threatening allergic reaction to any other pain reliever/fever reducer. Ask a doctor or pharmacist before use if you are taking any other drug. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: (stomach pain, vomit blood, have bloody or black stools, have stomach pain that does not go away), or (pain or fever signs of infection, such as sore throat, chest pain, cough, or fever). **Mean and all death of children. In case of overdose, get medical help or contact a Poison Control Center right away. *Directions:* Adults and children 12 years and older: take 1 capsule with water every 6 to 12 hours while symptoms last. For other first doses you may take 2 capsules with the first hour. Do not exceed 3 capsules in a 24-hour period. Children under 12 years, ask a doctor. *Questions are answered:* 1-800-235-2466 (Mon-Fri 9AM - 5PM EST)**

***This product is not manufactured or distributed by Bayer Pharmaceuticals Corporation or Aleve® Liquid Gels. Manufactured for: Bayer Pharmaceuticals Inc., 600 Aleve Center Road, Princeton, NJ 08540
Lot/EXP

02/17

Label Size : 5.25" x 1.875"
Track : 14/09/2016, 23/02/2017-II


Colors



(b) (4)

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	-
Drug Facts	N/A	8 pt	-
(continued):	N/A	8 pt	-
Headings:	N/A	8 pt	5 pt
Subheadings:	N/A	6 pt	5 pt
Text:	N/A	6 pt	5 pt
Bullets:	N/A	5 pt (no larger)	4 pt
Leading:	N/A	+0.5 pt	5.4 pt
Characters Per Inch:	N/A	No more than 39	43

NOT OFFICIAL SIZE/DIELINE



Carton Size : 1.875" x 1.812" x 4.18"
 Track : 15/09/2016, 08/02/2017,
 23/02/2017-11

Colors

CATEGORY	DRUG FACTS INFO.		
	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	10 pt
Drug Facts (continued):	N/A	8 pt	9 pt
Headings:	N/A	8 pt	9 pt
Subheadings:	N/A	6 pt	8 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no target)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	39

NO COATING

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

160 CAPSULES

Compare to the active ingredient of ALEVE® Liquid Gels ***

NDC 69452-179-79

Strength to last 12 hours

Drug Facts

Active ingredient (in each capsule) Purposes
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
 - temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

NO COATING

Naproxen Sodium Capsules, 220 mg (NSAID)
Pain Reliever/Fever Reducer

Do not use if printed label under cap is torn or missing.

3 69452 17979 6


Rev # 02/17
 Lot No.: **LOT & EXP**
 Exp. Date:

NOT OFFICIAL SIZE/DIELINE

Do not use if printed safety seal under cap is torn or missing.
***This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of ALEVE® Liquid Gels.

Manufactured for:
Bionpharma Inc.
800 Alexander Road,
Princeton, NJ 08540

02/17



3 69452 17979 6

LOT/EXP


NDC 69452-179-79 Compare to the active ingredient of ALEVE® Liquid Gels***

Naproxen Sodium Capsules, 220 mg (NSAID)

Pain Reliever/Fever Reducer

160 CAPSULES

Strength to last 12 hours

PEEL HERE 

Drug Facts

Active ingredient (in each capsule)
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
 - menstrual cramps
 - headache
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

TOP



BEHIND TOP

Drug Facts (continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - slurred speech
 - weakness in one part or side of body
 - leg swelling

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- you have difficulty swallowing
- it feels like the capsule is stuck in your throat

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- if taken with food, this product may take longer to work

adults and children 12 years and older:

- take 1 capsule every 8 to 12 hours while symptoms last
- for the first dose you may take 2 capsules within the first hour
- do not exceed 2 capsules in any 8- to 12-hour period
- do not exceed 3 capsules in a 24-hour period

children under 12 years:

- ask a doctor

Drug Facts (continued)

Other information

- each capsule contains: sodium 20 mg
- store at 20-25°C (68-77°F) avoid high humidity and excessive heat above 40°C (104°F)
- read all directions and warnings before use.

Inactive ingredients FD&C blue #1, gelatin, glycerin, lactic acid, mannitol, pharmaceutical ink, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

Questions or comments? 1-888-235-2466 (Mon-Fri 9AM – 5PM EST)

BASE

Label Size : 8.625" x 3.25"
Track : 15/09/2016, 08/02/2017,
23/02/2017-II

Colors

DRUG FACTS INFO.			
CATEGORY	PRIMARY PACKAGING REQUIRED	SECONDARY PACKAGING REQUIRED	ACTUAL SIZE IN FILE
Drug Facts Title:	N/A	>8.0 pt	10 pt
Drug Facts (continued):	N/A	8 pt	9 pt
Headings:	N/A	8 pt	8 pt
Subheadings:	N/A	6 pt	6 pt
Text:	N/A	6 pt	6 pt
Bullets:	N/A	5 pt (no larger)	5 pt
Leading:	N/A	+0.5 pt	6.5 pt
Characters Per Inch:	N/A	No more than 39	36

Reference ID: 4070416