

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
 Helvetica Condensed
 Variable horizontal scale adheres to 36 characters per inch

Drug Facts - 9 pt Helvetica Cnd Bold Italic
Drug Facts (continued)
 Drug Facts - 8 pt Helvetica Cnd Bold Italic (continued) - 8 pt Helvetica Cnd

Headings - 6 pt Helvetica Cnd Bold Italic
Sub Heads - 6 pt Helvetica Cnd Bold

Text - 6 pt Helvetica Cnd minimum leading 6.5
 Square Bullets - 3.5 pt Zaph Dingbats no compression
 Border - 1 pt rule 3/64" gap all around
 Barline - 1 pt rule
 Hairline - .5 pt rule

Certified by: _____

STANDARD FORMAT
 HIGHLIGHTS CONTAINED
 SOURCE: PHOTOCOPIED SCALE: 100% (100% OF ORIGINAL)
 FEEL FREE
 Drug Facts: 5 pt Helvetica Cond, Bold Italic
 Drug Facts: (continued)
 Drug Facts: 5 pt Helvetica Cond, Bold Italic
 (continued) - 5 pt Helvetica Cond
 Headings: 6 pt Helvetica Cond, Bold Italic
 Sub-Headings: 6 pt Helvetica Cond, Bold
 Text: 6 pt Helvetica Cond, minimum leading 6 s
 Source Effects: 5 pt Zaps, Emphasis on compression
 Border: 1 pt line, 2 pt gap all around
 Bottom: 1 pt line
 Warning: 5 pt line
 Certified by: [signature]

ATTACH LEGEND ON ALL
 DIGITAL MECHANICALS

30 ORALLY DISINTEGRATING TABLETS

Claritin RediTabs

loratadine 5 mg

TABLETS **12 Hour**

Claritin RediTabs
loratadine 5 mg

New 12 Hour!

Non-Drowsy*

Claritin

RediTabs

loratadine 5 mg / antihistamine

Allergy

Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

12 Hour

No Water Needed
Melts in Your Mouth

Actual Size

30 ORALLY DISINTEGRATING TABLETS

4 1100 80521 8

NO VARNISH
CODE AREA

NO VARNISH

NO VARNISH

NO VARNISH

30232500

30232500

VARNISH

Drug Facts (continued)

Indications and Usage: Claritin RediTabs is used to relieve the symptoms of seasonal allergic rhinitis (hay fever) and perennial allergic rhinitis.

Warnings: Do not use if you are allergic to any of the ingredients listed below. Do not use if you are taking other allergy medicine. Do not use if you are pregnant or breastfeeding. Do not use if you are taking other medicine for allergies. Do not use if you are taking other medicine for asthma. Do not use if you are taking other medicine for high blood pressure. Do not use if you are taking other medicine for heart disease. Do not use if you are taking other medicine for diabetes. Do not use if you are taking other medicine for thyroid disease. Do not use if you are taking other medicine for kidney disease. Do not use if you are taking other medicine for liver disease. Do not use if you are taking other medicine for stomach or intestinal disease. Do not use if you are taking other medicine for urinary tract infection. Do not use if you are taking other medicine for prostate disease. Do not use if you are taking other medicine for blood thinners. Do not use if you are taking other medicine for seizure medicine. Do not use if you are taking other medicine for antidepressants. Do not use if you are taking other medicine for anti-anxiety medicine. Do not use if you are taking other medicine for pain relievers. Do not use if you are taking other medicine for muscle relaxants. Do not use if you are taking other medicine for sedatives. Do not use if you are taking other medicine for alcohol. Do not use if you are taking other medicine for grapefruit juice. Do not use if you are taking other medicine for St. John's Wort. Do not use if you are taking other medicine for warfarin. Do not use if you are taking other medicine for digoxin. Do not use if you are taking other medicine for lithium. Do not use if you are taking other medicine for theophylline. Do not use if you are taking other medicine for tacrolimus. Do not use if you are taking other medicine for cyclosporin. Do not use if you are taking other medicine for sirolimus. Do not use if you are taking other medicine for everolimus. Do not use if you are taking other medicine for vardenafil. Do not use if you are taking other medicine for sildenafil. Do not use if you are taking other medicine for tadalafil. Do not use if you are taking other medicine forildenafil. Do not use if you are taking other medicine for avanafil. Do not use if you are taking other medicine for dapoxetine. Do not use if you are taking other medicine for tramadol. Do not use if you are taking other medicine for bupropion. Do not use if you are taking other medicine for mirtazapine. Do not use if you are taking other medicine for escitalopram. Do not use if you are taking other medicine for sertraline. Do not use if you are taking other medicine for fluoxetine. Do not use if you are taking other medicine for paroxetine. Do not use if you are taking other medicine for citalopram. Do not use if you are taking other medicine for desvenlafaxine. Do not use if you are taking other medicine for venlafaxine. Do not use if you are taking other medicine for duloxetine. Do not use if you are taking other medicine for milnacipran. Do not use if you are taking other medicine for levamisole. Do not use if you are taking other medicine for mebendazole. Do not use if you are taking other medicine for ivermectin. Do not use if you are taking other medicine for praziquantel. Do not use if you are taking other medicine for albendazole. Do not use if you are taking other medicine for fenbendazole. Do not use if you are taking other medicine for oxfendazole. Do not use if you are taking other medicine for moxidectin. Do not use if you are taking other medicine for selamectin. Do not use if you are taking other medicine for moxidectin. Do not use if you are taking other medicine for imidacloprid. Do not use if you are taking other medicine for deltamethrin. Do not use if you are taking other medicine for permethrin. Do not use if you are taking other medicine for cyfluthrin. Do not use if you are taking other medicine for fenprothion. Do not use if you are taking other medicine for fenitrothion. Do not use if you are taking other medicine for chlorpyrifos. Do not use if you are taking other medicine for diazinon. Do not use if you are taking other medicine for malathion. Do not use if you are taking other medicine for methidathion. Do not use if you are taking other medicine for propoxur. Do not use if you are taking other medicine for spinetor. Do not use if you are taking other medicine for acetaminophen. Do not use if you are taking other medicine for aspirin. Do not use if you are taking other medicine for ibuprofen. Do not use if you are taking other medicine for naproxen. Do not use if you are taking other medicine for celecoxib. Do not use if you are taking other medicine for rofecoxib. Do not use if you are taking other medicine for valdecoxib. Do not use if you are taking other medicine for etoricoxib. Do not use if you are taking other medicine for meloxicam. Do not use if you are taking other medicine for piroxicam. Do not use if you are taking other medicine for tenoxicam. Do not use if you are taking other medicine for flurbiprofen. Do not use if you are taking other medicine for aceclofenac. Do not use if you are taking other medicine for paracetamol. Do not use if you are taking other medicine for acetaminophen. Do not use if you are taking other medicine for aspirin. Do not use if you are taking other medicine for ibuprofen. Do not use if you are taking other medicine for naproxen. Do not use if you are taking other medicine for celecoxib. Do not use if you are taking other medicine for rofecoxib. Do not use if you are taking other medicine for valdecoxib. Do not use if you are taking other medicine for etoricoxib. Do not use if you are taking other medicine for meloxicam. Do not use if you are taking other medicine for piroxicam. Do not use if you are taking other medicine for tenoxicam. Do not use if you are taking other medicine for flurbiprofen. Do not use if you are taking other medicine for aceclofenac.



ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
 Helvetica Condensed
 Variable horizontal scale adheres to 28 characters per inch

Drug Facts - 8 pt Helvetica Cond. Bold Italic
Drug Facts - continued:
Drug Facts - 8 pt Helvetica Cond. Bold Italic
 continued - 8 pt Helvetica Cond

Headings - 8 pt Helvetica Cond. Bold Italic
Sub Heads - 6 pt Helvetica Cond. Bold

Text - 8 pt Helvetica Cond. minimum leading 6.5
 Square Endlets - 3.5 pt Zapf Dingbats no compression
 Eocent - 1 pt rule, 5.64" gap all around
 Eocline - 1 pt rule
 Hairline - 5 pt rule

Certified by:

Claritin RediTabs
loratadine 5 mg

12 Hour

Non-Drowsy* Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

Allergy

When taken as directed, Claritin RediTabs can help relieve your allergy symptoms.

5-MINUTE DISINTEGRATING TABLETS

Claritin RediTabs
loratadine 5 mg

12 Hour

No Water Needed Melts in Your Mouth

CODE AREA

5-MINUTE DISINTEGRATING TABLETS

UPC 100%

0 000000 000000

© Copyright & Distributed by Schering-Plough HealthCare Products, Inc., Kenilworth, NJ 07033 USA. All Rights Reserved. Made in United Kingdom.

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
 Helvetica Condensed
 Variable horizontal scale adheres to 38 characters per inch.

Drug Facts - 8 pt Helvetica Cond Bold Italic
Drug Facts (continued) - 8 pt Helvetica Cond Bold Italic
Drug Facts - 8 pt Helvetica Cond Bold Italic (continued) - 8 pt Helvetica Cond
Headings - 8 pt Helvetica Cond Bold Italic
Sub Heads - 6 pt Helvetica Cond Bold
Text - 6 pt Helvetica Cond minimum leading 8 E
Source Bullets - 2 E 5 pt Zaph Dingbats no compression
Border - 1 pt rule 3 E 4" gap all around
Baseline - 1 pt rule _____
Headline - 3 pt rule _____

Certified by: ym

Claritin RediTabs
loratadine 5 mg

12 Hour

5-MINUTE DISINTEGRATING TABLETS

Directions
 Take 1 tablet orally once daily with or without food. Do not take more than 1 tablet in 24 hours. Do not take more than 12 tablets in 30 days. Do not take more than 12 tablets in 30 days. Do not take more than 12 tablets in 30 days.

Warnings
 Do not take if you are allergic to loratadine or any of the ingredients. Do not take if you are taking other antihistamines. Do not take if you are taking other sedatives. Do not take if you are taking other drugs that may interact with loratadine. Do not take if you are pregnant or breastfeeding. Do not take if you are taking other drugs that may interact with loratadine.

Other Information
 Contains 12 tablets. Do not use if the seal is broken. Do not use if the tablets are discolored or broken.

Questions or Comments?
 Call 1-800-CLARITIN (1-800-252-7441) or visit www.claritin.com

10 ORALLY DISINTEGRATING TABLETS

Claritin RedItabs
 loratadine 5 mg

12 Hour

Age 6 years and older

Claritin RedItabs
 loratadine 5 mg / antihistamine

Allergy

10 ORALLY DISINTEGRATING TABLETS

UPC 100%
 0 00000 000000 0

12 Hour

Non-Drowsy[®]
 Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

No Water Needed
 Melts in Your Mouth

CODE AREA


10 ORALLY DISINTEGRATING TABLETS

© Copyright © and Distributed by Johnson & Johnson HealthCare Products, Inc., Kenilworth, NJ 07033 USA. All Rights Reserved. Made in United Kingdom.

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

- STANDARD FORMAT**
- Helvetica Condensed
- Variable horizontal scale adheres to 35 characters per inch
- Drug Facts** - 8 pt Helvetica Cond Bold Italic
- Drug Facts** - continued
- Drug Facts** - 8 pt Helvetica Cond Bold Italic
- recontinued - 8 pt Helvetica Cond
- Headings** - 8 pt Helvetica Cond Bold Italic
- Sub Heads** - 8 pt Helvetica Cond Bold
- Text** - 8 pt Helvetica Cond minimum leading 8/3
- Square Bullets** - 3.5 pt Zaph Clingbats no compression
- Border** - 1 pt rule, 3/16" gap all around
- Baseline** - 1 pt rule
- Headline** - 5 pt rule

Certified by: **vm**




DESCRIPTION: Claritin RediTabs 5mg 5ct 2.205" X 2.717" (56mm X 69mm) DIE IS AT 100%

FILE NAME: PA27284.00_K.ai		APO 0472-06		 <input type="checkbox"/> Verify UPC (if applicable) <input type="checkbox"/> Verify SP PDF <input type="checkbox"/> Verify Illustrator version
COLORS... <div style="background-color: black; width: 15px; height: 10px; display: inline-block; margin-right: 5px;"></div> Black	FONTS... 1. Helvetica 2. 3. 4. 5. 6.	OUTPUT SPECIFICATIONS... Illustrator CS	PIF: OTC050915001 Item Code: 27284-00 UPC: N/A RIC#: 30502604	
• SEPARATOR NOTE: IF ELECTRONIC ART SUPPLIED DOES NOT MATCH THIS PRINT – DO NOT PROCEED! CONTACT S/P MARKETING SERVICES AT ONCE. • THIS LASER PROOF INDICATES COLOR BREAK ONLY AND MAY NOT ACCURATELY REFLECT ACTUAL PRODUCTION COLORS. • DIE LINES SHOWN FOR REFERENCE ONLY – DO NOT PRINT.			PROOFREAD: _____ DATE: _____	

POL Claritin RediTabs 5mg 5ct			SCHERING-PLOUGH ICOM REVISION LOG
REVISION NUMBER: 1			
Item Code	Date	Artist/Vendor	Nature of Revisions
27284-00	09.06.06		PU 026140 as base; PU elements from 030138 and put into this file



copy is 5 pt.




Version 2

DESCRIPTION: Claritin RediTabs 5mg 10ct 2.205" X 4.528" (56mm X 115mm)

DIE IS AT 100%

FILE NAME: PA27276.00_K.ai		APO 0473-06	
COLORS...  Black	FONTS... 1. Helvetica 2. 3. 4. 5. 6.	OUTPUT SPECIFICATIONS... Illustrator CS	PIF: OTC050915001 Item Code: 27276-00 UPC: N/A RIC#: 30502701
• SEPARATOR NOTE: IF ELECTRONIC ART SUPPLIED DOES NOT MATCH THIS PRINT – DO NOT PROCEED! CONTACT S/P MARKETING SERVICES AT ONCE. • THIS LASER PROOF INDICATES COLOR BREAK ONLY AND MAY NOT ACCURATELY REFLECT ACTUAL PRODUCTION COLORS. • DIE LINES SHOWN FOR REFERENCE ONLY – DO NOT PRINT.			 <input type="checkbox"/> Verify UPC (if applicable) <input type="checkbox"/> Verify SP PDF <input type="checkbox"/> Verify Illustrator version PROOFREAD _____ DATE: _____

POL Claritin RediTabs 5mg 5ct			SCHERING-PLOUGH ICOM REVISION LOG
REVISION NUMBER: 1			
Item Code	Date	Artist/Vendor	Nature of Revisions
27276-00	09.06.06		PU 026165 as base; PU logo layout from client files; this becomes version 2
27276-00	09.06.06		Make 5mg Bold in all.