



NO VARNISH

00000000

NO VARNISH

00000000

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b> Loratadine 5 mg</p> <p><b>Purpose</b> Antihistamine</p> <p><b>Uses</b> temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:</p> <ul style="list-style-type: none"> <li>■ runny nose</li> <li>■ sneezing</li> <li>■ itchy, watery eyes</li> <li>■ itching of the nose or throat</li> </ul> <p><b>Warnings</b> Do not use if you have ever had an allergic reaction to this product or any of its ingredients.</p> <p>Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.</p> <p>When using this product do not take more than directed. Taking more than directed may cause drowsiness.</p> <p>Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <table style="width: 100%; border: none;"> <tr> <td style="border: none;"> <ul style="list-style-type: none"> <li>■ place 1 tablet on tongue; tablet disintegrates, with or without water</li> </ul> </td> <td style="border: none; padding-left: 10px;"> <table style="border: none;"> <tr> <td style="border: none;">adults and children 6 years and over</td> <td style="border: none;">1 tablet every 12 hours; not more than 2 tablets in 24 hours</td> </tr> <tr> <td style="border: none;">children under 6 years of age</td> <td style="border: none;">ask a doctor</td> </tr> <tr> <td style="border: none;">consumers with liver or kidney disease</td> <td style="border: none;">ask a doctor</td> </tr> </table> </td> </tr> </table>	<ul style="list-style-type: none"> <li>■ place 1 tablet on tongue; tablet disintegrates, with or without water</li> </ul>	<table style="border: none;"> <tr> <td style="border: none;">adults and children 6 years and over</td> <td style="border: none;">1 tablet every 12 hours; not more than 2 tablets in 24 hours</td> </tr> <tr> <td style="border: none;">children under 6 years of age</td> <td style="border: none;">ask a doctor</td> </tr> <tr> <td style="border: none;">consumers with liver or kidney disease</td> <td style="border: none;">ask a doctor</td> </tr> </table>	adults and children 6 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours	children under 6 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor	<p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn</li> <li>■ store between 20° to 25° C (68° to 77° F)</li> <li>■ use tablet immediately after opening individual blister</li> <li>■ complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration</li> </ul> <p><b>Inactive ingredients</b> anhydrous citric acid, gelatin, mannitol, mint flavor</p> <p><b>Questions or comments?</b> 1-800-CLARITIN (1-800-252-7484) or <a href="http://www.claritin.com">www.claritin.com</a></p> <div style="text-align: center;"> <p>• No Water Needed • Melts in your mouth</p> <p>Follow these directions carefully. Do not attempt to push the tablet through the foil.</p> <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div style="text-align: center;"> <p>1. Peel back outer edge.</p> </div> <div style="text-align: center;"> <p>2. Gently push tablet out.</p> </div> <div style="text-align: center;"> <p>3. Place the tablet on tongue and close mouth. The tablet will disintegrate.</p> </div> </div> </div>
<ul style="list-style-type: none"> <li>■ place 1 tablet on tongue; tablet disintegrates, with or without water</li> </ul>	<table style="border: none;"> <tr> <td style="border: none;">adults and children 6 years and over</td> <td style="border: none;">1 tablet every 12 hours; not more than 2 tablets in 24 hours</td> </tr> <tr> <td style="border: none;">children under 6 years of age</td> <td style="border: none;">ask a doctor</td> </tr> <tr> <td style="border: none;">consumers with liver or kidney disease</td> <td style="border: none;">ask a doctor</td> </tr> </table>	adults and children 6 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours	children under 6 years of age	ask a doctor	consumers with liver or kidney disease	ask a doctor		
adults and children 6 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours								
children under 6 years of age	ask a doctor								
consumers with liver or kidney disease	ask a doctor								

00000000

NO VARNISH

00000000

0 4110080531 7

00000-00

© Copyright & Distributed by MSD Consumer Care, Inc., a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA. All Rights Reserved. Product of Ireland.

The graphics on the front panel of this carton constitute trademarks of MSD Consumer Care, Inc.

NO VARNISH CODE AREA

NO VARNISH

NDC 11523-7202-3

NO VARNISH

NO VARNISH

(b) (4)

**Claritin® RediTabs**

**24 Hour**

10 ORALLY DISINTEGRATING TABLETS FOR 10 DAYS OF RELIEF

Original Prescription Strength

NDC 11523-7157-2

Claritin® RediTabs

Non-Drowsy\*

**Claritin®**

**RediTabs®**

loratadine 10 mg/antihistamine

**Indoor & Outdoor Allergies**

\* When taken as directed. See Drug Facts Panel.

**24 Hour**

**Relief of:**

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

**No Water Needed Melts in Your Mouth**

NO VARNISH CODE AREA

**10 ORALLY DISINTEGRATING TABLETS**

The graphics on the front panel of this carton constitute trademarks of MSD Consumer Care, Inc.

© Copyright & Distributed by MSD Consumer Care, Inc.,  
PO Box 377, Memphis, TN 38151 USA,  
a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA.  
All rights reserved. Product of Ireland.

00000-00

0 41100 80602 4

recyclable carton

NO VARNISH

**Drug Facts**

**Active ingredient (in each tablet)** Loratadine 10 mg, Antihistamine

**Purpose** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat

**Warnings** Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

**Do not use if you have liver or kidney disease.** Your doctor should determine if you need a different dose.

**Ask a doctor before use if you have liver or kidney disease.** Taking more than directed may cause drowsiness.

**When using this product** do not take more than directed. Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if an allergic reaction to this product occurs.** Seek medical help right away.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ place 1 tablet on tongue; tablet disintegrates, with or without water adults and children 6 years and over 1 tablet daily; not more than 1 tablet in children under 6 years of age ask a doctor ■ ask a doctor

**Other information** ■ safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn ■ store between 20° to 25° C (68° to 77° F) ■ use tablet immediately after opening individual blister ■ complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

**Inactive ingredients** antihydrous citric acid, gelatin, mannitol, mint flavor

**Questions or comments?** 1-800-CLARITIN (1-800-232-7184) or [www.claritin.com](http://www.claritin.com)

**Follow these directions carefully. Do not attempt to push the tablet through the foil.**

**No Water Needed • Melts in your mouth**

1. Peel back outer edge. 2. Gently push tablet out. 3. Place the tablet on tongue and close mouth. The tablet will disintegrate.

NO VARNISH

NO VARNISH

**Drug Facts**

**Active ingredient (in each tablet)** Loratadine 10 mg, Antihistamine

**Purpose** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat

**Warnings** Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

**Do not use if you have liver or kidney disease.** Your doctor should determine if you need a different dose.

**Ask a doctor before use if you have liver or kidney disease.** Taking more than directed may cause drowsiness.

**When using this product** do not take more than directed. Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if an allergic reaction to this product occurs.** Seek medical help right away.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ place 1 tablet on tongue; tablet disintegrates, with or without water adults and children 6 years and over 1 tablet daily; not more than 1 tablet in children under 6 years of age ask a doctor ■ ask a doctor

**Other information** ■ safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn ■ store between 20° to 25° C (68° to 77° F) ■ use tablet immediately after opening individual blister ■ complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

**Inactive ingredients** antihydrous citric acid, gelatin, mannitol, mint flavor

**Questions or comments?** 1-800-CLARITIN (1-800-232-7184) or [www.claritin.com](http://www.claritin.com)

**Follow these directions carefully. Do not attempt to push the tablet through the foil.**

**No Water Needed • Melts in your mouth**

1. Peel back outer edge. 2. Gently push tablet out. 3. Place the tablet on tongue and close mouth. The tablet will disintegrate.

NO VARNISH

00000000



