



**Drug Facts**

Active ingredient	Purpose
Levonorgestrel 1.5 mg	Emergency contraceptive

**Use** for women to reduce chance of pregnancy after unprotected sex (if a contraceptive failed or if you did not use birth control)

**Warnings**  
 Allergy alert: Do not use if you have ever had an allergic reaction to levonorgestrel  
 Sexually transmitted diseases (STDs) alert: This product does not protect against HIV/AIDS or other STDs  
 Do not use  
 ■ if you are already pregnant (because it will not work)  
 ■ for regular birth control  
 Ask a doctor or pharmacist before use if you are taking efavirenz (HIV medication) or rifampin (tuberculosis treatment) or medication for seizures (epilepsy). These medications may reduce the effectiveness of levonorgestrel.  
 When using this product you may have  
 ■ menstrual changes ■ tiredness ■ breast pain  
 ■ nausea ■ headache ■ vomiting  
 ■ lower stomach (abdominal) pain ■ dizziness  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control center right away.

**Directions**  
 ■ take as soon as possible within 72 hours (3 days) after unprotected sex. The sooner you take it the better it will work.  
 ■ if you vomit within 2 hours after taking the medication, call a healthcare professional to find out if you should repeat the dose

**Other information** ■ read the instructions, warnings and enclosed product leaflet before use

**Drug Facts (continued)**

**Questions?** Call 1-800-330-1271 or visit [www.PlanBOneStep.com](http://www.PlanBOneStep.com)

**Drug Facts (continued)**  
 ■ do not use if carton is open or blister seal is broken or missing  
 ■ store at 20-25°C (68-77°F)

**Inactive ingredients** colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, potato starch, talc



emergency contraceptive  
One Tablet. One Step.

NDC 69536-162-99



levonorgestrel 1.5 mg

emergency contraceptive

Reduces chance of pregnancy after unprotected sex.  
 One Tablet. One Step.  
 Not for regular birth control.



1 Tablet Levonorgestrel 1.5 mg



3 169536 16300 0

**Plan B One-Step**  
 emergency contraceptive  
 One Tablet. One Step.

The sooner you take it, the more effective it will be




Take as soon as possible within 72 hours (3 days) after unprotected sex

Will not harm an existing pregnancy

Visit us at:  
[www.PlanBOneStep.com](http://www.PlanBOneStep.com)

Distributed by:  
 Foundation Consumer Healthcare, LLC (FCH)  
 Pittsburgh, PA 15212  
 ©2022  
 Made in Hungary  
 0000000000



<p><b>What is Plan B One-Step®?</b>                  Plan B One-Step is emergency contraception that helps prevent pregnancy after birth control failure or unprotected sex. It is a <b>backup</b> method of preventing pregnancy and should not be used as regular birth control.</p> <p><b>What Plan B One-Step is not.</b>                  Plan B One-Step will not work if you are already pregnant and will not affect an existing pregnancy. Plan B One-Step will not protect you from HIV infection (the virus that causes AIDS) and other sexually transmitted diseases (STDs).</p> <p><b>When should I use Plan B One-Step?</b>                  The sooner you take emergency contraception, the better it works. You should use Plan B One-Step within 72 hours (3 days) <b>after you have had unprotected sex.</b> Plan B One-Step is a backup or emergency method of birth control you can use when:</p> <ul style="list-style-type: none"> <li>• your regular birth control was used incorrectly or failed</li> <li>• you did not use any birth control method</li> </ul> <p><b>When not to use Plan B One-Step.</b>                  Plan B One-Step should not be used:</p> <ul style="list-style-type: none"> <li>• as a regular birth control method, because it's not as effective as regular birth control.</li> <li>• if you are already pregnant, because it will not work.</li> <li>• if you are allergic to levonorgestrel or any other ingredients in Plan B One-Step.</li> </ul>	<p><b>When should I talk to a doctor or pharmacist?</b>                  Ask a doctor or pharmacist before use if you are taking efavirenz (HIV medication) or rifampin (tuberculosis treatment) or medication for seizures (epilepsy). These medications may reduce the effectiveness of Plan B One-Step and increase your chance of becoming pregnant. Your doctor may prescribe another form of emergency contraception that may not be affected by these medications.</p> <p><b>How does Plan B One-Step work?</b>                  Plan B One-Step works before release of an egg from the ovary. As a result, Plan B One-Step usually stops or delays the release of an egg from the ovary. Plan B One-Step is one tablet that contains a higher dose of levonorgestrel than birth control pills and works in a similar way to prevent pregnancy.</p> <p><b>How can I get the best results from Plan B One-Step?</b>                  You have 72 hours (3 days) to try to prevent pregnancy after birth control failure or unprotected sex. <b>The sooner you take Plan B One-Step, the better it works.</b></p> <p><b>How effective is Plan B One-Step?</b>                  If Plan B One-Step is taken as directed, it can significantly decrease the chance that you will get pregnant. <b>About 7 out of every 8 women</b> who would have gotten pregnant will not become pregnant.</p> <p><b>How will I know Plan B One-Step worked?</b>                  You will know Plan B One-Step has been effective when you get your next period, which should come at the expected time, or within a week of the expected time. If your period is delayed beyond 1 week, it is possible you may be pregnant. You should get a pregnancy test and follow up with your healthcare professional.</p> <p><b>Will I experience any side effects?</b></p> <ul style="list-style-type: none"> <li>• some women may have changes in their period, such as a period that is heavier or lighter or a period that is early or late. <b>If your period is more than a week late, you may be pregnant.</b></li> <li>• if you have severe abdominal pain, you may have an ectopic pregnancy, and should get immediate medical attention.</li> <li>• when used as directed, Plan B One-Step is safe and effective. Side effects may include changes in your period, nausea, lower stomach (abdominal) pain, tiredness, headache, dizziness, and breast tenderness.</li> <li>• if you vomit within 2 hours of taking the medication, call a healthcare professional to find out if you should repeat the dose.</li> </ul>
	<p><b>What if I still have questions about Plan B One-Step?</b>                  If you have questions or need more information, call our toll-free number, 1-800-330-1271, or visit our website at <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a>.</p> <p><b>Other Information</b>  <b>Keep out of reach of children:</b>                  In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.</p> <p><b>Do not use if the blister seal is opened.</b>                  Store at room temperature 20–25°C (68–77°F).</p> <p><b>Active ingredient:</b> levonorgestrel 1.5 mg</p> <p><b>Inactive ingredients:</b> colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, potato starch, talc</p> <p>1-800-330-1271  <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a></p>
	<p><b>What if I still have questions about Plan B One-Step?</b>                  If you have questions or need more information, call our toll-free number, 1-800-330-1271, or visit our website at <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a>.</p> <p><b>Other Information</b>  <b>Keep out of reach of children:</b>                  In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.</p> <p><b>Do not use if the blister seal is opened.</b>                  Store at room temperature 20–25°C (68–77°F).</p> <p><b>Active ingredient:</b> levonorgestrel 1.5 mg</p> <p><b>Inactive ingredients:</b> colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, potato starch, talc</p> <p>1-800-330-1271  <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a></p>
<p>Found at <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a>                  Rev. 0/0000 ©2022 Printed in USA</p>	<p><b>What if I still have questions about Plan B One-Step?</b>                  If you have questions or need more information, call our toll-free number, 1-800-330-1271, or visit our website at <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a>.</p> <p><b>Other Information</b>  <b>Keep out of reach of children:</b>                  In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.</p> <p><b>Do not use if the blister seal is opened.</b>                  Store at room temperature 20–25°C (68–77°F).</p> <p><b>Active ingredient:</b> levonorgestrel 1.5 mg</p> <p><b>Inactive ingredients:</b> colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, potato starch, talc</p> <p>1-800-330-1271  <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a></p>
<p>If you are sexually active, you should see a healthcare provider for routine checkups. Your healthcare provider will talk to you about and, if necessary, test you for sexually transmitted diseases. Teach you about effective methods of routine birth control, and answer any other questions you may have.</p>	<p><b>What if I still have questions about Plan B One-Step?</b>                  If you have questions or need more information, call our toll-free number, 1-800-330-1271, or visit our website at <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a>.</p> <p><b>Other Information</b>  <b>Keep out of reach of children:</b>                  In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.</p> <p><b>Do not use if the blister seal is opened.</b>                  Store at room temperature 20–25°C (68–77°F).</p> <p><b>Active ingredient:</b> levonorgestrel 1.5 mg</p> <p><b>Inactive ingredients:</b> colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, potato starch, talc</p> <p>1-800-330-1271  <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a></p>
	<p><b>What if I still have questions about Plan B One-Step?</b>                  If you have questions or need more information, call our toll-free number, 1-800-330-1271, or visit our website at <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a>.</p> <p><b>Other Information</b>  <b>Keep out of reach of children:</b>                  In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.</p> <p><b>Do not use if the blister seal is opened.</b>                  Store at room temperature 20–25°C (68–77°F).</p> <p><b>Active ingredient:</b> levonorgestrel 1.5 mg</p> <p><b>Inactive ingredients:</b> colloidal silicon dioxide, corn starch, lactose monohydrate, magnesium stearate, potato starch, talc</p> <p>1-800-330-1271  <a href="http://www.PlanBOneStep.com">www.PlanBOneStep.com</a></p>