

**Twin Pack!**  
2 x 34 Doses

**EASY to Pour Bottle**

**MiraLAX<sup>®</sup>**  
*Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative*

**UNFLAVORED  
POWDER  
GRIT FREE**

**34 ONCE-DAILY DOSES**      NET WT 20.4 OZ (578 g)

**Bayer**

*MiraLAX<sup>®</sup> increases frequency of bowel movements and softens the stool.*

Bayer, the Bayer Cross, and MiraLAX are registered trademarks of Bayer. © 2015 Bayer. Dist by: Bayer HealthCare LLC Whippany, NJ 07981

00000-00


Lot  
Expiry Area

00000-00      Bayer

0 41100 80926 1

• *Dissolves in Any Beverage*

FDA Submission: 05.06.15



(b) (4)





**Drug Facts**

**Active ingredient (in each dose) Purpose**  
 Polyethylene Glycol 3350,  
 17 g (cap filled to line)..... Osmotic Laxative

**Use**  
 ■ relieves occasional constipation (irregularity)  
 ■ generally produces a bowel movement in 1 to 3 days

**Warnings**  
**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**  
 ■ nausea, vomiting or abdominal pain  
 ■ a sudden change in bowel habits that lasts over 2 weeks  
 ■ irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**  
 ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.  
 ■ you get diarrhea  
 ■ you need to use a laxative for longer than 1 week

**Drug Facts (continued)**

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not take more than directed unless advised by your doctor  
 ■ the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)  
 ■ adults and children 17 years of age and older:  
   ■ fill to top of white section in cap which is marked to indicate the correct dose (17g)  
   ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink  
   ■ use once a day  
   ■ use no more than 7 days  
 ■ children 16 years of age or under: ask a doctor

**Other information**  
 ■ store at 20° - 25°C (68° - 77°F)  
 ■ tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

**Inactive ingredients** none

**Questions or comments?**  
 1-800-MiraLAX (1-800-647-2529) or [www.MiraLAX.com](http://www.MiraLAX.com)

• Sugar Free 38500-00

**Drug Facts**

**Active ingredient (in each dose) Purpose**  
 Polyethylene Glycol 3350,  
 17 g (cap filled to line)..... Osmotic Laxative

**Use**  
 ■ relieves occasional constipation (irregularity)  
 ■ generally produces a bowel movement in 1 to 3 days

**Warnings**  
**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**  
 ■ nausea, vomiting or abdominal pain  
 ■ a sudden change in bowel habits that lasts over 2 weeks  
 ■ irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**  
 ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.  
 ■ you get diarrhea  
 ■ you need to use a laxative for longer than 1 week

**Drug Facts (continued)**

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not take more than directed unless advised by your doctor  
 ■ the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)  
 ■ adults and children 17 years of age and older:  
   ■ fill to top of white section in cap which is marked to indicate the correct dose (17g)  
   ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink  
   ■ use once a day  
   ■ use no more than 7 days  
 ■ children 16 years of age or under: ask a doctor

**Other information**  
 ■ store at 20° - 25°C (68° - 77°F)  
 ■ tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken


**Inactive ingredients** none

**Questions or comments?**  
 1-800-MiraLAX (1-800-647-2529) or [www.MiraLAX.com](http://www.MiraLAX.com)

• Sugar Free 38500-00



FDA Submission: 05.06.15



ATTACH LEGEND ON ALL  
 DIGITAL MECHANICALS

(b) (4)





**Drug Facts**

**Active ingredient (in each dose) Purpose**  
 Polyethylene Glycol 3350,  
 17 g (cap filled to line)..... Osmotic Laxative

**Use**  
 ■ relieves occasional constipation (irregularity)  
 ■ generally produces a bowel movement in 1 to 3 days

**Warnings**  
**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**  
 ■ nausea, vomiting or abdominal pain  
 ■ a sudden change in bowel habits that lasts over 2 weeks  
 ■ irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**  
 ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.  
 ■ you get diarrhea  
 ■ you need to use a laxative for longer than 1 week

**Drug Facts (continued)**

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ **do not take more than directed unless advised by your doctor**  
 ■ the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)  
 ■ adults and children 17 years of age and older:  
   ■ fill to top of white section in cap which is marked to indicate the correct dose (17g)  
   ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink  
   ■ use once a day  
   ■ use no more than 7 days  
 ■ children 16 years of age or under: ask a doctor

**Other information**  
 ■ store at 20° - 25°C (68° - 77°F)  
 ■ tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

**Inactive ingredients** none

**Questions or comments?**  
 1-800-MiraLAX (1-800-647-2529) or [www.MiraLAX.com](http://www.MiraLAX.com)

• Sugar Free 38500-00

**Drug Facts**

**Active ingredient (in each dose) Purpose**  
 Polyethylene Glycol 3350,  
 17 g (cap filled to line)..... Osmotic Laxative

**Use**  
 ■ relieves occasional constipation (irregularity)  
 ■ generally produces a bowel movement in 1 to 3 days

**Warnings**  
**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**  
 ■ nausea, vomiting or abdominal pain  
 ■ a sudden change in bowel habits that lasts over 2 weeks  
 ■ irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**  
 ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.  
 ■ you get diarrhea  
 ■ you need to use a laxative for longer than 1 week

**Drug Facts (continued)**

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ **do not take more than directed unless advised by your doctor**  
 ■ the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)  
 ■ adults and children 17 years of age and older:  
   ■ fill to top of white section in cap which is marked to indicate the correct dose (17g)  
   ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink  
   ■ use once a day  
   ■ use no more than 7 days  
 ■ children 16 years of age or under: ask a doctor

**Other information**  
 ■ store at 20° - 25°C (68° - 77°F)  
 ■ tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

**Inactive ingredients** none

**Questions or comments?**  
 1-800-MiraLAX (1-800-647-2529) or [www.MiraLAX.com](http://www.MiraLAX.com)

• Sugar Free 38500-00

---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/

---

KAREN M MAHONEY

11/12/2015