

00000000



ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd, Bold Italic

Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 8 pt Helvetica Cnd.

Headings - 8 pt Helvetica Cnd, Bold Italic

Sub Heads - 6 pt Helvetica Cnd, Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Square Bullets - 5 pt Zaph Dingbats no compression

Border - 2.5 pt rule, 3/64" gap all around

Barline - 2.5 pt rule

Hairline - .5 pt rule

(b) (4)

NEW PACKAGING REVISIONS:

- 1** Replaced the side panel purse with new image (b) (4)
- 2** Changed (b) (4) to **Mix-In Pax**
- 3** Deleted the website "or www.MiraLAX.com" from the last section of the Drug Facts



ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch
Drug Facts - 9 pt Helvetica Cnd, Bold Italic
Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 8 pt Helvetica Cnd,
Headings - 8 pt Helvetica Cnd, Bold Italic
Sub Heads - 6 pt Helvetica Cnd, Bold
Text - 6 pt Helvetica Cnd, minimum leading 6.5
Square Bullets - 5 pt Zaph Dingbats no compression
Border - 2.5 pt rule, 3/64" gap all around
Barline - 2.5 pt rule
Hairline - .5 pt rule

NEW PACKAGING REVISIONS:

- 1** Replaced the side panel purse with new image 6067UAD
- 2** Changed ^{(b) (4)} to **Mix-In Pax**
- 3** Deleted the website "or www.MiraLAX.com" from the last section of the Drug Facts

(b) (4)



MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

- Unflavored
- Sugar Free
- Dissolves in Any Beverage



20
Single Doses

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

- Relieves Occasional Constipation/Irregularity
- Softens Stool



20
ONCE-DAILY DOSES

UNFLAVORED
POWDER
GRIT FREE

20 PACKETS
NET WT 0.5 OZ
(17g) EACH

MiraLAX
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax



Travel Size

20
PRE-MASURED, UNFLAVORED,
SINGLE DOSES

Drug Facts
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g
Purpose Osmotic Laxative

Use
relieves occasional constipation (irregularity)
generally produces a bowel movement in 1 to 3 days

Warnings
Allergy alert: Do not use if you are allergic to polyethylene glycol
Do not use if you have kidney disease, except under the advice and supervision of a doctor
Ask a doctor before use if you have
nausea, vomiting or abdominal pain
a sudden change in bowel habits that lasts over 2 weeks
irritable bowel syndrome
Ask a doctor or pharmacist before use if you are taking a
prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if
you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
you get diarrhea
you need to use a laxative for longer than 1 week

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
do not take more than directed unless advised by your doctor
adults and children 17 years of age and older:
sift and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
use once a day
use no more than 7 days
children 18 years of age or under: ask a doctor

Other information
store at 20° - 25°C (68° - 77°F)
tamper-evident: do not use if foil is open or broken

Inactive ingredients none

Questions or comments?
1-800-MiraLAX (1-800-647-2329)

Bayer, the Bayer Cross, and MiraLAX are registered trademarks of Bayer.
© 2016 Bayer.
Distributed by Bayer HealthCare LLC, Whippany, NJ 07981
XXXX-XX

Lot & Expiry
(Non - Varnish Area)
1.75"W X .75"H



Bayer

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd, Bold Italic

Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 8 pt Helvetica Cnd

Headings - 8 pt Helvetica Cnd, Bold Italic

Sub Heads - 6 pt Helvetica Cnd, Bold

Text - 6 pt Helvetica Cnd, minimum leading 6,5

Square Bullets - 5 pt Zaph Dingbats no compression

Border - 2.5 pt rule, 3/64" gap all around

Barline - 2.5 pt rule

Hairline - .5 pt rule

- NEW PACKAGING REVISIONS:**
- 1 Replaced the side panel purse with new image 6067UAD
 - 2 Changed  to Mix-In Pax
 - 3 Deleted the website "or www.MiraLAX.com" from the last section of the Drug Facts



ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch
Drug Facts - 9 pt Helvetica Cnd, Bold Italic
Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 8 pt Helvetica Cnd.
Headings - 8 pt Helvetica Cnd, Bold Italic
Sub Heads - 6 pt Helvetica Cnd, Bold
Text - 6 pt Helvetica Cnd, minimum leading 6.5
Square Bullets - 5 pt Zaph Dingbats no compression
Border - 2.5 pt rule, 3/64" gap all around
Barline - 2.5 pt rule
Hairline - .5 pt rule

NEW PACKAGING REVISIONS:

- 1** Replaced the side panel purse with new image 6067UAD
- 2** Changed  to **Mix-In Pax**
- 3** Deleted the website "or www.MiraLAX.com" from the last section of the Drug Facts



Drug Facts
Active Ingredient (in each dose) Purpose
Polyethylene Glycol 3350, 17 g Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings
Allergy alert: Do not use if you are allergic to polyethylene glycol.
Do not use if you have kidney disease, except under the advice and supervision of a doctor.
Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug.

When using this product you may have loose, watery, more frequent stools.
Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

Drug Facts (continued)
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- adults and children 17 years of age and older: stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- use once a day
- use no more than 7 days
- children 16 years of age or under: ask a doctor

Other Information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil is open or broken

Inactive ingredients none

Questions or comments?
1-800-MiraLAX (1-800-647-2529)

Bayer, the Bayer Cross, and MiraLAX are registered trademarks of Bayer.
© 2016 Bayer.
Dist by: Bayer HealthCare LLC, Whippany, NJ 07981
XXXX-XX Bayer

No Varnish
No Copy
Lot/Exp Area


00000000

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch
Drug Facts - 9 pt Helvetica Cnd, Bold Italic
Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 8 pt Helvetica Cnd.
Headings - 8 pt Helvetica Cnd, Bold Italic
Sub Heads - 8 pt Helvetica Cnd, Bold
Text - 8 pt Helvetica Cnd, minimum leading 7.5
Square Bullets - 5 pt Zaph Dingbats no compression
Border - 2.5 pt rule, 3/16" gap all around
Barline - 2.5 pt rule
Hairline - .5 pt rule

- NEW PACKAGING REVISIONS:**
- 1 Replaced the side panel purse with new image 6067UAD
 - 2 Changed (b) (4) to **Mix-In Pax**
 - 3 Deleted the website "or www.MiraLAX.com" from the last section of the Drug Facts

(b) (4) MiraLAX 17g 24ct



- NEW PACKAGING REVISIONS:**
- 1 Replaced the back panel purse with new image 6067UAD
 - 2 Changed  to **Mix-In Pax**
 - 3 Deleted the website "or www.MiraLAX.com" from the last section of the Drug Facts

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Indicates Content
Variable horizontal scale adheres to 50 characters per inch

Drug Facts - 9 pt Helvetica Cond. Bold Italic

Warnings - 9 pt Helvetica Cond. Bold Italic

Directions - 9 pt Helvetica Cond. Bold

Other information - 9 pt Helvetica Cond. Bold

Questions or comments? - 9 pt Helvetica Cond. Bold

Visit www.MiraLAX.com - 9 pt Helvetica Cond. Bold

Side Panels - 5 pt Helvetica Cond. Medium

Back Panel - 5 pt Helvetica Cond. Medium

Front Panel - 5 pt Helvetica Cond. Medium

Top Panel - 5 pt Helvetica Cond. Medium

Bottom Panel - 5 pt Helvetica Cond. Medium

00118

Not For Individual Sale

- **Unflavored**
- **Sugar Free**
- **Dissolves in Any Beverage**

Drug Facts	Drug Facts (continued)
<p>Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g.....Osmotic Laxative</p> <p>Purpose relieves occasional constipation (irregularity) and generally produces a bowel movement in 1 to 3 days</p> <p>Use relieves occasional constipation (irregularity) and generally produces a bowel movement in 1 to 3 days</p> <p>Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor Ask a doctor before use if you have nausea, vomiting or abdominal pain, a sudden change in bowel habits that lasts over 2 weeks, or irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug When using this product you may have loose, watery, more frequent stools Stop use and ask a doctor if you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. you get diarrhea you need to use a laxative for longer than 1 week</p>	<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions do not take more than directed unless advised by your doctor adults and children 17 years of age and older: stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink use once a day use no more than 7 days children 16 years of age or under: ask a doctor</p> <p>Other information store at 20° - 25°C (68° - 77°F) tamper-evident: do not use if foil is open or broken</p> <p>Inactive ingredients none</p> <p>Questions or comments? 1-800-MiraLAX (1-800-647-2529)</p> <p>Bayer, the Bayer Cross, and MiraLAX are registered trademarks of Bayer, © 2016 Bayer. Dist by: Bayer HealthCare LLC, Whippany, NJ 07981</p> <p style="text-align: center;">Bayer XXXXXXXX</p>

FOLD AT LINE
TEAR AT ARROW



MiraLAX[®]
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax Travel Size

- Relieves Occasional Constipation/Irregularity
- Softens Stool

UNFLAVORED POWDER GRIT FREE

1 DOSE

NET WT 0.5 OZ (17 g)

NO COPY - EYEMARK LANE

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd, Bold Italic

Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 8 pt Helvetica Cnd.

Headings - 8 pt Helvetica Cnd, Bold Italic

Sub Heads - 6 pt Helvetica Cnd, Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Square Bullets - 3.5 pt Zaph Dingbats no compression

Border - 1 pt rule, 3/64" gap all around

Barline - 1 pt rule

Hairline - .5 pt rule

(b) (4)

NEW PACKAGING REVISIONS:

- 1** Changed (b) (4) to **Mix-In Pax**
- 2** Deleted the website "or www.MiraLAX.com" from the last section of the Drug Facts

4.0000

4.5000

 N O 00000-00000 0	<ul style="list-style-type: none"> • Unflavored • Sugar Free • Dissolves in any Beverage
Drug Facts	Drug Facts (continued)
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g.....Osmotic Laxative	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Use ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days	Directions ■ do not take more than directed unless advised by your doctor ■ adults and children 17 years of age and older: ■ stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ use once a day ■ use no more than 7 days ■ children 16 years of age or under: ask a doctor
Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor Ask a doctor before use if you have ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug When using this product you may have loose, watery, more frequent stools Stop use and ask a doctor if ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week	Other information ■ store at 20° - 25°C (68° - 77°F) ■ tamper-evident: do not use if foil is open or broken Inactive ingredients none Questions or comments? 1-800-MiraLAX (1-800-647-2529)
Bayer	Bayer, the Bayer Cross, and MiraLAX are registered trademarks of Bayer. © 2016 Bayer. Dist by: Bayer HealthCare LLC, Whippany, NJ 07981 XXXXXXXX



MiraLAX[®]

Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

Travel Size

- Relieves Occasional Constipation/Irregularity
- Softens Stool

UNFLAVORED
POWDER
 GRIT FREE



1 DOSE

NET WT 0.5 OZ (17 g)

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT Helvetica Condensed Variable horizontal scale adheres to 39 characters per inch Drug Facts - 9 pt Helvetica Cnd, Bold Italic Drug Facts (continued) Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 8 pt Helvetica Cnd. Headings - 8 pt Helvetica Cnd, Bold Italic Sub Heads - 6 pt Helvetica Cnd, Bold Text - 6 pt Helvetica Cnd, minimum leading 6.5 Square Bullets - 3.5 pt Zaph Dingbats no compression Border - 1 pt rule, 3/64" gap all around Bartline - 1 pt rule Hairline - .5 pt rule
--

(b) (4)

NEW PACKAGING REVISIONS:

- 1** Changed (b) (4) to **Mix-In Pax**
- 2** Deleted the website "or www.MiraLAX.com" from the last section of the Drug Facts


4.0000

4.5000

Sample-Not For Sale

- Unflavored
- Sugar Free
- Dissolves in any Beverage

Drug Facts	Drug Facts (continued)								
<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <th style="text-align: left;">Active ingredient (in each dose)</th> <th style="text-align: left;">Purpose</th> </tr> <tr> <td>Polyethylene Glycol 3350, 17 g.....Osmotic Laxative</td> <td></td> </tr> <tr> <th style="text-align: left;">Use</th> <td> <ul style="list-style-type: none"> ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days </td> </tr> <tr> <th style="text-align: left;">Warnings</th> <td> <p>Allergy alert: Do not use if you are allergic to polyethylene glycol</p> <p>Do not use if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome <p>Ask a doctor or pharmacist before use if you are taking a prescription drug</p> <p>When using this product you may have loose, watery, more frequent stools</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week </td> </tr> </table>	Active ingredient (in each dose)	Purpose	Polyethylene Glycol 3350, 17 g.....Osmotic Laxative		Use	<ul style="list-style-type: none"> ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days 	Warnings	<p>Allergy alert: Do not use if you are allergic to polyethylene glycol</p> <p>Do not use if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome <p>Ask a doctor or pharmacist before use if you are taking a prescription drug</p> <p>When using this product you may have loose, watery, more frequent stools</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week 	<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed unless advised by your doctor ■ adults and children 17 years of age and older: <ul style="list-style-type: none"> ■ stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ use once a day ■ use no more than 7 days ■ children 16 years of age or under: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ store at 20° - 25°C (68° - 77°F) ■ tamper-evident: do not use if foil is open or broken <p>Inactive ingredients none</p> <p>Questions or comments? 1-800-MiraLAX (1-800-647-2529)</p> <p><small>Bayer, the Bayer Cross, and MiraLAX are registered trademarks of Bayer. © 2016 Bayer. Dist by: Bayer HealthCare LLC, Whippany, NJ 07981</small></p>
Active ingredient (in each dose)	Purpose								
Polyethylene Glycol 3350, 17 g.....Osmotic Laxative									
Use	<ul style="list-style-type: none"> ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days 								
Warnings	<p>Allergy alert: Do not use if you are allergic to polyethylene glycol</p> <p>Do not use if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome <p>Ask a doctor or pharmacist before use if you are taking a prescription drug</p> <p>When using this product you may have loose, watery, more frequent stools</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week 								



MiraLAX®

Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

Mix-In Pax

Travel Size

- Relieves Occasional Constipation/Irregularity
- Softens Stool

UNFLAVORED
POWDER
 GRIT FREE

1 DOSE

NET WT 0.5 OZ (17 g)

(b) (4)

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

<p>STANDARD FORMAT Helvetica Condensed Variable horizontal scale adheres to 39 characters per inch</p> <p>Drug Facts - 9 pt Helvetica Cnd, Bold Italic</p> <p>Drug Facts (continued) Drug Facts - 8 pt Helvetica Cnd, Bold Italic (continued) - 8 pt Helvetica Cnd.</p> <p>Headings - 8 pt Helvetica Cnd, Bold Italic</p> <p>Sub Heads - 6 pt Helvetica Cnd, Bold</p> <p>Text - 6 pt Helvetica Cnd, minimum leading 6.5</p> <p>Square Bullets - 3.5 pt Zaph Dingbats no compression</p> <p>Border - 1 pt rule, 3/64" gap all around</p> <p>Barline - 1 pt rule</p> <p>Hairline - .5 pt rule</p>

(b) (4)

NEW PACKAGING REVISIONS:

- 1 Changed (b) (4) to **Mix-In Pax**
- 2 Deleted the website "or www.MiraLAX.com" from the last section of the Drug Facts

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY
09/30/2016