

**2 FREE Single Dose Packets**  
**MIX-IN PAX** SEE BACK

**2 FREE Single Dose Packets**  
**MIX-IN PAX**

**MIRALAX**  
Polyethylene Glycol 3350, Powder for Solution, Osmotic Laxative

**UNFLAVORED POWDER**  
GRIT FREE

- Relieves Occasional Constipation/Irregularity
- Softens Stool

**16** ONCE-DAILY DOSES

1 BOTTLE - NET WT 8.3 OZ (238 g)  
2 PACKETS - NET WT 0.5 OZ (17 g) EACH  
TOTAL NET WT 9.3 OZ (272 g)

**Drug Facts**

**Active ingredient (in each dose)** Polyethylene Glycol 3350, 17g (cap filled to line)..... Osmotic Laxative

**Purpose**

**Use**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

**Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are taking a prescription drug**

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

**Drug Facts (continued)**

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - fill to top of white section in cap which is marked to indicate the correct dose (17g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once a day
  - use no more than 7 days
  - children 16 years of age or under: ask a doctor

**Other information**

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

**Inactive ingredients** none

**Questions or comments?**  
1-800-MiraLAX (1-800-647-2529)

**LOT/EXP AREA**

**Actual Size 100%**

DRUG FACTS INFO		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt **	8 pt
Drug Facts Continued	8 pt	8 pt
Headings*	8 pt bold italic*	8 pt
Subheadings	≥6 pt	6 pt
Body Text	≥6 pt	6 pt
Bullets	5 pt	3.5 pt
Leading	≥0.5 pt	0.5 pt
Hairlines	0.5 pt rule	.5 pt
Box and Barlines†	>0.5 pt rule	1 pt
Characters Per Inch:	No more than 38	> 38

\* Headings are to be at least 20% larger than Body Text.  
\*\* Drug Facts Title to be larger than all other type size used in DFP  
† FDA Recommends 2.5pt rule

  
 0 000000 000000

Bayer  
 00000-00  
 Whippany, NJ 07981  
 Dist by: Bayer HealthCare LLC,  
 © 2018 Bayer,  
 are registered trademarks of Bayer,  
 Bayer, the Bayer Cross, and MiraLAX





**Drug Facts**

**Active ingredient (in each dose)**      **Purpose**  
Polyethylene Glycol 3350,  
17g (cap filled to line) ..... Osmotic Laxative

**Use**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

**Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

**Drug Facts (continued)**

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - fill to top of white section in cap which is marked to indicate the correct dose (17g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once a day
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

**Other information**

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

**Inactive ingredients**      none

**Questions or comments?**  
1-800-MiraLAX (1-800-647-2529)

---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/  
-----

KAREN M MAHONEY  
05/04/2018