



Actual Size 100%

FDA60502-1

DRUG FACTS INFO (Updated 5/18/18)		
CATEGORY	REQUIRED	ACTUAL IN FILE
Drug Facts Title	>8 pt. **	9 pt.
Drug Facts Continued	8 pt.	8 pt.
Headings*	8 pt.-bold italic*	8 pt.
Subheadings	≥6 pt.	6 pt.
Body Text	≥6 pt.	6 pt.
Bullets	5 pt.	5 pt.
Leading	≥0.5 pt.	0.5 pt.
Headlines	0.5 pt. rule	0.5 pt.
Box and Barlines†	>0.5 pt. rule	2.5 pt.
Characters Per Inch:	No more than 39	≤39

* Headings are to be at least 2pts larger than Body Text.
** Drug Facts Title to be larger than all other type size used in DFP
† FDA Recommends 2.5pt rule

PRE-MEASURED
3 SINGLE DOSES



Drug Facts
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 g
Purpose Osmotic Laxative
Use
■ relieves occasional constipation (irregularity)
■ generally produces a bowel movement in 1 to 3 days
Warnings
Allergy alert: Do not use if you are allergic to polyethylene glycol.
Do not use if you have kidney disease, except under the advice and supervision of a doctor.
Ask a doctor before use if you have
■ nausea, vomiting or abdominal pain
■ a sudden change in bowel habits that lasts over 2 weeks
■ irritable bowel syndrome
Ask a doctor or pharmacist before use if you are taking a prescription drug
When using this product you may have loose, watery, more frequent stools

Drug Facts (continued)
Stop use and ask a doctor if
■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
■ you get diarrhea
■ you need to use a laxative for longer than 1 week
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ do not take more than directed unless advised by your doctor
■ adults and children 17 years of age and older:
■ use once a day
■ stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
■ do not combine with starch-based thickeners used for difficulty swallowing
■ ensure that the powder is fully dissolved before drinking
■ do not drink if there are any clumps
■ do not use more than 7 days
■ children 16 years of age or under: ask a doctor
Other information
■ store at 20° - 25°C (68° - 77°F)
■ tamper-evident: do not use if foil is open or broken
Inactive ingredients none
Questions or comments?
1-800-MiraLAX (1-800-647-2529)

Lot & Expiry (Non-varnish Area)
1.50"W x 0.75"H

Bayer, the Bayer Cross, Mix-In Pax and MiraLAX are registered trademarks of Bayer.
© 2023 Bayer. Pat. patents.livewell.bayer.com
Dist by: Bayer HealthCare LLC, Whippany, NJ 07981
00000

Bayer



This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
10/27/2023 04:41:12 PM