

FS67_6b_BP3_130606

Labeling Format Information:			
Fonts : Helvetica Roman , Bold , Bold Oblique			
Drug Facts :	9.5pt	Leading :	6.5pt
Header :	8pt	Bullets :	5pt
Subheader :	6pt	Barlines :	2.5pt
Body Text :	6pt	Hairlines :	0.5pt
Horizontal Scale: 80%-100%		Average Kerning: 0	

100%出力



Labeling Format Information:		
Fonts : Helvetica BoldOblique Helvetica 77 Bold Condensed Oblique Helvetica 57 Condensed Helvetica Condensed Bold		
Drug Facts :	9.5pt	Leading : 2pt
Header :	8pt	Bullets : 5pt
Subheader :	6.5pt	Barlines : 2pt
Body Text :	6pt	Hairlines : 0,4pt
Drug Facts (Continued) NA		
Horizontal Scale: 80%-100% Average Kerning: 0		

100%出力
 FS67_3s_f_130606

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

FOLD TWICE AFTER OPENING

Discard unused patches 14 days after opening

FOLD TWICE AFTER OPENING



1
Peel the center film.



2
Place on affected area.



3
Slide the film forward.



4
Slide the other film backward.



Salonpas®
Pain Relief Patch

Menthol 3%
Methyl Salicylate 10%
Topical Analgesic

3 patches
 2 3/4" X 3 15/16"
 (7cm X 10cm)

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
 EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

※ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Hisamitsu®
 Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
 JAPAN SAGA TOSU
 MADE IN JAPAN

Distributed by
Hisamitsu America, Inc.
 Florham Park, NJ 07932

LOT

EXP.

Package not child resistant.
 Keep out of reach of children.

© : Registered Trade Mark

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

Salonpas® Pain Relief Patch Menthol 3%, Methyl Salicylate 10% minty scent

Drug Facts	Purpose
Active ingredients (in each patch)	
Menthol 3%	Topical analgesic
Methyl salicylate 10%(NSAID : nonsteroidal anti-inflammatory drug)	Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
 ■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

Warnings
 For external use only
Stomach bleeding warning:
 This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Do not use
 ■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs
 ■ with a heating pad ■ when sweating (such as from exercise or heat)
 ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

Ask a doctor before use if
 ■ you are allergic to topical products ■ the stomach bleeding warning applies to you
 ■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease

When using this product ■ wash hands after applying or removing patch. Avoid contact with eyes.
 If eye contact occurs, rinse thoroughly with water.
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if ■ stomach pain or upset gets worse or lasts
 ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
 ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days
 ■ symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions
Adults 18 years and older: ■ only use one patch at a time
 ■ clean and dry affected area ■ remove patch from backing film and apply to skin (see illustration)
 ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours
 ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
 ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row
 ■ the used patch should be removed from the skin when a new one is applied
Children under 18 years of age: do not use; this product has not been shown to work in children

Other information
 ■ store at 20-25°C (68-77°F) ■ avoid storing product in direct sunlight
 ■ protect product from excessive moisture
 ■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments? www.salonpas.us
 Toll free 1-87-SALONPAS

1L835T-0

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
01/31/2014