

Salonpas[®] Pain Relief Patch **LARGE**

Menthol 3%, Methyl Salicylate 10%

3 patches

NDC#46561-675-03

Hisamitsu[®]
LARGE

for temporary relief of mild to moderate pain

- Backache
- Sore Muscle
- Strains
- Bruises



Salonpas[®] Pain Relief Patch

APPLY FOR
8-12
HOURS

Menthol 3%
Methyl Salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

3 **LARGE** patches

3 15/16" X 5 1/2" (10cm X 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Package not child resistant.
Keep out of reach of children.

LOT
EXP.
Salonpas[®] Pain Relief Patch **LARGE** patches
Menthol 3%, Methyl Salicylate 10%



Manufactured by Hisamitsu Pharmaceutical Co., Inc. TORANCE, CA 90503
Distributed by Hisamitsu America, Inc. TORANCE, CA 90503
© Registered Trade Mark

Salonpas[®] Pain Relief Patch **LARGE** patches
Menthol 3%, Methyl Salicylate 10%



press open
menthol 3%, Methyl Salicylate 10%
minimally scented

Drug Facts

Active Ingredients (in each patch)
Menthol 3%..... Topical analgesic
Methyl salicylate 10%(NSAID : nonsteroidal anti-inflammatory drug)..... Topical analgesic

Uses

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
• strains • sprains • simple backache • arthritis • bruises

Warnings

For external use only
Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
• have 3 or more alcoholic drinks every day while using this product.
Take more or for a longer time than directed

Do not use

- on the face or rashes
- with a heating pad
- any patch from a pouch that has been open for 14 or more days
- if allergic to topical products
- you are allergic to aspirin or other NSAIDs
- you are taking a diuretic
- you have high blood pressure, heart disease, or kidney disease
- you are taking a diuretic

When using this product

- wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- rash, itching or skin irritation develops
- symptoms clear up and occur again within a few days
- symptoms last for more than 3 days
- symptoms clear up and occur again within a few days

Directions

Adults 18 years and older:
• only use one patch at a time
• clean and dry affected area
• apply one patch to the affected area and leave in place for up to 8 to 12 hours
• if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
• do not use more than 2 patches per day
• do not use for more than 3 days in a row
• the used patch should be removed from the skin when a new one is applied
Children under 18 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid storing product in direct sunlight
- protect product from excessive moisture
- some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetix: aluminum silicate

Questions or comments?

Toll free 1-87-SALONPAS, Weekdays, 9AM to 5PM (PST) | www.salonpas.us

24m
0.49m
3mning-0

CUT OPEN HERE ----- CUT OPEN HERE

FOLD TWICE AFTER OPENING
Discard unused patches 14 days after opening
FOLD TWICE AFTER OPENING



1 Peel the center film.



2 Place on affected area.



3 Slide a film.



4 Slide the other film.



Salonpas®
Pain Relief Patch

**Menthol 3%
Methyl Salicylate 10%
Topical Analgesic**

3 LARGES
3 1⁵/₁₆" X 5 1²/₁₆"
(10cm X 14cm)

3 PATCHES

EFFE CTIVENESS CONFIRMED IN CLINICAL TRIAL

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Hisamitsu®

Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
JAPAN SAGA TOSU
MADE IN JAPAN
Distributed by
Hisamitsu America, Inc.
Irvine, CA 92618


Package not child resistant.
Keep out of reach of children.

LOT: _____
EXP: _____

© Registered Trade Mark

CUT OPEN HERE ----- CUT OPEN HERE

FOLD TWICE AFTER OPENING
Discard unused patches 14 days after opening
FOLD TWICE AFTER OPENING



Salonpas® Pain Relief Patch LARGE Menthol 3%, Methyl Salicylate 10%, Methyl Salicylate 10%, Methyl Salicylate 10%, Methyl Salicylate 10%

**Menthol 3%
Methyl Salicylate 10%
Topical Analgesic**

3 LARGES
3 1⁵/₁₆" X 5 1²/₁₆"
(10cm X 14cm)

3 PATCHES

EFFE CTIVENESS CONFIRMED IN CLINICAL TRIAL

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Hisamitsu®

Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
JAPAN SAGA TOSU
MADE IN JAPAN
Distributed by
Hisamitsu America, Inc.
Irvine, CA 92618

Package not child resistant.
Keep out of reach of children.

LOT: _____
EXP: _____

© Registered Trade Mark

Drug Facts

Active ingredients (in each patch)

Menthol 3%
Methyl salicylate 10% (NSAID - nonsteroidal anti-inflammatory drug)

Purpose
Topical analgesic
Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
 ■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

Warnings
For external use only

Stomach bleeding warning:
This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Do not use
 ■ on the face or rashes ■ on wounds or damaged skin ■ If allergic to aspirin or other NSAIDs
 ■ with a heating pad ■ when sweating (such as from exercise or heat)
 ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

Ask a doctor before use if
 ■ you are allergic to topical products ■ the stomach bleeding warning applies to you
 ■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease

When using this product ■ wash hands after applying or removing patch. Avoid contact with eyes.
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
 ■ you feel faint, vomit blood, or have bloody or upset gets worse or lasts ■ stomach pain or upset gets worse or lasts
 ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days
 ■ symptoms clear up and occur again within a few days

If pregnant or breastfeeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child-resistant. Dispose of the used patches after folding sticky ends together.

Directions
Adults 18 years and older:
 ■ only use one patch at a time
 ■ clean and dry affected area
 ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours
 ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
 ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row
 ■ the used patch should be removed from the skin when a new one is applied

Children under 18 years of age: do not use

Other information
 ■ avoid storing product in direct sunlight
 ■ store at 20-25°C (68-77°F)
 ■ protect product from excessive moisture
 ■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients atyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
 toll free 1-877-SALONPAS Weekdays 9AM to 5PM (PST) www.salonpas.us

1L7721-0

Salonpas[®] Pain Relief Patch **LARGE** 2 patches

Menthol 3%, Methyl Salicylate 10%

Hisamitsu[®]

LARGE

for temporary relief of mild to moderate pain

- Backache
- Sore Muscle
- Strains
- Bruises



NDC#46581-675-02

Salonpas[®] Pain Relief Patch **LARGE** 2 patches
Menthol 3%, Methyl Salicylate 10%

Salonpas[®] Pain Relief Patch

APPLY FOR
8-12
HOURS

Menthol 3%
Methyl salicylate 10%
Topical Analgesic

2 LARGE patches

3 15/16" X 5 1/2" (10cm X 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS.

Package not child resistant.
Keep out of reach of children.

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

LOT
EXP

Salonpas[®] Pain Relief Patch **LARGE** 2 patches
Menthol 3%, Methyl Salicylate 10%



Manufactured by Hisamitsu Pharmaceutical Co., Inc. Torrance, CA 90503
Distributed by Hisamitsu America, Inc. Torrance, CA 90503
Menthol 3%, Methyl Salicylate 10%
MADE IN JAPAN
© Registered Trade Mark



press open
menthy scent

Drug Facts

Active Ingredients (in each patch)
Menthol 3%..... Topical analgesic
Methyl salicylate 10%(NSAID : nonsteroidal anti-inflammatory drug)..... Topical analgesic

Uses
Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

Warnings

For external use only
Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
take more or for a longer time than directed

Do not use:
■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs
■ with a heating pad ■ when sweating (such as from exercise or heat)
■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery
Ask a doctor before use if
■ you are allergic to topical products ■ the stomach bleeding warning applies to you
■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease
When using this product:
■ wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Directions

Stop use and ask a doctor if:
■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days
■ symptoms clear up and occur again within a few days
If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Adults 18 years and older:

- only use one patch at a time
- clean and dry affected area
- apply one patch to the affected area and leave in place for up to 8 to 12 hours
- if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- do not use more than 2 patches per day
- do not use for more than 3 days in a row
- the used patch should be removed from the skin when a new one is applied

Children under 18 years of age: do not use

Other Information

- store at 20-25°C (68-77°F)
- avoid storing product in direct sunlight
- protect product from excessive moisture
- some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?

Toll free 1-87-SALONPAS, Weekdays, 9AM to 5PM (PST) | www.salonpas.us

2pr
0.4pr
spring: 0

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

Discard unused patches 14 days after opening

FOLD TWICE AFTER OPENING



Salonpas®

Pain Relief Patch

Menthol 3%
Methyl Salicylate 10%
Topical Analgesic

2 LARGE
patches

3 1/8" x 5 1/2"
(10cm x 14cm)

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

✳ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Hisamitsu®

Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
1000 N. Main St.
MADISON, NJ 07040
MADE IN JAPAN
Distributed by
Hisamitsu America, Inc.
Torrance, CA 90503

Package not child resistant.
Keep out of reach of children.

LOT

EXP.

® : Registered Trade Mark

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

Salonpas® Pain Relief Patch LARGE Menthol 3%, Methyl Salicylate 10%, Minty Scent

Drug Facts

Active ingredients (in each patch)

Menthol 3%
Methyl salicylate 10% (NSAID - nonsteroidal anti-inflammatory drug)
Topical analgesic
Topical analgesic

Uses

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
 ■ sprains ■ simple backache ■ arthritis ■ bruises

Warnings

For external use only
Stomach bleeding warning:
 This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Do not use

■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs
 ■ with a heating pad ■ when sweating (such as from exercise or heat)
 ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

Ask a doctor before use if

■ you are taking a diuretic ■ the stomach bleeding warning applies to you
 ■ you are allergic to topical products ■ you have high blood pressure, heart disease, or kidney disease
 When using this product ■ wash hands after applying or removing patch. Avoid contact with eyes.
 If eye contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if

■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
 ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days
 ■ symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. D spouse of the used patches after folding sticky ends together.

Directions

Adults 18 years and older:
 ■ only use one patch at a time
 ■ clean and dry affected area ■ remove patch from backing film and apply to skin (see illustration)
 ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours
 ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
 ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row
 ■ the used patch should be removed from the skin when a new one is applied
Children under 18 years of age: do not use

Other information

■ store at 20-25°C (68-77°F) ■ avoid storing product in direct sunlight
 ■ protect product from excessive moisture
 ■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200 000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments? www.salonpas.us
 Toll free 1-877-SALONPAS, Weekdays, 9AM to 5PM (PST) 1L771T-0



2PR
 0.4Pr
 Printing: 0



✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

Drug Facts

Active ingredients (in each patch)

Menthol 3%..... Topical analgesic
 Methyl salicylate 10%(NSAID) : nonsteroidal anti-inflammatory drug..... Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
 ■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

Warnings

For external use only

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Do not use
 ■ on the face or rashes
 ■ on wounds or damaged skin
 ■ with a heating pad
 ■ when sweating (such as from exercise or heat)
 ■ any patch from a pouch that has been open for 14 or more days
 ■ right before or after heart surgery

Ask a doctor before use if
 ■ you are allergic to topical products
 ■ the stomach bleeding warning applies to you
 ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, or kidney disease

When using this product
 ■ wash hands after applying or removing patch. Avoid contact with eyes.
 ■ If eye contact occurs, rinse thoroughly with water.
 ■ If the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
 ■ stomach pain or upset gets worse or lasts
 ■ rash, itching or skin irritation develops
 ■ condition worsens
 ■ symptoms last for more than 3 days
 ■ symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If used in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions

Adults 18 years and older:
 ■ only use one patch at a time
 ■ clean and dry affected area
 ■ remove patch from backing film and apply to skin (see illustration)
 ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours
 ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
 ■ do not use more than 2 patches per day
 ■ do not use for more than 3 days in a row
 ■ the used patch should be removed from the skin when a new one is applied

Children under 18 years of age: do not use

Other information
 ■ store at 20-25°C (68-77°F) ■ avoid storing product in direct sunlight
 ■ protect product from excessive moisture
 ■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients
 alkylic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
 Toll free 1-877-SALONPAS, Weekdays, 9AM to 5PM (PST)
www.salonpas.us

LOT
 EXP

© : Registered Trade Mark 1L7701-0

2pr
 0.4pr
 printing: 0

Salonpas
Pain Relief Patch

APPLY FOR 8-12 HOURS

1 LARGE patch
 3 1/8" x 5 1/8" (10cm x 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

**MENTHOL 3%
 METHYL SALICYLATE 10%
 Topical Analgesic**

**FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
 EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL**

FREE SAMPLE NOT FOR RE-SALE

minty scent

for temporary relief of mild to moderate pain

Backache
 Sore Muscle
 Strains
 Bruises

1 Peel the center film.
 2 Place on affected area.
 3 Slide a film.
 4 Slide the other film.

Manufactured by
 Hisamitsu Pharmaceutical Co., Inc.
 Hiratsuka, Japan
 MADE IN JAPAN

Distributed by
 Hisamitsu America, Inc.
 Torrance, CA 90503

DRUG FACTS

Active ingredients (in each patch)
 Menthol 3%
 Methyl salicylate 10% (NSAID: nonsteroidal anti-inflammatory drug)
 Topical analgesic
 Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
 ■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

Warnings
 For external use only
 Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Do not use
 ■ on the face or rashes
 ■ on wounds or damaged skin
 ■ with a heating pad
 ■ when sweating (such as from exercise or heat)
 ■ any patch from a pouch that has been open for 14 or more days
 ■ right before or after heart surgery

Ask a doctor before use if
 ■ you are allergic to topical products
 ■ the stomach bleeding warning applies to you
 ■ you are taking a diuretic:
 ■ you have high blood pressure, heart disease, or kidney disease

When using this product
 ■ wash hands after applying or removing patch. Avoid contact with eyes.
 If eye contact occurs, rinse thoroughly with water.
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
 ■ stomach pain or upset gets worse or lasts
 ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
 ■ rash, itching or skin irritation develops
 ■ condition worsens
 ■ symptoms last for more than 3 days
 ■ symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions
Adults 18 years and older:
 ■ only use one patch at a time
 ■ clean and dry affected area
 ■ remove patch from backing film and apply to skin (see illustration)
 ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours
 ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
 ■ do not use more than 2 patches per day
 ■ do not use for more than 3 days in a row
 ■ the used patch should be removed from the skin when a new one is applied
Children under 18 years of age: do not use

Other information
 ■ store at 20-25°C (68-77°F)
 ■ avoid storing product in direct sunlight
 ■ protect product from excessive moisture
 ■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients
 salicylic acid, salicylic acid, hydrocodone resin, backing film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
 Toll free 1-877-SALONPAS, weekdays, 9AM to 5PM (PST)
 www.salonpas.us

LOT
EXP.

© : Registered Trade Mark 1L78910

Salonpas[®] Arthritis Pain Patch **LARGE** patches

Hisamitsu[®]

LARGE

for temporary relief of mild to moderate pain

- Arthritis
- Joint Pain
- Sprains
- Strains

NDC#46581-685-03

3 **Salonpas[®]** Arthritis Pain Patch **LARGE** patches
Menthol 3%, Methyl Salicylate 10%

Manufactured by Hisamitsu Pharmaceutical Co., Inc. JAPAN SAGA TOSU
Distributed by Hisamitsu America, Inc. Torrance, CA 90503
© : Registered Trade Mark



APPLY FOR **8-12** HOURS

Salonpas[®] Arthritis Pain Patch

Menthol 3%
Methyl salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

3 **LARGE** patches

3 1/8" x 5 1/2" (10cm x 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Package not child resistant.
Keep out of reach of children.

LOT
EXP.
Salonpas[®] Arthritis Pain Patch **LARGE** patches
Menthol 3%, Methyl salicylate 10%

3 **Salonpas[®]** Arthritis Pain Patch **LARGE** patches
Menthol 3%, Methyl Salicylate 10%



press open
menty scent

Drug Facts

Active ingredients (in each patch)
Menthol 3%..... Topical analgesic
Methyl salicylate 10%(NSAID : nonsteroidal anti-inflammatory drug)..... Topical analgesic

Uses
Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
■ arthritis ■ sprains ■ strains ■ bruises ■ simple backache

Warnings

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use

■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs
■ with a heating pad ■ when sweating (such as from exercise or heat)
■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

Ask a doctor before use if

■ you are allergic to topical products ■ the stomach bleeding warning applies to you
■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease

When using this product

■ wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
■ rash, itching or skin irritation develops. These are signs of stomach bleeding.
■ symptoms clear up and occur again within a few days ■ symptoms last for more than 3 days

Directions

Adults 18 years and older:
■ clean and dry affected area ■ remove patch from backing film and apply to skin (see illustration)
■ apply one patch to the affected area and leave in place for up to 8 to 12 hours
■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row
■ the used patch should be removed from the skin when a new one is applied

Other information

■ store at 20-25°C (68-77°F) ■ avoid storing product in direct sunlight
■ protect product from excessive moisture
■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene
1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?

Toll free 1-87-SALONPAS, Weekdays, 9AM to 5PM (PST) www.salonpas.us

2pr
0.4pr
mling: 0

CUT OPEN HERE ----- CUT OPEN HERE

FOLD TWICE AFTER OPENING
Discard unused patches 14 days after opening



1 Peel the patch from the inner film.



2 Place on affected area.



3 Slide the top film upward.



4 Slide the bottom film downward.

Salonpas®

Arthritis Pain Patch

**Menthol 3%
Methyl salicylate 10%
Topical Analgesic**

3 LARGE patches
3 1⁵/₁₆" X 5 1¹/₂"
(10cm X 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Hisamitsu®

Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
JAPAN SAGA TOSU
MADE IN JAPAN

Distributed by
Hisamitsu America, Inc.
Torrance, CA 90503

*Package not child resistant.
Keep out of reach of children.*

FDA APPROVED **NON-RESCRIPTION PAIN RELIEVING PATCH**
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

LOT

EXP.

® - Registered Trade Mark

CUT OPEN HERE ----- CUT OPEN HERE

FOLD TWICE AFTER OPENING
Discard unused patches 14 days after opening

3 LARGE patches
3 1⁵/₁₆" X 5 1¹/₂"
(10cm X 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Hisamitsu®

Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
JAPAN SAGA TOSU
MADE IN JAPAN

Distributed by
Hisamitsu America, Inc.
Torrance, CA 90503

*Package not child resistant.
Keep out of reach of children.*

FDA APPROVED **NON-RESCRIPTION PAIN RELIEVING PATCH**
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

LOT

EXP.

® - Registered Trade Mark

Drug Facts

Active ingredients (in each patch)

Menthol 3%
Methyl salicylate 10% (NSAID - nonsteroidal anti-inflammatory drug)

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:

- arthritis
- sprains
- strains
- bruises
- simple backache

Warnings

For external use only

Stomach bleeding warning:
This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- on the face or rashes
- with a healing pad
- any patch from a pouch that has been open for 14 or more days
- right before or after heart surgery

Ask a doctor before use if

- you are allergic to topical products
- you are taking a diuretic
- you have heart attack or stroke may increase if you use more than directed or for longer than directed
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
- if eye contact occurs, rinse thoroughly with water.
- if you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- rash, itching or skin irritation develops
- symptoms clear up and occur again within a few days

If pregnant or breast feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions

Adults 18 years and older:

- only use one patch at a time
- clean and dry affected area
- apply one patch to the affected area and leave in place for up to 8 to 12 hours
- if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- do not use more than 2 patches per day
- do not use for more than 3 days in a row
- the used patch should be removed from the skin when a new one is applied

Children under 18 years of age, do not use

Other information

- store at 20-25°C (68-77°F)
- avoid storing product in direct sunlight
- protect product from excessive moisture
- some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients alcoholic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200 000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments? Toll free 1-87-SALONPAS. Weekdays, 9AM to 5PM (PST) www.salonpas.us

1L77610

Salonpas[®] Arthritis Pain Patch **LARGE** patches
Menthol 3%, Methyl Salicylate 10%

Hisamisus[®]

LARGE

for temporary relief of mild to moderate pain

- Arthritis
- Joint Pain
- Sprains
- Strains



Salonpas

Arthritis Pain Patch

APPLY FOR
8-12
HOURS

Menthol 3%
Methyl Salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

2 LARGE patches
3 1/8" x 5 1/2" (10cm x 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Package not child resistant.
Keep out of reach of children.

LOT: _____
EXP: _____
Salonpas[®] Arthritis Pain Patch **LARGE** patches
Menthol 3%, Methyl Salicylate 10%



Salonpas[®] **2** patches
Arthritis Pain Patch **LARGE**
Menthol 3%, Methyl Salicylate 10%



Salonpas[®] Arthritis Pain Patch **LARGE** patches
Menthol 3%, Methyl Salicylate 10%
press open
minty scent

Drug Facts

Active ingredients (in each patch)
Menthol 3%,..... Topical analgesic
Methyl salicylate 10%(NSAID : nonsteroidal anti-inflammatory drug)..... Topical analgesic

Uses
Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
• arthritis • sprains • strains • bruises • simple backache

Warnings

For external use only
Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
• are age 60 or older • have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• take more or for a longer time than directed

Do not use
• on the face or rashes • on wounds or damaged skin • if allergic to aspirin or other NSAIDs
• with a heating pad • when sweating (such as from exercise or heat)
• any patch from a pouch that has been open for 14 or more days • right before or after heart surgery

Ask a doctor before use if
• you are allergic to topical products • the stomach bleeding warning applies to you
• you are taking a diuretic • you have high blood pressure, heart disease, or kidney disease

When using this product:
• wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
• the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
• you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
• rash, itching or skin irritation develops • condition worsens • symptoms last for more than 3 days
• symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions

Adults 18 years and older:
• clean and dry affected area • only use one patch at a time
• apply one patch to the affected area and leave in place for up to 12 hours • remove patch from backing film and apply to skin (see illustration)
• if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours • do not use more than 2 patches per day • do not use for more than 3 days in a row
• the used patch should be removed from the skin when a new one is applied

Other information

• store at 20-25°C (68-77°F) • avoid storing product in direct sunlight
• protect product from excessive moisture • some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients


alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene, 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments? Toll free 1-87-SALONPAS. Weekdays, 9AM to 5PM (PST) www.salonpas.us

4pk
0.4pk
Nimig 0

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

FOLD TWICE AFTER OPENING
Discard unused patches 14 days after opening
FOLD TWICE AFTER OPENING



Menthol 3%
Methyl salicylate 10%
Topical Analgesic

FDA APPROVED, NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

2 LARGE Patches
3 15/16" X 5 1/2"
(10cm X 14cm)

Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
JAPAN SAGATOSU
MADE IN JAPAN
Distributed by
Hisamitsu America, Inc.
Torrance, CA 90503
LOT
EXP.
Package not child resistant.
Keep out of reach of children.

® : Registered Trade Mark

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

Salonpas® Arthritis Pain Patch LARGE Menthol 3%, Methyl salicylate 10%, minty scent

Drug Facts

Active ingredients (in each patch)
Menthol 3%
Methyl salicylate 10% (NSAID: nonsteroidal anti-inflammatory drug)
Topical analgesic
Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
arthritis ■ sprains ■ strains ■ bruises ■ simple backache

Warnings
For external use only
Stomach bleeding warning:
This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use
■ on the face or rashes ■ on wounds or damaged skin
■ with a heating pad ■ when sweating (such as from exercise or heat)
■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

Ask a doctor before use if
■ you are allergic to topical products ■ the stomach bleeding warning applies to you
■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease

When using this product ■ wash hands after applying or removing patch. Avoid contact with eyes.
If eye contact occurs, rinse thoroughly with water.
If the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions
Adults 18 years and older:
■ only use one patch at a time
■ clean and dry affected area
■ apply one patch to the affected area and leave in place for up to 6 to 12 hours
■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row
■ the used patch should be removed from the skin when a new one is applied
Children under 18 years of age: do not use

Other information
■ avoid storing product in direct sunlight
■ store at 20-25°C (68-77°F)
■ protect product from excessive moisture
■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients alicyclic, saturated hydrocarbon, resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-sebacate-styrene block copolymer, synthetic aluminum silicate


Questions or comments?
Toll free 1-87-SALONPAS. Weekdays, 9AM to 5PM (PST) www.salonpas.us

1L7751-P0



Salonpas

Salonpas



Salonpas

Salonpas

Salonpas
Arthritis Pain Patch

APPLY FOR
8-12
HOURS

STRETCHABLE & FLEXIBLE

1 LARGE patch
3 1/4" X 5 1/2" (10cm X 14cm)

Menthol 3%
Methyl salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

1 LARGE patch X 12 sachets
3 1/4" X 5 1/2" (10cm X 14cm)

503 46567 68501 0

2pr
0.4pr
0

Hiemsuru® minty scent
NDC#4581-685-01

LARGE for temporary relief of mild to moderate pain

Arthritis
Joint Pain
Sprains
Strains

1 Peel the center film.
2 Place on affected area.
3 Slice the film upward.
4 Slice the film downward.

Salonpas®
Arthritis Pain Patch

APPLY FOR 8-12 HOURS

Menthol 3%
Methyl salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

1 LARGE patch
3.15" x 5.12" (10cm x 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Package not child resistant.
Keep out of reach of children.

MADE IN JAPAN

Hisamitsu Pharmaceutical Co., Inc.
Hisamitsu America, Inc.
4658168501

⌘ CUT OPEN HERE ----- CUT OPEN HERE ⌘

Drug Facts

Active ingredients (in each patch)

Menthol 3%
Methyl salicylate 10%
Topical analgesic
Topical analgesic

Purpose
Menthol 3%
Methyl salicylate 10%
Topical analgesic
Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
 ■ arthritis ■ sprains ■ strains ■ bruises ■ simple backache

Warnings
 For external use only
 Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The danger is small but higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Do not use
 ■ on the face or rashes
 ■ when sweating (such as from exercise or heat)
 ■ any patch from a pouch that has been open for 14 or more days
 ■ if allergic to aspirin or other NSAIDs

Ask a doctor before use if
 ■ you are allergic to topical products
 ■ the stomach bleeding warning applies to you
 ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, or kidney disease

When using this product
 ■ wash hands after applying or removing patch. Avoid contact with eyes.
 ■ if eye contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
 ■ stomach pain or upset gets worse or lasts
 ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
 ■ rash, itching or skin irritation develops
 ■ condition worsens
 ■ symptoms last for more than 3 days
 ■ symptoms clear up and occur again within a few days

Symptoms that may occur during pregnancy and during the first 6 months of pregnancy.
 Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If put in mouth, get medical help or contact Poison Control Center right away. Patchage not child resistant. Dispose of the used patches after folding sticky ends together.

Directions
 Adults 18 years and older:
 ■ only use one patch at a time
 ■ remove patch from backing film and apply to skin (see illustration)
 ■ clean and dry affected area
 ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours
 ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
 ■ do not use more than 2 patches per day
 ■ do not use for more than 3 days in a row
 ■ the used patch should be removed from the skin when a new one is applied
 Children under 18 years of age: do not use

Other information
 ■ store at 20-25°C (68-77°F)
 ■ avoid storing product in direct sunlight
 ■ protect product from excessive moisture
 ■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients
 ethylacrylate, saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
 Toll free 1-877-SALONPAS Weddays, 9AM to 5PM (PST)
 www.Salonpas.us

LOT
EXP.

© Registered Trade Mark 1L774T-0

241
0.4qt
mining: 0

Salonpas
Arthritis Pain Patch

APPLY FOR 8-12 HOURS

1 LARGE patch
3 1/4" x 5 1/2" (10cm x 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

**MENTHOL 3%
Methyl salicylate 10%
Topical Analgesic**

**FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL**

**Package not child resistant.
Keep out of reach of children.**

Manufactured by
H Santen Pharmaceutical Co., Inc.
MADE IN JAPAN

© 2013 H Santen
H Santen America, Inc.
Torrance, CA 90503

minty scent
NDC #45831-085-59

FREE SAMPLE NOT FOR RE-SALE

LARGE

For temporary relief of mild to moderate pain

Arthritis
Joint Pain
Sprains
Strains

1 Peel the center film.
2 Place on affected area.
3 Slide the film upward.
4 Slide the other film downward.

➤ CUT OPEN HERE ----- CUT OPEN HERE ➤

Drug Facts

Active ingredients (in each patch)

Menthol 3%
Methyl salicylate 10% (NSAID: nonsteroidal anti-inflammatory drug)
Topical analgesic
Topical anesthetic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
arthritis sprains strains bruises simple backache

Warnings

For external use only

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems
take a blood thinning (anticoagulant) or steroid drug
take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product.
Take more or for a longer time than directed

Do not use

- on the face or rashes
- on wounds or damaged skin
- with a heating pad
- any patch from a pouch that has been open for 14 or more days
- ask a doctor before use if you are allergic to topical products, the stomach bleeding warning applies to you
- you are taking a diuretic
- you have high blood pressure, heart disease, or kidney disease

When using this product, wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.

If the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:

- stomach pain or upset gets worse or lasts
- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- rash, itching or skin irritation develops
- condition worsens
- symptoms last for more than 3 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after holding sticky ends together.

Directions

Adults 18 years and older:

- only use one patch at a time
- clean and dry affected area
- remove patch from backing film and apply to skin (see illustration)
- apply one patch to the affected area and leave in place for up to 8 to 12 hours
- if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- do not use more than 2 patches per day
- do not use for more than 3 days in a row
- the used patch should be removed from the skin when a new one is applied

Children under 18 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid storing product in direct sunlight
- protect product from excessive moisture
- some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

salicylic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyethylene glycol, polyethylene glycol 200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?

Toll free 1-877-SALONPAS, Weekdays, 9AM to 5PM (PST)
www.salonpas.us

LOT

EXP

© : Registered Trade Mark 11.77310

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER

11/05/2012