


Labeling Format Information:

Font: Helvetica Roman, Bold, Bold Oblique	
Drug Facts: 9.5pt	Leading: 6.5pt
Header: 8pt	Bullets: 5pt
Subheader: 6pt	Barlines: 2.5pt
Body Text: 6pt	Hairlines: 0.5pt

Horizontal Scale: 80%-100% Average Kerning: 0



SARP 20ct clean

20 Salonpas Arthritis Pain Patch Patches

Menthol 3%, Methyl Salicylate 10%

Manufactured by Hisamitsu Pharmaceutical Co., Inc. 408, Tashirodai-kan-machi, Tosu, Saga, Japan

MADE IN JAPAN

Distributed by Hisamitsu America, Inc. Fortran Park, NJ 07932

® Registered Trade Mark

Salonpas® Arthritis Pain Patch 20 PATCHES

Menthol 3%, Methyl Salicylate 10%

for temporary relief of mild to moderate pain

- Arthritis
- Joint Pain
- Sprains
- Strains

Salonpas® Arthritis Pain Patch

Menthol 3% Methyl Salicylate 10% Topical Analgesic

APPLY FOR 8-12 HOURS

20 PATCHES 2 3/4" x 3 15/16" (7cm x 10cm)

⚠ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS.

LOT: [REDACTED] EXP: [REDACTED]

Salonpas® Arthritis Pain Patch 20 PATCHES

Menthol 3%, Methyl Salicylate 10%

Hisamitsu® NDC#46581-680-20

20 Salonpas Arthritis Pain Patch Patches

Menthol 3%, Methyl Salicylate 10%

HOW TO APPLY

- Peel the center film.
- Place on affected area.
- Slide the film upward.
- Slide the other film downward.

Salonpas® Arthritis Pain Patch 20 PATCHES

Menthol 3%, Methyl Salicylate 10%

1M447T-0

PRESS OPEN minty scent

Drug Facts	
Active ingredients (in each patch)	Purpose
Menthol 3%	Topical analgesic
Methyl salicylate 10% (NSAID: nonsteroidal anti-inflammatory drug)	Topical analgesic
Uses	
Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:	
<ul style="list-style-type: none"> ■ arthritis ■ sprains ■ strains ■ bruises ■ simple backache 	
Warnings	
For external use only	
<p>Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:</p> <ul style="list-style-type: none"> ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed 	
<p>Do not use</p> <ul style="list-style-type: none"> ■ on the face or rashes ■ on wounds or damaged skin ■ with a heating pad ■ when sweating (such as from exercise or heat) ■ any patch from a pouch that has been open for 14 or more days ■ if allergic to aspirin or other NSAIDs ■ right before or after heart surgery 	
<p>Ask a doctor before use if</p> <ul style="list-style-type: none"> ■ you are allergic to topical products ■ you are taking a diuretic ■ the stomach bleeding warning applies to you ■ you have high blood pressure, heart disease, or kidney disease 	
<p>When using this product</p> <ul style="list-style-type: none"> ■ wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water. ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed 	
<p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding. ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days ■ symptoms clear up and occur again within a few days 	
<p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use methyl salicylate at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.</p>	
Directions	
<p>Adults 18 years and older:</p> <ul style="list-style-type: none"> ■ clean and dry affected area ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row ■ the used patch should be removed from the skin when a new one is applied 	
<p>Children under 18 years of age: do not use; this product has not been shown to work in children</p>	
Other information	
<ul style="list-style-type: none"> ■ Store at 20-25°C (68-77°F) ■ Protect product from excessive moisture ■ Some individuals may not experience pain relief until several hours after applying the patch 	
Inactive ingredients	
<p>alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate</p>	
Questions or comments?	
<p>Toll free 1-87-SALONPAS www.salonpas.us</p>	

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
10/06/2025 02:01:57 PM