

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatf>

Rx Only **NDC 0078-0951-66**

120 capsules

 **Tasigna**[®]
(nilotinib) capsules
50 mg

Dispense with Medication Guide

TAKE TASIGNA ON EMPTY STOMACH.
AVOID FOOD FOR AT LEAST 2 HOURS
BEFORE TAKING A DOSE OF TASIGNA
AND AT LEAST 1 HOUR AFTER THE DOSE
IS TAKEN.

Each capsule contains
50 mg nilotinib.
Dosage: See Prescribing
Information.
Store at 20°C to 25°C
(68°F to 77°F),
excursions permitted
between 15°C and 30°C
(59°F and 86°F).
Keep all drugs out of
the reach of children.
Mfd by Novartis Pharma
Stein AG, Stein,
Switzerland
Dist by Novartis
Pharmaceuticals Corp.,
East Hanover, NJ 07936
Product of Ireland

3 N
0078-0951-66
1








 NOVARTIS 46292087 US


This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

 **Tasigna**[®] (nilotinib) capsules **200 mg per capsule**



**TAKE TASIGNA[®] ON EMPTY STOMACH.
PLEASE SEE PATIENT MEDICATION GUIDE FOR FULL INFORMATION.**

 **Tasigna**[®] (nilotinib) capsules **200 mg per capsule**



**TAKE TASIGNA[®] ON EMPTY STOMACH.
PLEASE SEE PATIENT MEDICATION GUIDE FOR FULL INFORMATION.**

Dosage: See Prescribing Information.

 **NOVARTIS**



46300682 US


©Novartis

2


This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



46300685 US



 **Tasigna**[®] (nilotinib) capsules **200 mg** per capsule



 **Tasigna**[®] (nilotinib) capsules **200 mg** per capsule



NOVARTIS

For oral use.
Dosage: See Prescribing Information.
Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].
1 capsule contains 200 mg nilotinib.
Keep this and all drugs out of the reach of children.
For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna

TAKE TASIGNA ON
EMPTY STOMACH.
AVOID FOOD FOR
AT LEAST 2 HOURS
BEFORE TAKING A
DOSE OF TASIGNA
AND AT LEAST
1 HOUR AFTER THE
DOSE IS TAKEN.

Rx Only

Contents:
4 individual blister packs containing
28 capsules each.

200 mg
per capsule

(nilotinib) capsules

Tasigna®


NDC 0078-0526-87

46300691 US

Contents:
4 individual blister packs containing
28 capsules each.

Tasigna®
(nilotinib) capsules
200 mg
per capsule

Manufactured by:
Novartis Pharma Stein AG, Stein, Switzerland
Distributed by:
Novartis Pharmaceuticals Corp., East Hanover, NJ 07936
Product of Ireland
For more information, visit www.US.TASIGNA.com or call
1-866-411-Tasigna
©Novartis



Tasigna®
(nilotinib) capsules
200 mg
per capsule
NOVARTIS

Contents:
4 individual blister packs containing
28 capsules each.

Rx Only
TAKE TASIGNA ON
EMPTY STOMACH.
AVOID FOOD FOR
AT LEAST 2 HOURS
BEFORE TAKING A
DOSE OF TASIGNA
AND AT LEAST
1 HOUR AFTER THE
DOSE IS TAKEN.

For oral use.
Dosage: See Prescribing Information.
Store at 20°C to 25°C (68°F to 77°F);
excursions permitted between
15°C and 30°C (59°F and 86°F) [see
USP Controlled Room Temperature].
1 capsule contains 200 mg nilotinib
Keep this and all drugs out of the reach of children.
For more information, visit www.US.TASIGNA.com
or call 1-866-411-Tasigna

NDC 0078-0526-87

Tasigna®
(nilotinib) capsules
200 mg
per capsule

DISPENSE WITH MEDICATION GUIDE
ATTACHED OR PROVIDED SEPARATELY.

Tasigna®
(nilotinib) capsules
200 mg
per capsule
NOVARTIS

205x82x89_E

Rx Only

PHYSICIAN SAMPLE - NOT FOR SALE

28 capsules

TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.

For oral use.
Dosage: See Prescribing Information.
Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].
1 capsule contains 200 mg nilotinib
Keep this and all drugs out of the reach of children.
For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna

Tasigna[®]
(nilotinib) capsules
200 mg
per capsule

DISPENSE WITH MEDICATION GUIDE ATTACHED OR PROVIDED SEPARATELY.

OPEN HERE


INSTRUCTIONS FOR USE

To open kit:

1. Firmly press and hold button **1** below. While holding this button, pull tab **2** below.
2. Once the card is fully out, unfold it completely.
3. To release the capsule, press gently on the capsule from the clear side of the card and push it through the paper backing.
4. Remove any loose paper backing from the back of the card.

Tasigna[®]
(nilotinib) capsules
200 mg
per capsule

To open kit:



1 Press firmly

While pressing firmly on **1**, pull on tab **2**. → Pull

To close kit:

1. The card should not be damaged, it should fit flat in the box.
2. Fold the card back. Push the folded card back into the box.
3. Fold the cover-flap over to complete closure.
4. Store in a safe place out of the reach of children.

Tasigna[®] (nilotinib) capsules 200 mg per capsule

Tasigna[®] (nilotinib) capsules 200 mg per capsule ©Novartis 0078-9526-90

PHYSICIAN SAMPLE - NOT FOR SALE

28 capsules

Tasigna[®]
(nilotinib) capsules
200 mg
per capsule


NOVARTIS

For oral use.
Dosage: See Prescribing Information.
Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].
1 capsule contains 200 mg nilotinib
Keep this and all drugs out of the reach of children.
For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna

Manufactured by:
Novartis Pharma Stein AG
Stein, Switzerland
Distributed by:
Novartis Pharmaceuticals Corp.
East Hanover, NJ 07936
Product of Ireland

46300689 US

Tasigna[®] (nilotinib) capsules 200 mg per capsule




Rx Only NDC 0078-0526-51

28 capsules

TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.

For oral use.
Dosage: See Prescribing Information.
 Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].
 1 capsule contains 200 mg nilotinib
 Keep this and all drugs out of the reach of children.
 For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna



Tasigna[®]
 (nilotinib) capsules
200 mg
 per capsule


DISPENSE WITH MEDICATION GUIDE ATTACHED OR PROVIDED SEPARATELY.

OPEN HERE ↓

INSTRUCTIONS FOR USE

To open kit:


1. Firmly press and hold button **1** below. While holding this button, pull tab **2** below.
2. Once the card is fully out, unfold it completely.
3. To release the capsule, press gently on the capsule from the clear side of the card and push it through the paper backing.
4. Remove any loose paper backing from the back of the card.



Tasigna[®]
 (nilotinib) capsules
200 mg
 per capsule

Tasigna[®] (nilotinib) capsules 200 mg

To open kit:



While pressing firmly on **1**, pull on tab **2**. → Pull

To close kit:

1. The card should not be damaged, it should fit flat in the box.
2. Fold the card back. Push the folded card back into the box.
3. Fold the cover-flap over to complete closure.
4. Store in a safe place out of the reach of children.

Tasigna[®] (nilotinib) capsules 200 mg per capsule

©Novartis



28 capsules



N 0078-0526-51 7

46300688 US

Manufactured by:
 Novartis Pharma Stein AG
 Stein, Switzerland
 Distributed by:
 Novartis Pharmaceuticals Corp.
 East Hanover, NJ 07936

For oral use.
Dosage: See Prescribing Information.
 Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].
 1 capsule contains 200 mg nilotinib
 Keep this and all drugs out of the reach of children.
 For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna

NDC 0078-0526-51




Tasigna[®]
 (nilotinib) capsules
200 mg
 per capsule

NOVARTIS

Tasigna[®] (nilotinib) capsules 200 mg per capsule




 **Tasigna**[®] (nilotinib) capsules 200 mg per capsule



PHYSICIAN SAMPLE - NOT FOR SALE

TAKE TASIGNA[®] ON EMPTY STOMACH.
PLEASE SEE PATIENT MEDICATION GUIDE FOR FULL INFORMATION.

 **Tasigna**[®] (nilotinib) capsules 200 mg per capsule



PHYSICIAN SAMPLE - NOT FOR SALE

2

Dosage: See Prescribing Information.

 NOVARTIS




46300683 US

©Novartis

This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>

 **Tasigna**[®] (nilotinib) capsules


150 mg per capsule



TAKE TASIGNA[®] ON EMPTY STOMACH.
PLEASE SEE PATIENT MEDICATION GUIDE FOR FULL INFORMATION.

 **Tasigna**[®] (nilotinib) capsules

150 mg per capsule



TAKE TASIGNA[®] ON EMPTY STOMACH.
PLEASE SEE PATIENT MEDICATION GUIDE FOR FULL INFORMATION.

Dosage: See Prescribing Information.

 NOVARTIS



46300680 US


©Novartis

2


This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



46300684 US



 **Tasigna**[®] (nilotinib) capsules **150 mg** per capsule



 **Tasigna**[®] (nilotinib) capsules **150 mg** per capsule

Rx Only

PHYSICIAN SAMPLE – NOT FOR SALE

Tasigna[®]
(nilotinib) capsules

150 mg
per capsule

28 capsules

TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.

For oral use.
Dosage: See Prescribing Information.
Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].
1 capsule contains 150 mg nilotinib
Keep this and all drugs out of the reach of children.
For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna

OPEN HERE

DISPENSE WITH MEDICATION GUIDE ATTACHED OR PROVIDED SEPARATELY.

INSTRUCTIONS FOR USE

Tasigna[®]
(nilotinib) capsules
150 mg
per capsule

To open kit:

1. Firmly press and hold button **1** below. While holding this button, pull tab **2** below.
2. Once the card is fully out, unfold it completely.
3. To release the capsule, press gently on the capsule from the clear side of the card and push it through the paper backing.
4. Remove any loose paper backing from the back of the card.

To open kit:



While pressing firmly on **1**, pull on tab **2**. → Pull


To close kit:

1. The card should not be damaged, it should fit flat in the box.
2. Fold the card back. Push the folded card back into the box.
3. Fold the cover-flap over to complete closure.
4. Store in a safe place out of the reach of children.

Tasigna[®] (nilotinib) capsules 150 mg

Tasigna[®] (nilotinib) capsules **150 mg** per capsule

©Novartis 0078-9592-90



28 capsules

Rx Only

Manufactured by:
Novartis Pharma Stein AG
Stein, Switzerland
Distributed by:
Novartis Pharmaceuticals Corp.
East Hanover, NJ 07936
Product of Ireland



46300687 US

PHYSICIAN SAMPLE – NOT FOR SALE

Tasigna[®]
(nilotinib) capsules

150 mg
per capsule

NOVARTIS

Tasigna[®] (nilotinib) capsules **150 mg** per capsule



Rx Only

NDC 0078-0592-51

Tasigna[®]
(nilotinib) capsules

150 mg
per capsule

28 capsules

TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.

For oral use.

Dosage: See Prescribing Information.

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].

1 capsule contains 150 mg nilotinib

Keep this and all drugs out of the reach of children.

For more information,

visit www.US.TASIGNA.com or call 1-866-411-Tasigna

OPEN HERE

DISPENSE WITH MEDICATION GUIDE ATTACHED OR PROVIDED SEPARATELY.

INSTRUCTIONS FOR USE

Tasigna[®]
(nilotinib) capsules

150 mg
per capsule

To open kit:

1. Firmly press and hold button **1** below. While holding this button, pull tab **2** below.
2. Once the card is fully out, unfold it completely.
3. To release the capsule, press gently on the capsule from the clear side of the card and push it through the paper backing.
4. Remove any loose paper backing from the back of the card.

To open kit:

1

Press firmly

While pressing firmly on **1**, pull on tab **2**. → Pull


To close kit:

1. The card should not be damaged, it should fit flat in the box.
2. Fold the card back. Push the folded card back into the box.
3. Fold the cover-flap over to complete closure.
4. Store in a safe place out of the reach of children.

Tasigna[®] (nilotinib) capsules **150 mg** per capsule

©Novartis

NDC 0078-0592-51



28 capsules

Rx Only



Manufactured by:
Novartis Pharma Stein AG
Stein, Switzerland
Distributed by:
Novartis Pharmaceuticals Corp.
East Hanover, NJ 07936

46300686x US

For oral use.

Dosage: See Prescribing Information.

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].

1 capsule contains 150 mg nilotinib

Keep this and all drugs out of the reach of children.

For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna


Tasigna[®]
(nilotinib) capsules

150 mg
per capsule

NOVARTIS

 **Tasigna**[®] (nilotinib) capsules

150 mg per capsule




PHYSICIAN SAMPLE - NOT FOR SALE

TAKE TASIGNA[®] ON EMPTY STOMACH.
PLEASE SEE PATIENT MEDICATION GUIDE FOR FULL INFORMATION.

 **Tasigna**[®] (nilotinib) capsules

150 mg per capsule



PHYSICIAN SAMPLE - NOT FOR SALE

Dosage: See Prescribing Information.

 **NOVARTIS**



46300681 US

©Novartis

2



NOVARTIS

For oral use.
Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].
1 capsule contains 150 mg nilotinib.
Keep this and all drugs out of the reach of children.
For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna

Rx Only
TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.

150 mg
per capsule

(nilotinib) capsules

Tasigna®

Contents:
4 individual blister packs containing 28 capsules each.

NDC 0078-0592-87

46300690 US

Contents:
4 individual blister packs containing 28 capsules each.




0078-0592-87

Tasigna®
(nilotinib) capsules

150 mg
per capsule

Manufactured by:
Novartis Pharma Stein AG, Stein, Switzerland
Distributed by:
Novartis Pharmaceuticals Corp., East Hanover, NJ 07936
Product of Ireland
For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna
©Novartis



Tasigna®
(nilotinib) capsules

150 mg
per capsule

NOVARTIS

Contents:
4 individual blister packs containing 28 capsules each.

Rx Only
TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.

For oral use.
Dosage: See Prescribing Information.
Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].
1 capsule contains 150 mg nilotinib.
Keep this and all drugs out of the reach of children.
For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna

NDC 0078-0592-87

Tasigna®
(nilotinib) capsules

150 mg
per capsule

DISPENSE WITH MEDICATION GUIDE ATTACHED OR PROVIDED SEPARATELY.

Tasigna®
(nilotinib) capsules

150 mg
per capsule

NOVARTIS

205x82x89_E