

NDA 022068/S-040
Page 3

Rx Only **NDC 0078-0951-66**
120 capsules

Tasigna[®]
(nilotinib) capsules
50 mg

Dispense with Medication Guide

TAKE TASIGNA ON EMPTY STOMACH.
AVOID FOOD FOR AT LEAST 2 HOURS
BEFORE TAKING A DOSE OF TASIGNA
AND AT LEAST 1 HOUR AFTER THE DOSE
IS TAKEN.

Each capsule contains 50 mg nilotinib.
Dosage: See Prescribing Information.
Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (59°F and 86°F).
Keep all drugs out of the reach of children.
Mfd. by: Novartis d.o.o. Ljubljana, Slovenia
Dist. by: Novartis Pharmaceuticals Corp., East Hanover, NJ 07936
Product of Ireland

NOVARTIS 300070877 US


Contents:
4 individual blister packs containing
28 capsules each.

Tasigna[®]
(nilotinib) capsules
150 mg
per capsule

Manufactured by:
Novartis d.o.o.
Ljubljana, Slovenia

Distributed by:
Novartis Pharmaceuticals Corp., East Hanover, NJ 07936
Product of Ireland

For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna
©Novartis




NDA 022068/S-040

Page 4

Contents:
4 individual blister packs containing
28 capsules each.

 **Tasigna**[®]
(nilotinib) capsules
200 mg
per capsule

Manufactured by:
Novartis d.o.o.
Ljubljana, Slovenia
Distributed by:
Novartis Pharmaceuticals Corp., East Hanover, NJ 07936
Product of Ireland
For more information, visit www.US.TASIGNA.com or call
1-866-411-Tasigna
©Novartis



This label may not be the latest approved by FDA.
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Ramesh
Raghavachari

Digitally signed by Ramesh Raghavachari
Date: 2/29/2024 12:15:20PM
GUID: 502d0913000029f375128b0de8c50020