

NDA 022068/S-039

Page 3

Carton and Container Labeling

**Rx Only**    **NDC 0078-0951-66**    Each capsule contains 50 mg nilotinib.  
120 capsules    Dosage: See Prescribing Information.

**Tasigna**<sup>®</sup>  
(nilotinib) capsules  
**50 mg**

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (59°F and 86°F). Keep all drugs out of the reach of children.

**Dispense with Medication Guide**

**TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.**



Mfd. by: Novartis Pharma Stein AG, Stein, Switzerland  
Dist. by: Novartis Pharmaceuticals Corp., East Hanover, NJ 07936  
Product of China

**NOVARTIS**    300069784 US

3 N  
0078-0951-66  
1

300069784 US


NDA 022068/S-039  
Page 4

<b>PHYSICIAN SAMPLE – NOT FOR SALE</b>			<b>Rx Only</b>
<b>28 capsules</b>	<p>For oral use. Dosage: See Prescribing Information. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature]. 1 capsule contains 150 mg nilotinib. Keep this and all drugs out of the reach of children. For more information, visit <a href="http://www.US.TASIGNA.com">www.US.TASIGNA.com</a> or call 1-866-411-Tasigna</p>		<b>DISPENSE WITH MEDICATION GUIDE ATTACHED OR PROVIDED SEPARATELY.</b>
<b>TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.</b>	<b>OPEN HERE</b> ▼		
<b>INSTRUCTIONS FOR USE</b>			
<b>To open kit:</b>			
<ol style="list-style-type: none"><li>1. Firmly press and hold button <b>1</b> below. While holding this button, pull tab <b>2</b> below.</li><li>2. Once the card is fully out, unfold it completely.</li><li>3. To release the capsule, press gently on the capsule from the clear side of the card and push it through the paper backing.</li><li>4. Remove any loose paper backing from the back of the card.</li></ol>			

NDA 022068/S-039  
Page 5



NDA 022068/S-039  
 Page 6



Rx Only

**PHYSICIAN SAMPLE - NOT FOR SALE**

**28 capsules**

**TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.**

**For oral use.**  
**Dosage:** See Prescribing Information. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].  
 1 capsule contains 200 mg nilotinib  
 Keep this and all drugs out of the reach of children.  
 For more information, visit [www.US.TASIGNA.com](http://www.US.TASIGNA.com) or call 1-866-411-Tasigna

**OPEN HERE**

**Tasigna®**  
 (nilotinib) capsules  
**200 mg**  
 per capsule

**DISPENSE WITH MEDICATION GUIDE ATTACHED OR PROVIDED SEPARATELY.**

**INSTRUCTIONS FOR USE**

**To open kit:**

1. Firmly press and hold button **1** below. While holding this button, pull tab **2** below.
2. Once the card is fully out, unfold it completely.
3. To release the capsule, press gently on the capsule from the clear side of the card and push it through the paper backing.
4. Remove any loose paper backing from the back of the card.

**Tasigna®**  
 (nilotinib) capsules  
**200 mg**  
 per capsule

**To open kit:**



While pressing firmly on **1**, pull on tab **2**. → Pull

**To close kit:**

1. The card should not be damaged, it should fit flat in the box.
2. Fold the card back. Push the folded card back into the box.
3. Fold the cover-flap over to complete closure.
4. Store in a safe place out of the reach of children.

---

**Tasigna® (nilotinib) capsules 200 mg per capsule** ©Novartis 0078-9526-90



**28 capsules**

Manufactured by:  
Novartis Pharma Stein AG  
Stein, Switzerland  
Distributed by:  
Novartis Pharmaceuticals Corp.  
East Hanover, NJ 07936  
Product of China

300069778 US

**PHYSICIAN SAMPLE - NOT FOR SALE**



**Tasigna®**  
(nilotinib) capsules  
**200 mg**  
per capsule



For oral use.  
Dosage: See Prescribing Information.  
Store at 20°C to 25°C (68°F to 77°F);  
excursions permitted between  
15°C and 30°C (59°F and 86°F) [see  
USP Controlled Room Temperature].

1 capsule contains 200 mg nilotinib  
Keep this and all drugs out of the  
reach of children.

For more information, visit  
[www.US.TASIGNA.com](http://www.US.TASIGNA.com) or call  
1-866-411-Tasigna

---


**Tasigna® (nilotinib) capsules 200 mg per capsule**

NOVARTIS

**Tasigna® (nilotinib) capsules 200 mg per capsule**

NDC 0078-0526-97

Contents: 4 individual blister packs containing 28 capsules each.




3 078-0526-87

Manufactured by:  
Novartis Pharma Stein AG, Stein, Switzerland  
Distributed by:  
Novartis Pharmaceuticals Corp., East Hanover, NJ 07936  
Product of China  
For more information, visit [www.US.TASIGNA.com](http://www.US.TASIGNA.com) or call  
1-866-411-Tasigna  
©Novartis

200069777 US

**Tasigna® (nilotinib) capsules 200 mg per capsule**



NOVARTIS

**Tasigna® (nilotinib) capsules 200 mg per capsule**

NDC 0078-0526-97


Contents: 4 individual blister packs containing 28 capsules each.

**Ex. Only**

**TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.**

For oral use.  
Dosage: See Prescribing Information.  
Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see USP Controlled Room Temperature].  
1 capsule contains 200 mg nilotinib.  
Keep this and all drugs out of the reach of children.  
For more information, visit [www.US.TASIGNA.com](http://www.US.TASIGNA.com) or call 1-866-411-Tasigna

**DISPENSE WITH MEDICATION GUIDE ATTACHED OR PROVIDED SEPARATELY.**



This label may not be the latest approved by FDA.  
For current labeling information, please visit <https://www.fda.gov/drugsatfda>



Rohit  
Kolhatkar

Digitally signed by Rohit Kolhatkar  
Date: 2/06/2024 03:01:16PM  
GUID: 57bf531500b52a2eccd5395bec77ebc2