

UNPRINTED AREA FOR
LOT & EXP CODING
(UNVARNISHED)

Pfizer, Madison, NJ 07640 USA © 2019 Pfizer Inc.
For most recent product information, visit www.advil.com

3 0573 0190 40 4

81% For Position Only

Drug Facts

Active ingredients (in each tablet)	Purposes
Chlorpheniramine maleate 4 mg	Antihistamine
Ibuprofen 200 mg (NSAID)*	Pain reliever/Fever reducer
Phenylephrine HCl 10 mg	Nasal decongestant
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves these symptoms associated with the common cold or flu:

headache	fever	minor body aches and pains
nasal congestion	runny nose	sinus pressure
		sneezing

PAA124485.FDA01

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- in children under 12 years of age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

- you have a breathing problem such as emphysema or chronic bronchitis
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke

Drug Facts (continued)

- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking sedatives or tranquilizers
- taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- may cause excitability especially in children

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
 - pain gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days

Drug Facts (continued)

Inactive ingredients: acacia, croscarmellose sodium, glycerol dibehenate, hypromellose, lactic acid, lactin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polyethylene glycol, pregelatinized starch, propyl gallate, silicon dioxide, sucrose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

Questions or comments? Call weekdays from 9 AM to 5 PM EST toll free at 1-800-98-ADVIL

Drug Facts (continued)

- nasal congestion lasts for more than 7 days
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- symptoms continue or get worse
- any new symptoms appear

if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- adults and children 12 years of age and over:
 - take 1 tablet every 4 hours while symptoms persist
 - do not take more than 6 tablets in any 24-hour period, unless directed by a doctor
- children under 12 years of age: do not use

Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

PAA124485.FDA01

128 CODE = 124485

FPO

Advil®

MULTI-SYMPTOM
COLD & FLU

Ibuprofen 200 mg Pain Reliever/Fever Reducer (NSAID)
Phenylephrine HCl 10 mg Nasal Decongestant
Chlorpheniramine Maleate 4 mg Antihistamine

✓ Headache	✓ Nasal Congestion
✓ Fever	✓ Runny Nose
✓ Body Aches	✓ Sneezing
✓ Sinus Pressure	✓ Nasal Swelling

ONE
TABLET
DOSE!

40 Coated
Tablets

128 CODE = 124485

FPO

090177e1911929f8\Final\Final On: 13-Jun-2019 18:29 (GMT)

FDA SUBMISSION (NOT COLOR ACCURATE)

Job Information	Color Legend
Date 05-MAR-2019 PASS # 1 Component # PAA124485.FDA01 Drawing # DWG-50054091 v01	(b) (4)
Identification	Printer & User Notes
	PRINTER TO REPLACE ALL FPO (For Position Only) CODES WITH SCANNABLE ART.
(b) (4)	PRINTER RESPONSIBLE FOR ALL TRAPPING, BLEEDS, COATINGS AND RESOLUTION.
Graphic Artist: (b) (4)	PRINTER AND USERS RESPONSIBLE FOR PROPER LICENSING OF ALL FONTS USED IN THIS ARTWORK.

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

SILVER FOIL -

Reference ID: 4487775

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

THERESA M MICHELE
09/06/2019 12:12:51 PM