

BACK

AREA FOR LOT AND EXP
TO BE PRINTED IN DARK COLOR
26 mm x 5.5 mm

taking pain relievers or fever reducers, you have a history of stomach problems, such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke; you are taking a diuretic. **Ask a doctor or pharmacist before use** if you are under a doctor's care for any serious condition, taking sedatives or tranquilizers, taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or an histamine, taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin or taking any other drug. **When using this product** take with food or milk, stomach upset occurs, avoid alcohol; drinks, be careful when driving a motor vehicle or operating machinery, drowsiness may occur, alcohol, sedatives, and tranquilizers may increase drowsiness, may cause excitability especially in children. **Stop use and ask a doctor** if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools or have stomach pain that does not get better; you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, or leg swelling; pain gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; nasal congestion lasts for more than 7 days; redness or swelling is present in the painful area; you get nervous, dizzy, or sleepless; symptoms continue or get worse; any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** Do not take more than directed. Adults and children 12 years of age and over: take 1 tablet every 4 hours while symptoms persist. Do not take more than 6 tablets in any 24-hour period unless directed by a doctor. Children under 12 years of age: do not use because this product contains too much medication for children under this age. **Other information** Read all warnings and directions before use. Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). **Active ingredients:** aceclofenac, potassium, artificial flavors, carnauba wax, coloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactic acid, lecithin, malbolondrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polyethylene, polyvinyl alcohol, pregelatinized starch, pronyl gelatin, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum. **Call toll free at 1-800-88-ADWILL** For most recent product information, visit www.adwill.com GSK CH, Warren, NJ 07059 ©2021 GSK or its affiliates

000065788

Advil ALLERGY & CONGESTION RELIEF

Chlorpheniramine Maleate 4 mg Antihistamine
 Ibuprofen 200 mg (NSAID)..... Pain reliever/Fever reducer
 Phenylephrine HCl 10 mg Nasal decongestant

1 Tablet *nonsteroidal anti-inflammatory drug

Uses temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies and the common cold: runny nose, itchy, watery eyes, itching of the nose or throat, sneezing, nasal congestion, sinus pressure, headache, minor body aches and pains, fever, reduces swelling of the nasal passages; temporarily restores freer breathing through the nose.

Warnings
Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. In an allergic reaction occur, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** in children under 12 years of age because this product contains too much medication for children under this age, if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery, if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. **Ask a doctor before use** if you have a breathing problem such as emphysema or chronic bronchitis, stomach bleeding, warning applies to you, you have problems or serious side effects from taking pain relievers or fever reducers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke; you are taking a diuretic. **Ask a doctor or pharmacist before use** if you are under a doctor's care for any serious condition, taking sedatives or tranquilizers, taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine, taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin or taking any other drug. **When using this product** take with food or milk. If stomach upset occurs, avoid alcoholic drinks, be careful when driving a motor vehicle or operating machinery. Drowsiness may occur, alcohol, sedatives, and tranquilizers may increase drowsiness, may cause excessive drowsiness in children. **Stop use and ask a doctor** if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools or have stomach pain that does not get better, you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, or leg swelling; pain gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; nasal congestion lasts for more than 7 days; redness or swelling is present in the painful area; you get nervous, dizzy, or sleepless; symptoms continue or get worse; any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions:** Do not take more than directed. Adults and children 12 years of age and over: take 1 tablet every 4 hours while symptoms persist. Do not take more than 6 tablets in any 24-hour period unless directed by a doctor. Children under 12 years of age: do not use because this product contains too much medication for children under this age. **Other information:** Read all warnings and directions before use. Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). **Inactive ingredients:** acacia, fumed potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactac acid, lecithin, magnesium stearate, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucrose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum. **Call toll free at 1-800-88-ADVIL.** For most recent product information, visit www.advil.com. Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensor 000065791

TO OPEN, FOLD BACK ON DOTTED LINE
 TEAR ON ARROW OR USE SCISSORS.
 Do not use if pouch is broken or torn.

FRONT

effects from taking pain relievers or fever reducers; you have a history of stomach problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke; you are taking a diuretic. **Ask a doctor or pharmacist before use** if you are under a doctor's care for any serious condition, taking sedatives or tranquilizers, taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine, taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin or taking any other drug. **When using this product** take with food or milk. If stomach upset occurs, avoid alcoholic drinks, be careful when driving a motor vehicle or operating machinery. Drowsiness may occur, alcohol, sedatives, and tranquilizers may increase drowsiness, may cause excessive drowsiness in children. **Stop use and ask a doctor** if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools or have stomach pain that does not get better, you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, or leg swelling; pain gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; nasal congestion lasts for more than 7 days; redness or swelling is present in the painful area; you get nervous, dizzy, or sleepless; symptoms continue or get worse; any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions:** Do not take more than directed. Adults and children 12 years of age and over: take 1 tablet every 4 hours while symptoms persist. Do not take more than 6 tablets in any 24-hour period unless directed by a doctor. Children under 12 years of age: do not use because this product contains too much medication for children under this age. **Other information:** Read all warnings and directions before use. Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). **Inactive ingredients:** acacia, fumed potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactac acid, lecithin, magnesium stearate, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucrose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum. **Call toll free at 1-800-88-ADVIL.** For most recent product information, visit www.advil.com. Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensor 000065791

BACK

LASER ENCODING AREA FOR LOT AND EXPIRATION INFORMATION. PRINTED IN ONE SOLID COLOR, BLACK OR DARK BLUE.

BACK

000065791

PANEL
1

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

LIFT HERE
For More Drug Facts

<p>Drug Facts</p> <p>Active ingredients (in each tablet)</p> <p>Chlorpheniramine maleate 4 mgAntihistamine Ibuprofen 200 mg (NSAID)* Pain reliever/Fever reducer Phenylephrine HCl 10 mgNasal decongestant *nonsteroidal anti-inflammatory drug</p> <p>Purposes</p> <p>temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:</p> <ul style="list-style-type: none"> ■ runny nose ■ itchy, watery eyes ■ itching of the nose or throat ■ sneezing ■ nasal congestion ■ sinus pressure ■ headache ■ minor body aches and pains ■ fever <p>reduces swelling of the nasal passages temporarily restores freer breathing through the nose</p>	<p>Drug Facts (continued)</p> <p>Warnings</p> <p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> ■ hives ■ asthma (wheezing) ■ skin reddening ■ blisters ■ facial swelling ■ shock ■ rash <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
---	--

000065956

PANEL
2

<p>Drug Facts (continued)</p> <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ in children under 12 years of age because this product contains too much medication for children under this age ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> ■ you have a breathing problem such as emphysema or chronic bronchitis ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke ■ you are taking a diuretic <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ under a doctor's care for any serious condition ■ taking sedatives or tranquilizers ■ taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug <p>When using this product</p> <ul style="list-style-type: none"> ■ take with food or milk if stomach upset occurs ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ may cause excitability especially in children
--	---

BASE
PANEL

<p>Drug Facts (continued)</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: <ul style="list-style-type: none"> ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ nasal congestion lasts for more than 7 days ■ redness or swelling is present in the painful area ■ you get nervous, dizzy, or sleepless ■ symptoms continue or get worse ■ any new symptoms appear <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed ■ adults and children 12 years of age and over: <ul style="list-style-type: none"> ■ take 1 tablet every 4 hours while symptoms persist ■ do not take more than 6 tablets in any 24-hour period, unless directed by a doctor ■ children under 12 years of age: do not use because this product contains too much medication for children under this age <p>Other information</p> <ul style="list-style-type: none"> ■ read all warnings and directions before use. Keep carton. ■ store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F) <p>Inactive ingredients acesulfame potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum</p> <p>Questions or comments? Call weekdays from 9 AM to 5 PM EST toll free at 1-800-88-ADVIL</p>
---	--

000065956

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

LIFT HERE

Drug Facts

Active ingredients (in each tablet) **Purposes**

Chlorpheniramine maleate 4 mg Antihistamine
Ibuprofen 200 mg (NSAID)* Pain reliever/Fever reducer
Phenylephrine HCl 10 mg Nasal decongestant
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
 - runny nose
 - itchy, watery eyes
 - itching of the nose or throat
 - sneezing
 - nasal congestion
 - sinus pressure
 - headache
 - minor body aches and pains
 - fever
- reduces swelling of the nasal passages
- temporarily restores freer breathing through the nose

000065955

PANEL 1

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ facial swelling
- asthma (wheezing)
- shock ■ skin reddening
- rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- in children under 12 years of age because this product contains too much medication for children under this age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer

PANEL 2

Drug Facts (continued)

- right before or after heart surgery
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

- you have a breathing problem such as emphysema or chronic bronchitis
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn

Drug Facts (continued)

- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking sedatives or tranquilizers
- taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

PANEL 3

Drug Facts (continued)

When using this product

- take with food or milk if stomach upset occurs
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- may cause excitability especially in children

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - you have symptoms of heart problems or stroke:

Drug Facts (continued)

- chest pain
- trouble breathing
- weakness in one part or side of body ■ slurred speech
- leg swelling
- pain gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- symptoms continue or get worse
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child

PANEL 4

Drug Facts (continued)

or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- adults and children 12 years of age and over:
 - take 1 tablet every 4 hours while symptoms persist
 - do not take more than 6 tablets in any 24-hour period, unless directed by a doctor
- children under 12 years of age: do not use because this product contains too much medication for children under this age

Other information

- read all warnings and directions before use. Keep carton.

Drug Facts (continued)

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)


Inactive ingredients

acesulfame potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

Questions or comments?

Call weekdays from 9 AM to 5 PM EST toll free at 1-800-88-ADVIL

BASE PANEL



000065955

BACK

problems, such as heartburn, you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke; you are taking a diuretic. **Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition, taking sedatives or tranquilizers, taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine, taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin or taking any other drug.** **When using this product** take with food or milk if stomach upset occurs, avoid alcoholic drinks, be careful when driving a motor vehicle or operating machinery, drowsiness may occur, alcohol, sedatives, and tranquilizers may increase drowsiness, may cause excitability especially in children. **Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:** feel faint, vomit blood, have bloody or black stools or have stomach pain that does not get better; you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, or leg swelling; pain gets worse or lasts more than 7 days; fever gets worse or lasts more than 5 days; nasal congestion lasts for more than 7 days; redness or swelling is present in the painful area; you get nervous, dizzy, or sleepless; symptoms continue or get worse; any new symptoms appear. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions Do not take more than directed.** Adults and children 12 years of age and over: take 1 tablet every 4 hours while symptoms persist. Do not take more than 6 tablets in any 24-hour period unless directed by a doctor. Children under 12 years of age: do not use. **Other information** Read all warnings and directions before use. Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). **Inactive ingredients:** croscarmellose sodium, potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, vanillin gum

☎ toll free at 1-800-88-ADV L GSK CH, Warren, NJ 07059

00065877

AREA FOR LOT AND EXP
TO BE PRINTED IN DARK COLOR
26 mm x 5.5 mm

00065877

Advil **MULTI-SYMP TOM COLD & FLU**

Ibuprofen 200 mg (NSAID).....Pain Reliever/Fever Reducer
 Phenylephrine HCl 10 mg.....Nasal Decongestant
 Chlorpheniramine Maleate 4 mg.....Antihistamine
1 Tablet.....Nonsteroidal anti-inflammatory drug

Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Do not use** in children under 12 years of age, if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery, if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. **Ask a doctor before use** if you have a breathing problem such as emphysema or chronic bronchitis; stomach bleeding warning applies to you; you have problems or serious side effects from taking pain relievers or fever reducers; you have a history of stomach

Uses temporarily relieves these symptoms associated with the common cold or flu: headache, fever, minor body aches and pains, nasal congestion, runny nose, sinus pressure, sneezing, reduces swelling of the nasal passages; temporarily restores free breathing through the nose

TEAR OPEN ABOVE OR USE SCISSORS. DO NOT USE IF POUCH IS DAMAGED OR TORN.

UPPER END OF BACK OF CONTAINER LINE

FRONT

problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke; you are taking a diuretic. **Ask a doctor or pharmacist before use** if you are under a doctor's care for any serious condition, taking sedatives or tranquilizers, taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine, taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin or taking any other drug. **When using this product** take with food or milk if stomach upset occurs, avoid alcoholic drinks, be careful when driving a motor vehicle or operating machinery, drowsiness may occur, alcohol, sedatives, and tranquilizers may increase drowsiness, may cause excitability especially in children. **Stop use and ask a doctor** if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools or have stomach pain that does not get better; you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, or leg swelling; pain gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; nasal congestion lasts for more than 7 days; redness or swelling is present in the painful area; you get nervous, dizzy, or sleepless; symptoms continue or get worse; any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions Do not take more than directed.** Adults and children 12 years of age and over: take 1 tablet every 4 hours while symptoms persist. Do not take more than 6 tablets in any 24-hour period unless directed by a doctor. Children under 12 years of age: do not use. **Other information** Read all warnings and directions before use. Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). **Inactive ingredients** acacia, acesulfame potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

© 2021 GSK or licensee 000065886

Dist. by: GSK, CH, Warren, NJ 07059

problems, such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke; you are taking a diuretic. **Ask a doctor or pharmacist before use** if you are under a doctor's care for any serious condition, taking sedatives or tranquilizers, taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine, taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin or taking any other drug. **When using this product** take with food or milk if stomach upset occurs, avoid alcoholic drinks, be careful when driving a motor vehicle or operating machinery, drowsiness may occur, alcohol, sedatives, and tranquilizers may increase drowsiness, may cause excitability especially in children. **Stop use and ask a doctor** if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools or have stomach pain that does not get better; you have symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, or leg swelling; pain gets worse or lasts more than 7 days; fever gets worse or lasts more than 3 days; nasal congestion lasts for more than 7 days; redness or swelling is present in the painful area; you get nervous, dizzy, or sleepless; symptoms continue or get worse; any new symptoms appear. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions Do not take more than directed.** Adults and children 12 years of age and over: take 1 tablet every 4 hours while symptoms persist. Do not take more than 6 tablets in any 24-hour period unless directed by a doctor. Children under 12 years of age: do not use. **Other information** Read all warnings and directions before use. Store at 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F). **Inactive ingredients** acacia, acesulfame potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

© 2021 GSK or licensee 000065886

Dist. by: GSK, CH, Warren, NJ 07059

BACK

LASER ENCODING AREA FOR LOT AND EXPIRATION INFORMATION. PRINTED IN ONE SOLID COLOR, BLACK OR DARK BLUE.

BACK

000065886

PANEL
1

000065785

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

LIFT HERE
For More Drug Facts

<p>Drug Facts</p> <p>Active ingredients (in each tablet) Purposes</p> <p>Chlorpheniramine maleate 4 mg Antihistamine Ibuprofen 200 mg (NSAID)* Pain reliever/Fever reducer Phenylephrine HCl 10 mg Nasal decongestant *nonsteroidal anti-inflammatory drug</p> <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves these symptoms associated with the common cold or flu: <ul style="list-style-type: none"> headache fever minor body aches and pains nasal congestion runny nose sinus pressure sneezing reduces swelling of the nasal passages temporarily restores freer breathing through the nose <p>Warnings</p> <p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> hives facial swelling asthma (wheezing) shock skin reddening rash blisters <p>If an allergic reaction occurs, stop use and seek medical help right away.</p>	<p>Drug Facts (continued)</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p>
--	--

PANEL
2

<p>Drug Facts (continued)</p> <p>Do not use</p> <ul style="list-style-type: none"> in children under 12 years of age if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if</p> <ul style="list-style-type: none"> you have a breathing problem such as emphysema or chronic bronchitis stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke you are taking a diuretic 	<p>Drug Facts (continued)</p> <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> under a doctor's care for any serious condition taking sedatives or tranquilizers taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug <p>When using this product</p> <ul style="list-style-type: none"> take with food or milk if stomach upset occurs avoid alcoholic drinks be careful when driving a motor vehicle or operating machinery drowsiness may occur alcohol, sedatives, and tranquilizers may increase drowsiness may cause excitability especially in children <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> feel faint vomit blood have bloody or black stools have stomach pain that does not get better
---	--

BASE
PANEL

<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> you have symptoms of heart problems or stroke: <ul style="list-style-type: none"> chest pain trouble breathing weakness in one part or side of body slurred speech leg swelling pain gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days nasal congestion lasts for more than 7 days redness or swelling is present in the painful area you get nervous, dizzy, or sleepless symptoms continue or get worse any new symptoms appear <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed adults and children 12 years of age and over: <ul style="list-style-type: none"> take 1 tablet every 4 hours while symptoms persist do not take more than 6 tablets in any 24-hour period, unless directed by a doctor children under 12 years of age: do not use <p>Other information</p> <ul style="list-style-type: none"> read all warnings and directions before use. Keep carton. store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F) <p>Inactive ingredients acesulfame potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum</p> <p>Questions or comments? Call weekdays from 9 AM to 5 PM EST toll free at 1-800-88-ADVIL</p>
--	--

000065785

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

LIFT HERE

Drug Facts

Active ingredients (in each tablet) **Purposes**

Chlorpheniramine maleate 4 mgAntihistamine
Ibuprofen 200 mg (NSAID)*Pain reliever/Fever reducer
Phenylephrine HCl 10 mgNasal decongestant
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms associated with the common cold or flu:
 - headache
 - fever
 - minor body aches and pains
 - nasal congestion
 - runny nose
 - sinus pressure
 - sneezing
- reduces swelling of the nasal passages
- temporarily restores freer breathing through the nose

000065787

PANEL 1

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

Drug Facts (continued)

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- in children under 12 years of age
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

PANEL 2

Drug Facts (continued)

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if

- you have a breathing problem such as emphysema or chronic bronchitis
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn

Drug Facts (continued)

- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, thyroid disease, diabetes, glaucoma, have trouble urinating due to an enlarged prostate gland, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking sedatives or tranquilizers
- taking any other product that contains phenylephrine, chlorpheniramine or any other nasal decongestant or antihistamine
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

PANEL 3

Drug Facts (continued)

When using this product

- take with food or milk if stomach upset occurs
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- may cause excitability especially in children

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:

Drug Facts (continued)

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- nasal congestion lasts for more than 7 days
- redness or swelling is present in the painful area
- you get nervous, dizzy, or sleepless
- symptoms continue or get worse
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely

PANEL 4

Drug Facts (continued)

directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- adults and children 12 years of age and over:
 - take 1 tablet every 4 hours while symptoms persist
 - do not take more than 6 tablets in any 24-hour period, unless directed by a doctor
- children under 12 years of age: do not use

Other information

- read all warnings and directions before use. Keep carton.

Drug Facts (continued)

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)


Inactive ingredients

acesulfame potassium, artificial flavors, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, glycerin, glyceryl dibehenate, hypromellose, lactic acid, lecithin, maltodextrin, medium-chain triglycerides, microcrystalline cellulose, pharmaceutical ink, polydextrose, polyvinyl alcohol, pregelatinized starch, propyl gallate, silicon dioxide, sucralose, synthetic iron oxide, talc, titanium dioxide, triacetin, xanthan gum

Questions or comments?

Call weekdays from 9 AM to 5 PM EST toll free at 1-800-88-ADVIL

BASE PANEL



000065787

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
11/23/2021 01:42:29 PM