This label may not be the latest approved by FDA.
For current labeling information, please visit https://www.fda.gov/drugsatfda

Drug Facts

Active ingredient — **Purpose**

Diclofenac sodium (NSAID*) 1% (equivalent to 0.93% diclofenac) ..... Arthritis pain reliever

*nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of arthritis pain ONLY in the following areas:
  - hand, wrist, elbow (upper body areas) • foot, ankle, knee (lower body areas)
- this product may take up to 7 days to work for arthritis pain; it is not for immediate relief. If no pain relief in 7 days, stop use.

Warnings

For external use only

Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:
- hives • asthma (wheezing) • skin reddening • blisters • facial swelling • shock • rash

If an allergic reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains diclofenac. Liver damage may occur if you apply
- more or for a longer time than directed • when using other drugs containing diclofenac

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you
- are age 60 or older • have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- apply more or for longer than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever or to a fever reducer
- for sprains, strains, bruises or sports injuries. This product has not been shown to work for these types of injuries.
- right before or after heart surgery
- on more than 2 body areas at the same time • in the eyes, nose or mouth

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition • taking any other drug

▶

Drug Facts (continued)

When using this product

- avoid contact with eyes, nose, or mouth • if eye contact occurs, rinse thoroughly with water

Stop use and ask a doctor if

- pain gets worse or lasts more than 21 days • redness or swelling is present in the painful area
- fever occurs • skin irritation occurs
- any new symptoms appear. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding:
  - feel faint • have bloody or black stools • vomit blood
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke: • chest pain • trouble breathing • leg swelling
  - weakness in one part or side of body • slurred speech

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use this product during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Use up to 21 days unless directed by your doctor

Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

Daily	Per Dose
For your arthritis pain: • Use 4 times per day every day • Do not use on more than 2 body areas at the same time	**Use ENCLOSED DOSING CARD to measure a dose** • For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams)

Read the enclosed User Guide for complete instructions:

- use only as directed • do not use more than directed or for longer than directed
- apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
- do not apply in same area as any other product • do not apply with external heat such as heating pad
- do not apply a bandage over the treated area • store ENCLOSED DOSING CARD with your Voltaren Arthritis Pain. The dosing card is re-usable.

Other information

- store at 20-25°C (68-77°F). Keep from freezing.
- read all product information before using. Keep the dosing card; this carton and accompanying User Guide for important information.

Inactive ingredients

carbomer homopolymer Type C, cocoyl caprylocaprate, fragrance, isopropyl alcohol, mineral oil, polyoxyl 20 cetostearyl ether, propylene glycol, purified water, strong ammonia solution

Questions or comments?

1-855-297-3031 www.voltarengel.com

1-855-297-3031 A001245 Made in Canada
Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
Trademarks owned or licensed by GSK. ©2020 GSK or licensor.

A001245

TREATMENT AREAS

Medicated Gel Clinically Proven to Relieve Arthritis Pain

Not for use on any other body area (such as back, hip, or shoulder)

ORIGINAL PRESCRIPTION STRENGTH

NDC 0067-8152-08

gsk

Voltaren
ARTHRITIS PAIN

diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever

For Daily Treatment of Arthritis Pain
Anti-Inflammatory

TWIN VALUE PACK

Medicated Gel Clinically Proven to Relieve Arthritis Pain

Anti-Inflammatory

ORIGINAL PRESCRIPTION STRENGTH

gsk

Voltaren
ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever
For Daily Treatment of Arthritis Pain
Anti-Inflammatory

NET WT 2 x 100 g (3.53 oz) EACH, 200 g (7.06 oz) TOTAL

For external use only

This package contains 2 tubes of Voltaren Arthritis Pain

A001245

EXP./LOT A001245

2D = 001245

Use Enclosed Dosing Card to Measure a Dose



How to Use

- ☑ For arthritis pain only
- ☑ Use 4 times a day
- ☑ May take up to 7 days to work for your arthritis pain
- ☑ For use on no more than 2 body areas
- ☑ Use up to 21 days unless directed by your doctor

Tamper evident statement: Do not use if safety seal on tube nozzle is broken or missing.

GSK Regulatory Spec Box		Verified Date: 06.18.20
Drug Facts Info		
Drug Facts	Helvetica Bold Condensed Italic	9.38 pt.
Drug Facts (continued)	Helvetica Bold Condensed Italic / Condensed	8.34 pt.
Headings	Helvetica Bold Condensed Italic	8.34 pt.
Subheadings	Helvetica Bold Condensed	6.25 pt.
Body text	Helvetica Condensed / Bold Condensed	6.25 pt.
Bullets	Helvetica Condensed	5.25 pt.
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hair lines from edge of box-i.e. Minimum of 2 spaces either side of Drug Fact Box		Yes
Tracking: 0	Horizontal Scale:	90-100%
Leading: 6.78 pt.	Maximum Characters/Inch:	< 39
Barlines: 2.5 pt.	Hairlines:	0.5 pt.
Primary Display Panel Info		
Size of Net Wt/Contents (Smallest character height in inches)*		> 0.125 in.
PDP dimensions (in square inches)		5-25 sq. in.
Ratio of Statement of Identity to Logo/Largest Copy on PDP Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.) Vertical PDP: N/A	Horizontal PDP:	25%