

NDC 0067-8152-01

Voltaren
 ARTHRITIS PAIN
 diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

NET WT 0.7 oz (20 g)

To open tube: Unscrew cap and press the indent on the top of the cap onto the star-shaped seal on the tube. Firmly turn the cap to remove the safety seal.
 This tube does not contain full product information. Retain outer carton, dosing card, and User Guide for full product uses, directions and warnings.
 Tamper evident statement: Do not use if safety seal on tube nozzle is broken or missing.

Warnings

For external use only

Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains diclofenac. Liver damage may occur if you apply

- more or for a longer time than directed
- when using other drugs containing diclofenac

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- apply more or for longer than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

- Store at 20-25°C (68-77°F). Keep from freezing.

Questions or comments? 1-855-297-3031
 Dist. by GSK US, Warren, NJ 07059
 ©2021 GSK or licensor. Made in Canada NPA004474

Drug Facts (continued)
Read the enclosed User Guide for complete instructions:

- use only as directed
- do not use more than directed or for longer than directed
- apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
- do not apply in same area as any other product
- do not apply with external heat such as heating pad
- do not apply a bandage over the treated area
- store ENCLOSED DOSING CARD with your Voltaren Arthritis Pain. The dosing card is re-usable.

Other information
• store at 20-25°C (68-77°F). Keep from freezing.

PNL8

Drug Facts
Active ingredient Diclofenac sodium (NSAID) 1% (equivalent to 0.93% diclofenac) **Purpose** Arthritis pain reliever
*nonsteroidal anti-inflammatory drug

Uses
• for the temporary relief of arthritis pain (ONLY in the following areas:
• hand, wrist, elbow (upper body areas)
• foot, ankle, knee (lower body areas)
• this product may take up to 7 days to work for arthritis pain; it is not for immediate relief. If no pain relief in 7 days, stop use.

PNL1

Drug Facts (continued)
Warnings
For external use only
Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• hives • asthma (wheezing) • skin reddening • blisters • facial swelling • shock • rash
If an allergic reaction occurs, stop use and seek medical help right away.
Liver warning: This product contains diclofenac. Liver damage may occur if you apply
• more or for a longer time than directed
• when using other drugs containing diclofenac.

PNL2

Drug Facts (continued)
Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

Daily	Per Dose
For your arthritis pain: • Use 4 times per day every day • Do not use on more than 2 body areas at the same time	Use ENCLOSED DOSING CARD to measure a dose • For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams)

PNL7

Drug Facts (continued)

- feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- leg swelling
- weakness in one part or side of body
- slurred speech

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
Use up to 21 days unless directed by your doctor.

PNL6

Drug Facts (continued)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• apply more or for longer than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

PNL3

Drug Facts (continued)
Do not use
• if you have ever had an allergic reaction to any other pain reliever or to a fever reducer
• for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.
• right before or after heart surgery
• on more than 2 body areas at the same time
• in the eyes, nose or mouth

Ask a doctor before use if
• you have problems or serious side effects from taking pain relievers or fever reducers
• stomach bleeding warning applies to you
• you have a history of stomach problems, such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
• you are taking a diuretic
• you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.

PNL4

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are
• under a doctor's care for any serious condition
• taking any other drug

When using this product
• avoid contact with eyes, nose, or mouth
• if eye contact occurs, rinse thoroughly with water

Stop use and ask a doctor if
• pain gets worse or lasts more than 21 days
• redness or swelling is present in the painful area
• fever occurs
• skin irritation occurs
• any new symptoms appear. These could be signs of a serious condition.
• you experience any of the following signs of stomach bleeding:

PNL5

2D code= NPA004452

Drug Facts (continued)
• read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.

Inactive ingredients carbomer homopolymer Type C, cocoyl caprylocaprate, fragrance, isopropyl alcohol, mineral oil, polyvinyl 20 crosetate/ethyl ether, propylene glycol, purified water, strong anesthetic solution

Questions or comments? 1-855-297-3831 www.voltarenget.com

NPA004452

PNL9

To open tube: Unscrew cap and press the indent on the top of the cap onto the star-shaped seal on the tube.
Firmly turn the cap to remove the safety seal.

This tube does not contain full product information. Retain outer carton, dosing card, and User Guide for full product uses, directions and warnings.
Tamper evident statement: Do not use if safety seal on tube nozzle is broken or missing.

Warnings For external use only Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • asthma (wheezing) • skin reddening • blisters • facial swelling • shock • rash If an allergic reaction occurs, stop use and seek medical help right away. Liver warning: This product contains diclofenac. Liver damage may occur if you apply • more or for a longer time than directed • when using other drugs containing diclofenac Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription	NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • apply more or for longer than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. • Store at 20-25°C (68-77°F). Keep from freezing. Questions or comments? 1-855-297-3031 Dist. by: GSK CH, Warren, NJ 07059 ©2021 GSK or licensor. — Made in Canada — NPA004475
---	--

NDC 0067-8152-02

Voltaren

ARTHRITIS PAIN

diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

NET WT 1.76 oz (50 g)

EXP / LOT

NPA004475

To open tube: Unscrew cap and press the indent on the top of the cap onto the star-shaped seal on the tube. Firmly turn the cap to remove the safety seal.
This tube does not contain full product information.
Retain outer carton, dosing card, and User Guide for full product uses, directions and warnings.
Tamper evident statement: Do not use if safety seal on tube nozzle is broken or missing.

Warnings
For external use only
Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• hives • asthma (wheezing) • skin reddening
• blisters • facial swelling • shock • rash
If an allergic reaction occurs, stop use and seek medical help right away.
Liver warning: This product contains diclofenac. Liver damage may occur if you apply
• more or for a longer time than directed
• when using other drugs containing diclofenac
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)


• have 3 or more alcoholic drinks every day while using this product
• apply more or for longer than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
• Store at 20-25°C (68-77°F). Keep from freezing.
Questions or comments? 1-855-297-3031
Dist. by: GSK CH, Warren, NJ 07059
©2021 GSK or licensor. Made in Canada NPA004454

NPA004454
EXP / LOT

NDC 0067-8152-03

Voltaren
ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

NET WT 3.53 oz (100 g)



2D = NPA004454

To open tube: Unscrew cap and press the indent on the top of the cap onto the star-shaped seal on the tube. Firmly turn the cap to remove the safety seal.	
This tube does not contain full product information. Retain outer carton, dosing card, and User Guide for full product uses, directions and warnings.	
Tamper evident statement: Do not use if safety seal on tube nozzle is broken or missing.	
Warnings For external use only Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • asthma (wheezing) • skin reddening • blisters • facial swelling • shock • rash If an allergic reaction occurs, stop use and seek medical help right away. Liver warning: This product contains diclofenac. Liver damage may occur if you apply • more or for a longer time than directed • when using other drugs containing diclofenac Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)	• have 3 or more alcoholic drinks every day while using this product • apply more or for longer than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. • Store at 20-25°C (68-77°F). Keep from freezing. Questions or comments? 1-855-297-3031 Dist. by: GSK CH , Warren, NJ 07059 ©2021 GSK or licensor. Made in Canada NPA004481
NDC 0067-8152-07	EXP / LOT
	
Voltaren ARTHRITIS PAIN diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever	
NET WT 4.23 oz (120 g)	
gsk	

To open tube: Unscrew cap and press the indent on the top of the cap onto the star-shaped seal on the tube. Firmly turn the cap to remove the safety seal.	This tube does not contain full product information. Retain outer carton, dosing card, and User Guide for full product uses, directions and warnings.
---	--

Tamper evident statement: Do not use if safety seal on tube nozzle is broken or missing

Warnings
For external use only
Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• hives • asthma (wheezing) • skin reddening • blisters
• facial swelling • shock • rash
If an allergic reaction occurs, stop use and seek medical help right away.
Liver warning: This product contains diclofenac. Liver damage may occur if you apply
• more or for a longer time than directed
• when using other drugs containing diclofenac
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription

NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• apply more or for longer than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
• Store at 20-25°C (68-77°F). Keep from freezing.

Questions or comments? 1-855-297-3031
Dist. by: **GSK CH**, Warren, NJ 07059
©2021 GSK or licensor. Made in Canada NPA004471

NDC 0067-8152-04

Voltaren
ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

NET WT 5.29 oz (150 g)

gsk

EXP / LOT

NPA004471

Drug Facts
Keep this and the accompanying dosing card with you at all times. Read this important information about Voltaren Arthritis Pain every time you use this product. This information does not take the place of other directions you may receive from your doctor about this product. This information does not take the place of other directions you may receive from your doctor about this product. This information does not take the place of other directions you may receive from your doctor about this product.

Warnings
Do not use Voltaren Arthritis Pain if you are allergic to diclofenac sodium or any of the other ingredients listed under "Active ingredient." Do not use Voltaren Arthritis Pain if you are allergic to aspirin, salicylates, or other nonsteroidal anti-inflammatory drugs (NSAIDs). Do not use Voltaren Arthritis Pain if you are taking aspirin, salicylates, or other NSAIDs for heart, blood pressure, or other medical conditions. Do not use Voltaren Arthritis Pain if you are taking aspirin, salicylates, or other NSAIDs for heart, blood pressure, or other medical conditions.

Directions
Apply to the affected area 4 times a day. Do not use more than directed. Do not use Voltaren Arthritis Pain for more than 21 days unless directed by your doctor. Do not use Voltaren Arthritis Pain for more than 21 days unless directed by your doctor.

Other information
Keep this and the accompanying dosing card with you at all times. Read this important information about Voltaren Arthritis Pain every time you use this product. This information does not take the place of other directions you may receive from your doctor about this product. This information does not take the place of other directions you may receive from your doctor about this product.

Questions or comments?
1-855-297-3031
www.voltaren.com

TREATMENT AREAS

Medicated Gel Clinically Proven to Relieve Arthritis Pain

Not for use on any other body area (such as back, hip, or shoulder)

EASY TWIST CAP

ORIGINAL PRESCRIPTION STRENGTH

Voltaren
ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever
For Daily Treatment of Arthritis Pain
Anti-Inflammatory

ORIGINAL PRESCRIPTION STRENGTH

Voltaren
ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever
For Daily Treatment of Arthritis Pain
Anti-Inflammatory

For external use only
This package contains 2 tubes of Voltaren Arthritis Pain

TWIN VALUE PACK

Medicated Gel Clinically Proven to Relieve Arthritis Pain
Anti-Inflammatory

NET WT 2 x 100 g (3.53 oz) EACH, 200 g (7.06 oz) TOTAL

1001294

1-855-297-3031
Made in Canada
Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
Trademarks owned or licensed by GSK. ©2017 GSK or its affiliates.

Use Enclosed Dosing Card to Measure a Dose

Voltaren ARTHRITIS PAIN Dosing Card (NOT ACTUAL SIZE)
diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever
Read product User Guide before use
Not for immediate pain relief

Upper body dose 2.25 inches long (2 grams)
Lower body dose 4.5 inches long (4 grams)

How to Use

- For arthritis pain only
- Use 4 times a day
- May take up to 7 days to work for your arthritis pain
- For use on no more than 2 body areas
- Use up to 21 days unless directed by your doctor

Tamper evident statement: Do not use if safety seal on tube nozzle is broken or missing.

1001294

2D = 1001294

1001294

Voltaren

ARTHRITIS PAIN



diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

Use Enclosed Dosing Card to Measure a Dose

Voltaren ARTHRITIS PAIN Dosing Card (NOT ACTUAL SIZE)
diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

Read product User Guide before use
Not for immediate pain relief

Upper body dose 2.25 inches long (2 grams)

Lower body dose 4.5 inches long (4 grams)

How to Use

- For arthritis pain only
- Use 4 times a day
- May take up to 7 days to work for your arthritis pain
- For use on no more than 2 body areas
- Use up to 21 days unless directed by your doctor

<p>Drug Facts</p> <p>Active ingredient Diclofenac sodium (NSAID*) 1% (equivalent to 0.93% diclofenac) Arthritis pain reliever *nonsteroidal anti-inflammatory drug</p> <p>Purpose Arthritis pain reliever</p> <p>Uses • for the temporary relief of arthritis pain <u>ONLY</u> in the following areas: • hand, wrist, elbow (upper body areas) • foot, ankle, knee (lower body areas) • this product may take up to 7 days to work for arthritis pain; it is not for immediate relief. If no pain relief in 7 days, stop use.</p> <p>Warnings For external use only Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • asthma (wheezing) • skin reddening • blisters • facial swelling • shock • rash If an allergic reaction occurs, stop use and seek medical help right away. Liver warning: This product contains diclofenac. Liver damage may occur if you apply • more or for a longer time than directed • when using other drugs containing diclofenac Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • apply more or for longer than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use • if you have ever had an allergic reaction to any other pain reliever or to a fever reducer • for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries. • right before or after heart surgery • on more than 2 body areas at the same time • in the eyes, nose or mouth</p> <p>Ask a doctor before use if • you have problems or serious side effects from taking pain relievers or fever reducers • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke • you are taking a diuretic • you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.</p> <p>Ask a doctor or pharmacist before use if you are • under a doctor's care for any serious condition • taking any other drug</p>		<p>Drug Facts (continued)</p> <p>When using this product • avoid contact with eyes, nose, or mouth • if eye contact occurs, rinse thoroughly with water</p> <p>Stop use and ask a doctor if • pain gets worse or lasts more than 21 days • redness or swelling is present in the painful area • fever occurs • skin irritation occurs • any new symptoms appear. These could be signs of a serious condition. • you experience any of the following signs of stomach bleeding: • feel faint • have bloody or black stools • vomit blood • have stomach pain that does not get better • you have symptoms of heart problems or stroke: • chest pain • trouble breathing • leg swelling • weakness in one part or side of body • slurred speech</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions Use up to 21 days unless directed by your doctor Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.</p> <table border="1"> <thead> <tr> <th>Daily</th> <th>Per Dose</th> </tr> </thead> <tbody> <tr> <td> <p>For your arthritis pain: • Use 4 times per day every day • Do not use on more than 2 body areas at the same time</p> </td> <td> <p>Use ENCLOSED DOSING CARD to measure a dose • For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams)</p> </td> </tr> </tbody> </table> <p>Read the enclosed User Guide for complete instructions: • use only as directed • do not use more than directed or for longer than directed • apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes • do not apply in same area as any other product • do not apply with external heat such as heating pad • do not apply a bandage over the treated area • store ENCLOSED DOSING CARD with your Voltaren Arthritis Pain. The dosing card is re-usable.</p> <p>Other information • store at 20-25°C (68-77°F). Keep from freezing. • read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.</p> <p>Inactive ingredients carbomer homopolymer Type C, cocoyl caprylocaprate, fragrance, isopropyl alcohol, mineral oil, polyoxyl 20 cetostearyl ether, propylene glycol, purified water, strong ammonia solution</p> <p>Questions or comments? 1-855-297-3031 www.voltaren.com</p>		Daily	Per Dose	<p>For your arthritis pain: • Use 4 times per day every day • Do not use on more than 2 body areas at the same time</p>	<p>Use ENCLOSED DOSING CARD to measure a dose • For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams)</p>
Daily	Per Dose						
<p>For your arthritis pain: • Use 4 times per day every day • Do not use on more than 2 body areas at the same time</p>	<p>Use ENCLOSED DOSING CARD to measure a dose • For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams)</p>						

1369089



1-855-297-3031 Made in Canada
Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
Trademarks are owned by or licensed to the GSK group of companies. 1001283
©2021 GSK group of companies or its licensor.

Tamper evident statement: Do not use if safety seal on tube nozzle is broken or missing.

Lot:
Exp:



Voltaren

ARTHRITIS PAIN



diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

Use Enclosed Dosing Card to Measure a Dose

Voltaren ARTHRITIS PAIN Dosing Card (NOT ACTUAL SIZE)
diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

Read product User Guide before use
Not for immediate pain relief

Upper body dose 2.25 inches long (2 grams)

Lower body dose 4.5 inches long (4 grams)

How to Use

- For arthritis pain *only*
- Use 4 times a day
- May take up to 7 days to work for your arthritis pain
- For use on no more than 2 body areas
- Use up to 21 days unless directed by your doctor

<p>Drug Facts</p> <p>Active ingredient Diclofenac sodium (NSAID)* 1% (equivalent to 0.93% diclofenac) Arthritis pain reliever *nonsteroidal anti-inflammatory drug</p> <p>Purpose Arthritis pain reliever</p> <p>Uses • for the temporary relief of arthritis pain <u>ONLY</u> in the following areas: • hand, wrist, elbow (upper body areas) • foot, ankle, knee (lower body areas) • this product may take up to 7 days to work for arthritis pain; it is not for immediate relief. If no pain relief in 7 days, stop use.</p> <p>Warnings For external use only Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • asthma (wheezing) • skin reddening • blisters • facial swelling • shock • rash If an allergic reaction occurs, stop use and seek medical help right away. Liver warning: This product contains diclofenac. Liver damage may occur if you apply • more or for a longer time than directed • when using other drugs containing diclofenac Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • apply more or for longer than directed Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Do not use • if you have ever had an allergic reaction to any other pain reliever or to a fever reducer • for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries. • right before or after heart surgery • on more than 2 body areas at the same time • in the eyes, nose or mouth Ask a doctor before use if • you have problems or serious side effects from taking pain relievers or fever reducers • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke • you are taking a diuretic • you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years. Ask a doctor or pharmacist before use if you are • under a doctor's care for any serious condition • taking any other drug</p>		<p>Drug Facts (continued)</p> <p>When using this product • avoid contact with eyes, nose, or mouth • if eye contact occurs, rinse thoroughly with water</p> <p>Stop use and ask a doctor if • pain gets worse or lasts more than 21 days • redness or swelling is present in the painful area • fever occurs • skin irritation occurs • any new symptoms appear. These could be signs of a serious condition. • you experience any of the following signs of stomach bleeding: • feel faint • have bloody or black stools • vomit blood • have stomach pain that does not get better • you have symptoms of heart problems or stroke: • chest pain • trouble breathing • leg swelling • weakness in one part or side of body • slurred speech</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</p> <p>Directions Use up to 21 days unless directed by your doctor Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.</p> <table border="1"> <thead> <tr> <th>Daily</th> <th>Per Dose</th> </tr> </thead> <tbody> <tr> <td> <p>For your arthritis pain:</p> <ul style="list-style-type: none"> • Use 4 times per day every day • Do not use on more than 2 body areas at the same time </td> <td> <p>Use ENCLOSED DOSING CARD to measure a dose</p> <ul style="list-style-type: none"> • For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams) </td> </tr> </tbody> </table> <p>Read the enclosed User Guide for complete instructions:</p> <ul style="list-style-type: none"> • use only as directed • do not use more than directed or for longer than directed • apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes • do not apply in same area as any other product • do not apply with external heat such as heating pad • do not apply a bandage over the treated area • store ENCLOSED DOSING CARD with your Voltaren Arthritis Pain. The dosing card is re-usable. <p>Other information • store at 20-25°C (68-77°F). Keep from freezing. • read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.</p> <p>Inactive ingredients carbomer homopolymer Type C, cocoyl caprylocaprate, fragrance, isopropyl alcohol, mineral oil, polyoxyl 20 cetostearyl ether, propylene glycol, purified water, strong ammonia solution</p> <p>Questions or comments? 1-855-297-3031 www.voltarengel.com</p>		Daily	Per Dose	<p>For your arthritis pain:</p> <ul style="list-style-type: none"> • Use 4 times per day every day • Do not use on more than 2 body areas at the same time 	<p>Use ENCLOSED DOSING CARD to measure a dose</p> <ul style="list-style-type: none"> • For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams)
Daily	Per Dose						
<p>For your arthritis pain:</p> <ul style="list-style-type: none"> • Use 4 times per day every day • Do not use on more than 2 body areas at the same time 	<p>Use ENCLOSED DOSING CARD to measure a dose</p> <ul style="list-style-type: none"> • For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) – Squeeze out 4.5 inches (4 grams) 						

0000000



1-855-297-3031 Made in Canada
Distributed by: GSK Consumer Healthcare, Warren, NJ 07059
Trademarks are owned by or licensed to the GSK group of companies. NPA004492
©2021 GSK group of companies or its licensor.

Tamper evident statement: Do not use if safety seal on tube nozzle is broken or missing.

Lot:
Exp:




Voltaren
ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

User Guide
How to use Voltaren Arthritis Pain, with answers to frequently asked questions

Quick Start
How to get started right away with Voltaren Arthritis Pain

Understanding Voltaren Arthritis Pain Gel
Voltaren Arthritis Pain (diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever) is an arthritis pain relief medication that you apply to your skin.

Before You Use Voltaren Arthritis Pain
Read the Drug Facts Label, which appears on the carton.



For each lower body area (a foot, an ankle, or a knee): Squeeze gel from the tube equal to the length shown on the lower body section of the dosing card (4.5 inches).


Lost Your Dosing Card? Don't worry! Call 1-855-297-3031 to get a free replacement.

Applying Voltaren Arthritis Pain
Gently rub Voltaren Arthritis Pain into the skin using your hand. Apply 4 times a day for best results. Do not use on more than 2 body areas at once.


Apply only to clean, dry skin that doesn't have any cuts, open wounds, infections or rashes. Do not apply in the same area as any other medicines or products that are applied to the skin.

Upper body areas	Lower body areas
Elbow Including sides, inner, and outer	Knee Including sides, kneecap, and back
Wrist Including sides, top, and bottom	Ankle Including sides, front, and back
Hand Including top, palm, and fingers	Foot Including sole, top of foot, and toes


After You Use Voltaren Arthritis Pain



Wash your hands after applying Voltaren Arthritis Pain. If the treated area is the hands wait up to 1 hour to wash the hands.



Wash the dosing card with water. Store dosing card with your Voltaren Arthritis Pain and keep out of reach of children.



Wait ten minutes before covering the treated areas with clothes or gloves.

Avoid:

- showering or bathing for at least 1 hour after use
- exposing the treated area to sunlight or artificial light (such as in tanning booths)
- use in same area as other products applied to the skin
- applying a bandage over the treated area
- eye, nose, or mouth contact
- if eye contact occurs, rinse thoroughly with water

Facts you need to know
Important information about Voltaren Arthritis Pain

Do not use Voltaren Arthritis Pain...

- if you have ever had an allergic reaction to any other pain reliever or to a fever reducer
- for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.
- right before or after heart surgery
- on more than 2 body areas at the same time
- in the eyes, nose or mouth

Ask a doctor before use if...

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.


Ask a doctor or pharmacist before use if...

- you are under a doctor's care for any serious condition
- you are taking any other drug
- you are pregnant or breast-feeding

NPA004472


Remove the dosing card from the inside of the carton. You should always use the dosing card to measure out the correct dose of Voltaren Arthritis Pain.

The first time you use a tube of Voltaren Arthritis Pain: You will have either one of the caps shown below.




- Take the cap off the tube. Open the safety seal by firmly pressing the indent on the top of the cap onto the star-shaped seal on the tube.
- Firmly turn the cap to remove the safety seal. Do not open the safety seal with scissors or other sharp objects.
- After use, put the cap back on the end of the tube.


Using Voltaren Arthritis Pain
How often to use Voltaren Arthritis Pain
Use 4 times a day for best results.
For example:



Where to apply Voltaren Arthritis Pain
Voltaren Arthritis Pain can be used on up to 2 body areas from the following list: a hand, a wrist, an elbow, a foot, an ankle, a knee. Do not use on more than 2 body areas at once.
Some examples include:



Measuring the correct amount using the dosing card



For each upper body area (a hand, a wrist, or an elbow): Squeeze gel from the tube equal to the length shown on the upper body section of the dosing card (2.25 inches).

Turn over for more information

It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Warnings to consider before use...

Allergy alert
Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- rash
- blisters
- facial swelling
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

Liver warning
This product contains diclofenac. Liver damage may occur if you apply

- more or for a longer time than directed
- when using other drugs containing diclofenac

Stomach bleeding warning
This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- apply more or for longer than directed

Heart attack and stroke warning
NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. This can be fatal. The risk is higher if you use more than directed or for longer than directed.

Follow these important guidelines when using Voltaren Arthritis Pain...

- use only as directed
- apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
- do not apply in same area as any other product used on the skin
- do not apply with external heat such as heating pad
- do not apply a bandage over the treated area
- do not get gel in eyes, nose or mouth
- store ENCLOSED DOSING CARD with your Voltaren Arthritis Pain. The dosing card is re-usable.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Stop using Voltaren Arthritis Pain and ask a doctor if...

- pain gets worse or lasts more than 21 days
- redness or swelling is present in the painful area
- fever occurs
- skin irritation occurs
- any new symptoms appear. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - leg swelling
 - weakness in one part or side of body
 - slurred speech


Questions and Answers
Answers to questions you may ask about Voltaren Arthritis Pain

Can I use Voltaren Arthritis Pain for the immediate relief of pain like that associated with sprains, strains, bruises or sports injuries?

No.
Do not use for sprains, sprains, bruises, or sports injuries. This product has not been shown to work for these types of injuries. It's intended only for the temporary relief of arthritis pain.

How quickly will Voltaren Arthritis Pain work for arthritis pain?
Voltaren Arthritis Pain may take up to 7 days to work for your arthritis pain; it is not for immediate relief. With 4 times-a-day use, you may start to feel relief within a few days. You should feel significant pain relief within 7 days of continuous use. If no pain relief in 7 days, stop use.

What is Voltaren Arthritis Pain used for?
Voltaren Arthritis Pain can be used for the temporary relief of arthritis pain in the hands, wrists, elbows, feet, ankles, and knees. Osteoarthritis (OA) is the most common form of arthritis. It mainly affects a type of tissue called cartilage, which cushions the joints and prevents the bones from rubbing against each other. With OA, cartilage begins to break down and wear away, resulting in joint pain and stiffness. OA occurs more frequently as you get older. Pain develops slowly and can gradually worsen over time.



How does Voltaren Arthritis Pain work?
Diclofenac sodium, the medicine in Voltaren Arthritis Pain, is a nonsteroidal anti-inflammatory drug (NSAID). NSAIDs are used to treat pain from medical conditions such as arthritis. Voltaren Arthritis Pain works similarly to oral NSAIDs like ibuprofen or naproxen by temporarily blocking the production of pain signaling chemicals called prostaglandins. However, when taking a pill, the medication is distributed to the site of pain through the bloodstream. Compared to oral diclofenac, only 6% of the medicine in Voltaren Arthritis Pain is absorbed in the bloodstream. It works by penetrating through your skin at the application site to deliver arthritis pain relief.

How does over-the-counter Voltaren Arthritis Pain differ from prescription Voltaren Gel?
Over-the-counter Voltaren Arthritis Pain is the same strength and formula as original full prescription strength Voltaren Gel but is available without a prescription.

Apply Voltaren Arthritis Pain 4 times a day to ensure you get the full effect of the medicine.

Can I apply Voltaren Arthritis Pain to upper and lower body areas at the same time?
Yes.
You can use Voltaren Arthritis Pain on up to 2 body areas at the same time. The areas can include the foot, ankle, knee, hand, wrist, and elbow.
Some correct examples include:



Do not use Voltaren Arthritis Pain on more than 2 body areas at once.

Can I apply Voltaren Arthritis Pain without measuring it out on the dosing card first?
No.
Use the dosing card to make sure you are getting the correct amount of medicine.

Can I use Voltaren Arthritis Pain on my spine, hips, or shoulders?
No.
The use of Voltaren Arthritis Pain on the spine, hips, and shoulders has not been studied.

Can I take any other pain medications with Voltaren Arthritis Pain?
Ask your doctor or pharmacist before using Voltaren Arthritis Pain if you are taking any other pain reliever or are planning on taking any other pain reliever, especially prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others).

Can I take any other pain medications during the first 7 days I use Voltaren Arthritis Pain?
Taking other pain medications is generally not recommended. Speak with your doctor or pharmacist before taking other pain medications when using Voltaren Arthritis Pain.

How long can I use Voltaren Arthritis Pain?
Use Voltaren Arthritis Pain 4 times a day every day for up to 21 days for treatment of arthritis pain or as directed by your doctor.
Voltaren Arthritis Pain should be part of an overall program for managing your arthritis pain. The program should also include activities such as appropriate movement or exercise, specific lifestyle changes, and weight maintenance. Regular visits and discussions with your doctor can help incorporate Voltaren Arthritis Pain into a complete program to help manage your arthritis pain.

What do I do if I get Voltaren Arthritis Pain in my eyes?
If eye contact occurs, rinse thoroughly with water and consult a doctor if irritation lasts more than an hour.

How do I clean the dosing card?
Wash the dosing card with water, and store dosing card and tube out of reach of children.

What if I miss a dose of Voltaren Arthritis Pain?
If you forget to apply Voltaren Arthritis Pain, don't double your dose. Just apply the next dose as scheduled.

How should I store Voltaren Arthritis Pain?

- Store at 20-25°C (68-77°F).
- Do not freeze Voltaren Arthritis Pain.
- Store the dosing card with your Voltaren Arthritis Pain.
- Keep out of reach of children.

Additional Questions?
Call the Voltaren Arthritis Pain consumer hotline at 1-855-297-3031.
www.voltarengel.com

NPA004472

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

VALERIE S PRATT
11/03/2021 04:39:07 PM