

This tube does not contain full product information.
Retain outer carton, dosing card, and User Guide for full product uses, directions and warnings.
Tamper evident statement: Do not use if plastic safety seals on sides of cap are broken or missing.

Warnings
For external use only
Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• hives • asthma (wheezing) • skin reddening
• blisters • facial swelling • shock • rash
If an allergic reaction occurs, stop use and seek medical help right away.
Liver warning: This product contains diclofenac. Liver damage may occur if you apply
• more or for a longer time than directed
• when using other drugs containing diclofenac
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

• have 3 or more alcoholic drinks every day while using this product
• apply more or for longer than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
• Store at 20-25°C (68-77°F). Keep from freezing.
Questions or comments? 1-855-297-3031
Dist. by: GSK Consumer Healthcare, Warren, NJ 07059
©2022 GSK or licensor. Made in Canada NPA007807

EXP. / LOT

AA027907

NDC 0067-8152-03

Voltaren
ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

NET WT 100 g (3.53 oz)

gsk

GSK Regulatory Spec Box		Verified Date: 06.01.22
Drug Facts Info		
Drug Facts		N/A
Drug Facts (continued)		N/A
Headings	Helvetica Bold Condensed Italic	6.75 pt.
Subheadings	Helvetica Bold Condensed	6.75 pt.
Body text	Helvetica Condensed / Bold Condensed	6.75 pt.
Bullets	Helvetica Condensed	6.75 pt.
Bullets on same lines: end of statement separated from bulleted statement by two ems		
Spacing of the hair lines from edge of box -i.e. Minimum of 2 spaces either side of Drug Fact Box		
Tracking:	0	Horizontal Scale: 83%
Leading:	7 pt.	Maximum Characters/Inch: < 39
Barlines:	0.76 pt	Hairlines: 0.665 pt
Primary Display Panel Info		
Size of Net Wt/Contents (Smallest character height in inches)*		> 0.125 in.
PDP dimensions (in square inches)		5-25 sq. in.
Ratio of Statement of Identity to Logo/Largest Copy on PDP		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		
Vertical PDP: N/A		Horizontal PDP: 25%

This tube does not contain full product information.
Retain outer carton, dosing card, and User Guide for full product uses, directions and warnings.
Tamper evident statement: Do not use if plastic safety seals on sides of cap are broken or missing.

Warnings
For external use only
Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• hives • asthma (wheezing) • skin reddening
• blisters • facial swelling • shock • rash
If an allergic reaction occurs, stop use and seek medical help right away.
Liver warning: This product contains diclofenac. Liver damage may occur if you apply
• more or for a longer time than directed
• when using other drugs containing diclofenac
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

• have 3 or more alcoholic drinks every day while using this product
• apply more or for longer than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
• Store at 20-25°C (68-77°F). Keep from freezing.
Questions or comments? 1-855-297-3031
Dist. by: GSK CH, Warren, NJ 07059
©2022 GSK or licensor. Made in Canada NPA007813

EXP/LOT
A007813
FPO

NDC 0067-8152-07

Voltaren
ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

NET WT 120 g (4.23 oz)

GSK Regulatory Spec Box		Verified Date: 06.02.22
Drug Facts Info		
Drug Facts	Helvetica Bold Condensed Italic	N/A
Drug Facts (continued)	Helvetica Bold Condensed Italic / Condensed	N/A
Headings	Helvetica Bold Condensed Italic	6.75 pt.
Subheadings	Helvetica Bold Condensed	6.75 pt.
Body text	Helvetica Condensed / Bold Condensed	6.75 pt.
Bullets	Helvetica Condensed	6.75 pt.
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hair lines from edge of box -i.e. Minimum of 2 spaces either side of Drug Fact Box		N/A
Tracking:	0	Horizontal Scale: 83%
Leading:	7 pt.	Maximum Characters/Inch: < 39
Barlines:	N/A	Hairlines: .665 pt.
Primary Display Panel Info		
Size of Net Wt/Contents (Smallest character height in inches)*		> 0.125 in.
PDP dimensions (in square inches)		5-25 sq. in.
Ratio of Statement of Identity to Logo/Largest Copy on PDP Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.) Vertical PDP: N/A Horizontal PDP: 25%		

This tube does not contain full product information.
Retain outer carton, dosing card, and User Guide for full product uses, directions and warnings.
Tamper evident statement: Do not use if plastic safety seals on sides of cap are broken or missing.

Warnings
For external use only
Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
• hives • asthma (wheezing) • skin reddening • blisters
• facial swelling • shock • rash
If an allergic reaction occurs, stop use and seek medical help right away.
Liver warning: This product contains diclofenac. Liver damage may occur if you apply
• more or for a longer time than directed
• when using other drugs containing diclofenac
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you
• are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription

NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• apply more or for longer than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
• Store at 20-25°C (68-77°F). Keep from freezing.

Questions or comments? 1-855-297-3031

Dist. by: **GSK Consumer Healthcare**, Warren, NJ 07059
©2022 GSK or licensor. Made in Canada NPA007808

NDC 0067-8152-04

EXP. / LOT

EA007808



Voltaren

ARTHRITIS PAIN

diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever

NET WT 150 g (5.29 oz)

GSK Regulatory Spec Box		Verified Date: 06.02.22
Drug Facts Info		
Drug Facts		N/A
Drug Facts (continued)		N/A
Headings	Helvetica Bold Condensed Italic	7.75 pt.
Subheadings	Helvetica Bold Condensed	7.75 pt.
Body text	Helvetica Condensed / Bold Condensed	7.75 pt.
Bullets	Helvetica Condensed	5 pt.
Bullets on same lines: end of statement separated from bulleted statement by two ems		Yes
Spacing of the hair lines from edge of box -i.e. Minimum of 2 spaces either side of Drug Fact Box		No
Tracking:	0	Horizontal Scale: 83%
Leading:	7.85 pt.	Maximum Characters/Inch: ≤ 39
Barlines:	0.5 pt.	Hairlines: 0.5 pt.
Primary Display Panel Info		
Size of Net Wt/Contents (Smallest character height in inches)*		> 0.125 in.
PDP dimensions (in square inches)		5-25 sq. in.
Ratio of Statement of Identity to Logo/Largest Copy on PDP Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		
Vertical PDP: N/A		Horizontal PDP: 25%

ORIGINAL PRESCRIPTION STRENGTH

Voltaren

ARTHRITIS PAIN

diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever

ANTI-INFLAMMATORY

For Daily Treatment of Arthritis Pain

NDC 0067-8153-01 

Medicated Gel Clinically Proven to Relieve Arthritis Pain

NEW FLIP TOP CAP*

*Cap for 150 g only

For external use only

TREATMENT AREAS



Hand Wrist
Elbow Foot
Ankle Knee

Not for use on any other body area (such as back, hip, or shoulder)

ORIGINAL PRESCRIPTION STRENGTH

Voltaren

ARTHRITIS PAIN

diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever

ANTI-INFLAMMATORY

For Daily Treatment of Arthritis Pain

Use Enclosed Dosing Card to Measure a Dose

Voltaren ARTHRITIS PAIN Dosing Card (NOT ACTUAL SIZE)

diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever

Upper body dose: 2.25 inches long (2 grams)

Lower body dose: 4.5 inches long (4 grams)

How to Use

- For arthritis pain only
- Use 4 times a day
- May take up to 7 days to work for your arthritis pain
- For use on no more than 2 body areas
- Use up to 21 days unless directed by your doctor

Drug Facts

Active ingredient: diclofenac sodium (NSAID) 1% (equivalent to 0.5% diclofenac) Arthritis pain reliever

Purpose: Nonsteroidal anti-inflammatory drug

Uses: For the temporary relief of arthritis pain (OAR) in the following areas: hand, wrist, elbow (upper body areas); foot, ankle, knee (lower body areas). This product may take up to 7 days to work for arthritis pain. It is not for immediate relief. If no pain relief in 7 days, stop use.

Warnings: For external use only. **Allergy alert:** Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Directions may include: • have • asthma (asthma) • skin rash • hives • facial swelling • shock • rash. If an allergic reaction occurs, stop use and seek medical help right away.

User warning: This product contains diclofenac. Liver damage may occur if you apply more or for longer time than directed • when using other drugs containing diclofenac.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinner (anticoagulant) or aspirin drug.

• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use: • if you have ever had an allergic reaction to any other pain reliever or to a liver reducer • for strains, sprains, tears or sports injuries. This product has not been shown to work for these types of injuries. • right before or after heart surgery • on more than 2 body areas at the same time • in the eyes, nose or mouth.

Ask a doctor before use if: • you have problems or serious side effects from taking pain relievers or fever reducers • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke • you are taking a diuretic • you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years. • you are taking a diuretic or aspirin before use if you are • under a doctor's care for any serious condition • taking any other drug.

When using this product: • avoid contact with eyes, nose, or mouth • if eye contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if: • pain gets worse or lasts more than 21 days • redness or swelling is present in the painful area • fever occurs • skin irritation occurs • any new symptoms appear. These could be signs of a serious condition.

If pregnant or breast-feeding: ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions: Use up to 21 days unless directed by your doctor. Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

Apply	Per Dose
For your arthritis pain:	Use ENCLOSED DOSING CARD to measure a dose
• Use 4 times per day every day	• For each upper body area (hand, wrist, or elbow): Space out 2.25 inches (2 grams)
• Do not use on more than 2 body areas at the same time	• For each lower body area (foot, ankle, or knee): Space out 4.5 inches (4 grams)

Read the enclosed Use Guide for complete instructions: • use only as directed • do not use more than directed or for longer than directed • apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes • do not apply in same area as any other product • do not apply with external heat such as heating pad • do not apply a bandage over the treated area • store ENCLOSED DOSING CARD with your Voltaren Arthritis Pain. The dosing card is re-usable.

Other information: • store at 20°-25° (68°-77°). Keep from freezing. • read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.

Inactive ingredients: caprylene nonperfluoropolyether, colorant (polyethylene glycol, titanium dioxide, iron oxides, iron hydroxide).

Questions or comments? 1-800-261-2031 www.voltaren.com

NPA007810

1-855-297-3031 NPA007810 Made in Canada
Distributed by: **GSK Consumer Healthcare**, Warren, NJ 07069
Trademarks are owned by or licensed to the GSK group of companies.
©2022 GSK group of companies or its licensee.

LOT EXP

PULL TAB

Voltaren ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever

Voltaren ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever

Voltaren ARTHRITIS PAIN
diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever

NET WT 150 g (5.29 oz)
NET WT 150 g (5.29 oz)
NET WT 50 g (1.76 oz)

3 TUBES: 150 g + 150 g + 50 g TOTAL NET WT 350 g (12.34 oz)

CARTON RECYCLE

GSK Regulatory Spec Box Verified Date: 10.20.22

Drug Facts Info

Drug Facts	Helvetica Bold Condensed Italic	11 pt.
Drug Facts (continued)	Helvetica Bold Condensed Italic / Condensed	10 pt.
Headings	Helvetica Bold Condensed Italic	10 pt.
Subheadings	Helvetica Bold Condensed	7 pt.
Body text	Helvetica Condensed / Bold Condensed	7 pt.
Bullets	Helvetica Condensed	6 pt.
Bullets on same line, end of statement separated from bulleted statement by two ems		Yes
Spacing of the first line from edge of box-ca. Minimum of 2 spaces after side of Drug Fact Box		Yes
Tracking	0	Horizontal Scale: 88 – 100%
Leading	0 pt.	Maximum Characters/Inch: < 30
Burline	2.5 pt.	Hairline
Primary Display Panel Info		
Size of Net Wt/Contents (Smallest character height in inches)		> 0.1875 in.
PDP dimensions (in square inches)		26-100 sq. in.
Ratio of Statement or Identity to Logo (Largest Copy on PDP)		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		
Vertical PDP: 27%		Horizontal PDP: N/A

ORIGINAL PRESCRIPTION STRENGTH

Voltaren

ARTHRITIS PAIN

diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever

ANTI-INFLAMMATORY

For Daily Treatment of Arthritis Pain



NDC 0067-8152-05

Medicated Gel
Clinically Proven
to Relieve
Arthritis Pain



NEW
FLIP TOP
CAP

For external use only

TREATMENT AREAS

Hand

Wrist

Elbow

Foot

Ankle

Knee

Not for use on any other body area (such as back, hip, or shoulder)

ORIGINAL PRESCRIPTION STRENGTH

Voltaren


ARTHRITIS PAIN

diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever

ANTI-INFLAMMATORY

For Daily Treatment of Arthritis Pain

Use Enclosed Dosing Card to Measure a Dose



Upper body dose 2.25 inches long (2 grams)
Lower body dose 4.5 inches long (4 grams)

How to Use

- For arthritis pain only
- Use 4 times a day
- May take up to 7 days to work for your arthritis pain
- For use on no more than 2 body areas
- Use up to 21 days unless directed by your doctor

Drug Facts

Active ingredient: Diclofenac sodium (NSAID) 1% (NSAID) – arthritis pain reliever

Purpose: Temporary relief of arthritis pain (OAT) in the following areas: hand, wrist, elbow (upper body areas); foot, ankle, knee (lower body areas)

Uses: For temporary relief of arthritis pain (OAT) in the following areas: hand, wrist, elbow (upper body areas); foot, ankle, knee (lower body areas)

Warnings: For external use only. Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Do not use if you have had an allergic reaction to any other pain reliever or to a fever reducer.

Directions: Use up to 4 times a day on no more than 2 body areas. For each body area, use the enclosed dosing card to measure a dose.

LOT EXP

1-855-297-3031

Made in Canada

Distributed by: **GSK Consumer Healthcare**, Warren, NJ 07069

Trademarks are owned by or licensed to the GSK group of companies.

©2022 GSK group of companies or its licensee.

PULL TAB



diclofenac sodium topical gel, 1% (NSAID) – arthritis pain reliever



3 TUBES of 150 g EACH, TOTAL NET WT 450 g (15.87 oz)



CARTON
RECYCLE

GSK Regulatory Spec Box		Verified Date: 10.20.22
Drug Facts Info		
Drug Facts	Helvetica Bold Condensed Italic	11 pt.
Drug Facts (continued)	Helvetica Bold Condensed Italic / Condensed	10 pt.
Headings	Helvetica Bold Condensed Italic	10 pt.
Subheadings	Helvetica Bold Condensed	7 pt.
Body text	Helvetica Condensed / Bold Condensed	7 pt.
Bullets	Helvetica Condensed	6 pt.
Bullets on same line, end of statement separated from bulleted statement by two ems		Yes
Spacing of the first line from edge of box ca. Minimum of 2 spaces after side of Drug Fact Box		Yes
Tracking	0	Horizontal Scale: 88 – 100%
Leading	0 pt.	Maximum Characters/Inch: < 30
Burline	2.5 pt.	Hairline: 0.5 pt.
Primary Display Panel Info		
Size of Net Wt/Contents (Smallest character height in inches)		> 0.1875 in.
PDP dimensions (in square inches)		25-100 sq. in.
Ratio of Statement or Identity to Logo (Largest Copy on PDP)		
Statement of Identity (pt.) (Divided by) Largest Logo Copy (pt.)		
Vertical PDP: 27%		Horizontal PDP: N/A

Voltaren

ARTHRITIS PAIN

diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever


User Guide

How to use Voltaren Arthritis Pain, with answers to frequently asked questions

Quick Start
How to get started right away with Voltaren Arthritis Pain

Understanding Voltaren Arthritis Pain Gel
Voltaren Arthritis Pain (diclofenac sodium topical gel, 1% (NSAID) - arthritis pain reliever) is an arthritis pain relief medication that you apply to your skin. It is intended to be used 4 times a day every day for up to 21 days.


Before You Use Voltaren Arthritis Pain
Read the Drug Facts Label, which appears on the carton.




Remove the dosing card from the inside of the carton. You should always use the dosing card to measure out the correct dose of Voltaren Arthritis Pain.

The first time you use a tube of Voltaren Arthritis Pain: You will have either one of the caps shown below.


Screw Cap
1. Take the cap off the tube. Open the safety seal by firmly pressing the indent on the top of the cap onto the star-shaped seal on the tube.
2. Firmly turn the cap to remove the safety seal. Do not open the safety seal with scissors or other sharp objects.
3. After use, put the cap back on the end of the tube.




Flip Cap
1. Flip the cap lid to open. Tamper evident plastic safety seals located on each side of the cap will break when the lid is opened for the first time. Before first use check if the tamper evident plastic safety seals are not broken or missing.
2. After use, close cap lid.



Using Voltaren Arthritis Pain
How often to use Voltaren Arthritis Pain
Use 4 times a day for best results.
For example:




Where to apply Voltaren Arthritis Pain
Voltaren Arthritis Pain can be used on up to 2 body areas from the following list: a hand, a wrist, an elbow, a foot, an ankle, a knee. Do not use on more than 2 body areas at once.
Some examples include:




NPA007817

Measuring the correct amount using the dosing card









For each upper body area (a hand, a wrist, or an elbow): Squeeze gel from the tube equal to the length shown on the upper body section of the dosing card (2.25 inches).




For each lower body area (a foot, an ankle, or a knee): Squeeze gel from the tube equal to the length shown on the lower body section of the dosing card (4.5 inches).

**Lost Your Dosing Card? Don't worry!
Call 1-855-297-3031 to get a free replacement.**


Applying Voltaren Arthritis Pain
Gently rub Voltaren Arthritis Pain into the skin using your hand. Apply 4 times a day for best results. Do not use on more than 2 body areas at once.
Apply only to clean, dry skin that doesn't have any cuts, open wounds, infections or rashes. Do not apply in the same area as any other medicines or products that are applied to the skin.

<p>Upper body areas</p>  <p>Elbow Including sides, inner, and outer</p>  <p>Wrist Including sides, top, and bottom</p>  <p>Hand Including top, palm, and fingers</p>	<p>Lower body areas</p>  <p>Knee Including sides, kneecap, and back</p>  <p>Ankle Including sides, front, and back</p>  <p>Foot Including sole, top of foot, and toes</p>
---	--


After You Use Voltaren Arthritis Pain



Wash your hands after applying Voltaren Arthritis Pain. If the treated area is the hands wait up to 1 hour to wash the hands.



Wash the dosing card with water. Store dosing card with your Voltaren Arthritis Pain and keep out of reach of children.



Wait ten minutes before covering the treated areas with clothes or gloves.

Avoid:

- showering or bathing for at least 1 hour after use
- exposing the treated area to sunlight or artificial light (such as in tanning booths)
- use in same area as other products applied to the skin
- applying a bandage over the treated area
- eye, nose, or mouth contact. If eye contact occurs, rinse thoroughly with water.

Facts you need to know
Important information about Voltaren Arthritis Pain

Do not use Voltaren Arthritis Pain...

- if you have ever had an allergic reaction to any other pain reliever or to a fever reducer
- for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.
- right before or after heart surgery
- on more than 2 body areas at the same time
- in the eyes, nose or mouth

Ask a doctor before use if...

- you have problems or serious side effects from taking pain relievers or fever reducers

Turn over for more information ➔

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.

Ask a doctor or pharmacist before use if...

- you are under a doctor's care for any serious condition
- you are taking any other drug
- you are pregnant or breast-feeding

It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Warnings to consider before use...
Allergy alert
Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- rash
- blisters
- facial swelling
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

Liver warning
This product contains diclofenac. Liver damage may occur if you apply

- more or for a longer time than directed
- when using other drugs containing diclofenac

Stomach bleeding warning
This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- apply more or for longer than directed

Heart attack and stroke warning
NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. This can be fatal. The risk is higher if you use more than directed or for longer than directed.

Follow these important guidelines when using Voltaren Arthritis Pain...

- use only as directed
- apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
- do not apply in same area as any other product used on the skin
- do not apply with external heat such as heating pad
- do not apply a bandage over the treated area
- do not get gel in eyes, nose or mouth
- store ENCLOSED DOSING CARD with your Voltaren Arthritis Pain. The dosing card is re-usable.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Stop using Voltaren Arthritis Pain and ask a doctor if...


- pain gets worse or lasts more than 21 days
- redness or swelling is present in the painful area
- fever occurs
- skin irritation occurs
- any new symptoms appear. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - leg swelling
 - weakness in one part or side of body
 - slurred speech

Questions and Answers
Answers to questions you may ask about Voltaren Arthritis Pain

Can I use Voltaren Arthritis Pain for the immediate relief of pain like that associated with sprains, strains, bruises or sports injuries?
No.
Do not use for sprains, sprains, bruises, or sports injuries. This product has not been shown to work for these types of injuries. It's intended only for the temporary relief of arthritis pain.

How quickly will Voltaren Arthritis Pain work for arthritis pain?
Voltaren Arthritis Pain may take up to 7 days to work for your arthritis pain; it is not for immediate relief. With 4 times-a-day use, you may start to feel relief within a few days. You should feel significant pain relief within 7 days of continuous use. If no pain relief in 7 days, stop use.

What is Voltaren Arthritis Pain used for?
Voltaren Arthritis Pain can be used for the temporary relief of arthritis pain in the hands, wrists, elbows, feet, ankles, and knees. Osteoarthritis (OA) is the most common form of arthritis. It mainly affects a type of tissue called cartilage, which cushions the joints and prevents the bones from rubbing against each other. With OA, cartilage begins to break down and wear away, resulting in joint pain and stiffness. OA occurs more frequently as you get older. Pain develops slowly and can gradually worsen over time.




How does Voltaren Arthritis Pain work?
Diclofenac sodium, the medicine in Voltaren Arthritis Pain, is a nonsteroidal anti-inflammatory drug (NSAID). NSAIDs are used to treat pain from medical conditions such as arthritis. Voltaren Arthritis Pain works similarly to oral NSAIDs like ibuprofen or naproxen by temporarily blocking the production of pain signaling chemicals called prostaglandins. However, when taking a pill, the medication is distributed to the site of pain through the bloodstream. Compared to oral diclofenac, only 6% of the medicine in Voltaren Arthritis Pain is absorbed in the bloodstream. It works by penetrating through your skin at the application site to deliver arthritis pain relief.

How does over-the-counter Voltaren Arthritis Pain differ from prescription Voltaren Gel?
Over-the-counter Voltaren Arthritis Pain is the same strength and formula as original full prescription strength Voltaren Gel but is available without a prescription.

Apply Voltaren Arthritis Pain 4 times a day to ensure you get the full effect of the medicine.

Can I apply Voltaren Arthritis Pain to upper and lower body areas at the same time?
Yes.
You can use Voltaren Arthritis Pain on up to 2 body areas at the same time. The areas can include the foot, ankle, knee, hand, wrist, and elbow.
Some correct examples include:



Do not use Voltaren Arthritis Pain on more than 2 body areas at once.

Can I apply Voltaren Arthritis Pain without measuring it out on the dosing card first?
No.
Use the dosing card to make sure you are getting the correct amount of medicine.

Can I use Voltaren Arthritis Pain on my spine, hips, or shoulders?
No.
The use of Voltaren Arthritis Pain on the spine, hips, and shoulders has not been studied.

Can I take any other pain medications with Voltaren Arthritis Pain?
Ask your doctor or pharmacist before using Voltaren Arthritis Pain if you are taking any other pain reliever or are planning on taking any other pain reliever, especially prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others).

Can I take any other pain medications during the first 7 days I use Voltaren Arthritis Pain?
Taking other pain medications is generally not recommended. Speak with your doctor or pharmacist before taking other pain medications when using Voltaren Arthritis Pain.

How long can I use Voltaren Arthritis Pain?
Use Voltaren Arthritis Pain 4 times a day every day for up to 21 days for treatment of arthritis pain or as directed by your doctor.
Voltaren Arthritis Pain should be part of an overall program for managing your arthritis pain. The program should also include activities such as appropriate movement or exercise, specific lifestyle changes, and weight maintenance. Regular visits and discussions with your doctor can help incorporate Voltaren Arthritis Pain into a complete program to help manage your arthritis pain.

What do I do if I get Voltaren Arthritis Pain in my eyes?
If eye contact occurs, rinse thoroughly with water and consult a doctor if irritation lasts more than an hour.

How do I clean the dosing card?
Wash the dosing card with water, and store dosing card and tube out of reach of children.

What if I miss a dose of Voltaren Arthritis Pain?
If you forget to apply Voltaren Arthritis Pain, don't double your dose. Just apply the next dose as scheduled.

How should I store Voltaren Arthritis Pain?

- Store at 20-25°C (68-77°F).
- Do not freeze Voltaren Arthritis Pain.
- Store the dosing card with your Voltaren Arthritis Pain.
- Keep out of reach of children.

Helpful Tip: If you forget to apply Voltaren Arthritis Pain, don't double your dose. Just apply the next dose as scheduled.

Additional Questions?
Call the Voltaren Arthritis Pain consumer hotline at 1-855-297-3031.
www.voltarengel.com

DOCUMENT FONT SIZES		
	TYPE STYLE	
BODY COPY	8 pt.	/ Diodrum Voltaren Family
IMAGE HEADERS	Min 14 pt.	/ Diodrum Voltaren Family & Helvetica Neue Condensed Bold
TITLE COPY	10 pt.	/ Diodrum Voltaren Family

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
11/23/2022 11:01:46 AM