

Drug Facts (Continued)

Warnings:

- Do not use if you are allergic to any of the ingredients listed on the label or to any of the following: aspirin, salicylic acid, salicylates, or other NSAIDs.
- Do not use if you are taking aspirin, salicylic acid, salicylates, or other NSAIDs.
- Do not use if you are taking blood thinners, such as warfarin, aspirin, or clopidogrel.
- Do not use if you are taking antiplatelet drugs, such as aspirin, clopidogrel, or prasugrel.
- Do not use if you are taking anticoagulants, such as warfarin, dabigatran, or rivaroxaban.
- Do not use if you are taking anti-infective drugs, such as amoxicillin, cefepime, or ceftriaxone.
- Do not use if you are taking anti-epileptic drugs, such as phenytoin, carbamazepine, or valproic acid.
- Do not use if you are taking anti-hypertensive drugs, such as lisinopril, amlodipine, or atenolol.
- Do not use if you are taking anti-diabetic drugs, such as insulin, metformin, or glimepiride.
- Do not use if you are taking anti-arrhythmic drugs, such as digoxin, amiodarone, or sotalolol.
- Do not use if you are taking anti-cancer drugs, such as tamoxifen, cyclophosphamide, or doxorubicin.
- Do not use if you are taking anti-viral drugs, such as zidovudine, zalcitabine, or didanosine.
- Do not use if you are taking anti-fungal drugs, such as itraconazole, voriconazole, or posaconazole.
- Do not use if you are taking anti-parasitic drugs, such as mebendazole, albendazole, or praziquantel.
- Do not use if you are taking anti-asthma drugs, such as albuterol, budesonide, or fluticasone.
- Do not use if you are taking anti-allergy drugs, such as cetirizine, loratadine, or fexofenadine.
- Do not use if you are taking anti-acid drugs, such as omeprazole, lansoprazole, or esomeprazole.
- Do not use if you are taking anti-nausea drugs, such as ondansetron, promethazine, or prochlorperazine.
- Do not use if you are taking anti-sickness drugs, such as dimenhydrinate, meclizine, or scopolamine.
- Do not use if you are taking anti-pain drugs, such as acetaminophen, ibuprofen, or naproxen.
- Do not use if you are taking anti-fever drugs, such as acetaminophen, ibuprofen, or naproxen.
- Do not use if you are taking anti-cough drugs, such as dextromethorphan, hydrocodone, or codeine.
- Do not use if you are taking anti-sleep drugs, such as zolpidem, zaleplon, or eszopiclone.
- Do not use if you are taking anti-anxiety drugs, such as alprazolam, clonazepam, or lorazepam.
- Do not use if you are taking anti-depression drugs, such as fluoxetine, sertraline, or escitalopram.
- Do not use if you are taking anti-psychotic drugs, such as risperidone, olanzapine, or aripiprazole.
- Do not use if you are taking anti-convulsant drugs, such as levetiracetam, lamotrigine, or gabapentin.
- Do not use if you are taking anti-seizure drugs, such as levetiracetam, lamotrigine, or gabapentin.
- Do not use if you are taking anti-epileptic drugs, such as levetiracetam, lamotrigine, or gabapentin.
- Do not use if you are taking anti-arrhythmic drugs, such as digoxin, amiodarone, or sotalolol.
- Do not use if you are taking anti-cancer drugs, such as tamoxifen, cyclophosphamide, or doxorubicin.
- Do not use if you are taking anti-viral drugs, such as zidovudine, zalcitabine, or didanosine.
- Do not use if you are taking anti-fungal drugs, such as itraconazole, voriconazole, or posaconazole.
- Do not use if you are taking anti-parasitic drugs, such as mebendazole, albendazole, or praziquantel.
- Do not use if you are taking anti-asthma drugs, such as albuterol, budesonide, or fluticasone.
- Do not use if you are taking anti-allergy drugs, such as cetirizine, loratadine, or fexofenadine.
- Do not use if you are taking anti-acid drugs, such as omeprazole, lansoprazole, or esomeprazole.
- Do not use if you are taking anti-nausea drugs, such as ondansetron, promethazine, or prochlorperazine.
- Do not use if you are taking anti-sickness drugs, such as dimenhydrinate, meclizine, or scopolamine.
- Do not use if you are taking anti-pain drugs, such as acetaminophen, ibuprofen, or naproxen.
- Do not use if you are taking anti-fever drugs, such as acetaminophen, ibuprofen, or naproxen.
- Do not use if you are taking anti-cough drugs, such as dextromethorphan, hydrocodone, or codeine.
- Do not use if you are taking anti-sleep drugs, such as zolpidem, zaleplon, or eszopiclone.
- Do not use if you are taking anti-anxiety drugs, such as alprazolam, clonazepam, or lorazepam.
- Do not use if you are taking anti-depression drugs, such as fluoxetine, sertraline, or escitalopram.
- Do not use if you are taking anti-psychotic drugs, such as risperidone, olanzapine, or aripiprazole.
- Do not use if you are taking anti-convulsant drugs, such as levetiracetam, lamotrigine, or gabapentin.
- Do not use if you are taking anti-seizure drugs, such as levetiracetam, lamotrigine, or gabapentin.
- Do not use if you are taking anti-epileptic drugs, such as levetiracetam, lamotrigine, or gabapentin.

Directions:

Apply a thin layer of gel to the affected area(s) 4 times a day. Do not use on more than 7 body areas at one time. Do not use on more than 21 days unless directed by your doctor.

Other information:

Keep out of reach of children. Store at room temperature (20° to 25°C).

Contains:

100 g (3.53 oz) of Voltaren Arthritis Pain gel.

Contains:

100 g (3.53 oz) of Voltaren Arthritis Pain gel.

Contains:

100 g (3.53 oz) of Voltaren Arthritis Pain gel.

Drug Facts (Continued)

Directions:

Apply a thin layer of gel to the affected area(s) 4 times a day. Do not use on more than 7 body areas at one time. Do not use on more than 21 days unless directed by your doctor.

Other information:

Keep out of reach of children. Store at room temperature (20° to 25°C).

Contains:

100 g (3.53 oz) of Voltaren Arthritis Pain gel.

Contains:

100 g (3.53 oz) of Voltaren Arthritis Pain gel.

Contains:

100 g (3.53 oz) of Voltaren Arthritis Pain gel.

Medicated Gel Clinically Proven to Relieve Arthritis Pain

TREATMENT AREAS

Hand Wrist Elbow Foot Ankle Knee

Not for use on any other body area (such as back, hip, or shoulder)

How to Use: ■ For arthritis pain apply 4 times a day ■ May take up to 7 days to work for your arthritis pain ■ For use on no more than 7 body areas at one time unless directed by your doctor

Use Enclosed Dosing Card to Measure a Dose

Voltaren NSAID **ARTHRITIS PAIN**

Dosing Card

Read product User Guide before use

Not for immediate pain relief

Upper body dose: 2.5 grams (0.09 oz)

Lower body dose: 1.25 grams (0.045 oz)

Temporarily avoid alcohol consumption while using this product.

ORIGINAL PRESCRIPTION STRENGTH

Voltaren **ARTHRITIS PAIN**

didolene sodium topical gel, 1% (NSAID) - arthritis pain reliever

ANTI-INFLAMMATORY

For Daily Treatment of Arthritis Pain

For external use only

NET WT 100 g (3.53 oz)

EXP. DATE

Drug Facts (Continued)

Directions:

Apply a thin layer of gel to the affected area(s) 4 times a day. Do not use on more than 7 body areas at one time. Do not use on more than 21 days unless directed by your doctor.

Other information:

Keep out of reach of children. Store at room temperature (20° to 25°C).

Contains:

100 g (3.53 oz) of Voltaren Arthritis Pain gel.

Contains:

100 g (3.53 oz) of Voltaren Arthritis Pain gel.

Contains:

100 g (3.53 oz) of Voltaren Arthritis Pain gel.

NPA012241

REGULATORY LEGEND	Actual Size
DRUG FACTS TITLE	9.38 pt
DRUG FACTS CONTINUED	8.34 pt
HEADINGS	8.34 pt
SUBHEADING/SBODY TEXT	6.25, 6.25 pt
LEADING	6.78 pt
BULLETS	5.25 pt
# OF CHARACTERS PER INCH	35
SPACE BEFORE BULLET	2 mm
BARLINES, HORIZONTAL	2.5, 0.5 pt
SPACE BETWEEN HORIZONTAL AND BOX END	1 space
STATEMENT OF IDENTITY (WHAT IT IS) Vertical	11 pt
STATEMENT OF IDENTITY (WHAT IT IS) Horizontal	13.13 pt

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
11/29/2023 10:17:32 AM