

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Feraheme safely and effectively. See full prescribing information for Feraheme.

FERAHEME® (ferumoxytol) Injection
For Intravenous (IV) use
Initial U.S. Approval: 2009

RECENT MAJOR CHANGES

Dosage and Administration (2)

12/2017

**WARNING: RISK FOR SERIOUS
HYPERSENSITIVITY/ANAPHYLAXIS REACTIONS**
See full prescribing information for complete boxed warning.

Fatal and serious hypersensitivity reactions including anaphylaxis have occurred in patients receiving Feraheme. Initial symptoms may include hypotension, syncope, unresponsiveness, cardiac/cardiorespiratory arrest.

- Only administer Feraheme when personnel and therapies are immediately available for the treatment of anaphylaxis and other hypersensitivity reactions. (5.1)
- Observe for signs or symptoms of hypersensitivity reactions during and for at least 30 minutes following Feraheme infusion including monitoring of blood pressure and pulse during and after Feraheme administration. (5.1)
- Hypersensitivity reactions have occurred in patients in whom a previous Feraheme dose was tolerated. (5.1)

INDICATIONS AND USAGE

Feraheme is an iron replacement product indicated for the treatment of iron deficiency anemia in adult patients with chronic kidney disease (CKD). (1)

DOSAGE AND ADMINISTRATION

- The recommended dose of Feraheme is an initial 510 mg dose followed by a second 510 mg dose 3 to 8 days later.

- Administer Feraheme as an intravenous infusion in 50-200 mL 0.9% Sodium Chloride Injection, USP or 5% Dextrose Injection, USP over at least 15 minutes

DOSAGE FORMS AND STRENGTHS

Injection: 510 mg iron per 17 mL (30 mg per mL) in single use vials. (3)

CONTRAINDICATIONS

- Known hypersensitivity to Feraheme or any of its components.
- History of allergic reaction to any intravenous iron product

WARNINGS AND PRECAUTIONS

- Greater risk of anaphylaxis in patients with multiple drug allergies. (5.1).
- Hypotension: Feraheme may cause hypotension. Monitor for signs and symptoms of hypotension following each administration of Feraheme. (5.2)
- Iron Overload: Regularly monitor hematologic responses during Feraheme therapy. Do not administer Feraheme to patients with iron overload. (5.3)
- Magnetic Resonance Imaging: Feraheme can alter magnetic resonance imaging (MRI) studies. (5.4)

ADVERSE REACTIONS

The most common adverse reactions ($\geq 2\%$) following the administration of Feraheme are diarrhea, nausea, dizziness, hypotension, constipation, and peripheral edema. (6.1)

To report SUSPECTED ADVERSE REACTIONS with Feraheme, contact AMAG Pharmaceuticals, Inc. at 1-877- 411-2510, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling

Revised: 12/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

1	INDICATIONS AND USAGE
2	DOSAGE AND ADMINISTRATION
3	DOSAGE FORMS AND STRENGTHS
4	CONTRAINDICATIONS
5	WARNINGS AND PRECAUTIONS
5.1	Serious Hypersensitivity Reactions
5.2	Hypotension
5.3	Iron overload
5.4	Magnetic Resonance (MR) Imaging
6	ADVERSE REACTIONS
6.1	Adverse Reactions in Clinical Studies
6.2	Postmarketing Experience
7	DRUG INTERACTIONS
8	USE IN SPECIFIC POPULATIONS
8.1	Pregnancy
8.3	Nursing Mothers
8.4	Pediatric Use
8.5	Geriatric Use

10	OVERDOSAGE
11	DESCRIPTION
12	CLINICAL PHARMACOLOGY
12.1	Mechanism of Action
12.2	Pharmacodynamics
12.3	Pharmacokinetics
13	NONCLINICAL TOXICOLOGY
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
14	CLINICAL STUDIES
16	HOW SUPPLIED/STORAGE AND HANDLING
16.1	How Supplied
16.2	Stability and Storage
17	PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: RISK FOR SERIOUS HYPERSENSITIVITY/ANAPHYLAXIS REACTIONS

Fatal and serious hypersensitivity reactions including anaphylaxis have occurred in patients receiving Feraheme. Initial symptoms may include hypotension, syncope, unresponsiveness, cardiac/cardiorespiratory arrest.

- Only administer Feraheme when personnel and therapies are immediately available for the treatment of anaphylaxis and other hypersensitivity reactions [see Warnings and Precautions (5.1)].
- Observe for signs or symptoms of hypersensitivity reactions during and for at least 30 minutes following Feraheme infusion including monitoring of blood pressure and pulse during and after Feraheme administration [see Warnings and Precautions (5.1)].
- Hypersensitivity reactions have occurred in patients in whom a previous Feraheme dose was tolerated [see Warnings and Precautions (5.1)].

1 INDICATIONS AND USAGE

Feraheme is indicated for the treatment of iron deficiency anemia in adult patients with chronic kidney disease (CKD).

2 DOSAGE AND ADMINISTRATION

The recommended dose of Feraheme is an initial 510 mg dose followed by a second 510 mg dose 3 to 8 days later. Administer Feraheme as an intravenous infusion in 50-200 mL 0.9% Sodium Chloride Injection, USP or 5% Dextrose Injection, USP over at least 15 minutes. Administer while the patient is in a reclined or semi-reclined position.

Feraheme does not contain antimicrobial preservatives. Feraheme, when added to intravenous infusion bags containing either 0.9% Sodium Chloride Injection, USP (normal saline), or 5% Dextrose Injection, USP, at concentrations of 2-8 mg elemental iron per mL, should be used immediately but may be stored at controlled room temperature ($25^{\circ}\text{C} \pm 2^{\circ}\text{C}$) for up to 4 hours or refrigerated ($2-8^{\circ}\text{C}$) for up to 48 hours.

The dosage is expressed in terms of mg of elemental iron, with each mL of Feraheme containing 30 mg of elemental iron. Evaluate the hematologic response (hemoglobin, ferritin, iron and transferrin saturation) at least one month following the second Feraheme infusion. The recommended Feraheme dose may be readministered to patients with persistent or recurrent iron deficiency anemia.

For patients receiving hemodialysis, administer Feraheme once the blood pressure is stable and the patient has completed at least one hour of hemodialysis. Monitor for signs and symptoms of hypotension following each Feraheme infusion.

Allow at least 30 minutes between administration of Feraheme and administration of other medications that could potentially cause serious hypersensitivity reactions and/or hypotension, such as chemotherapeutic agents or monoclonal antibodies.

Inspect parenteral drug products visually for the absence of particulate matter and discoloration prior to administration.

3 DOSAGE FORMS AND STRENGTHS

Feraheme Injection is available in single use vials. Each vial contains 510 mg of elemental iron in 17 mL (30 mg per mL).

4 CONTRAINDICATIONS

Feraheme is contraindicated in patients with:

- Known hypersensitivity to Feraheme or any of its components
- History of allergic reaction to any intravenous iron product

5 WARNINGS AND PRECAUTIONS

5.1 Serious Hypersensitivity Reactions

Fatal and serious hypersensitivity reactions including anaphylaxis, presenting with cardiac/ cardiorespiratory arrest, clinically significant hypotension, syncope, or unresponsiveness have occurred in patients receiving Feraheme [*see Boxed Warning*]. Other adverse reactions potentially associated with hypersensitivity have occurred (pruritus, rash, urticaria, and wheezing). These reactions have occurred following the first dose or subsequent doses in patients in whom a previous Feraheme dose was tolerated.

Patients with a history of multiple drug allergies may have a greater risk of anaphylaxis with parenteral iron products. Carefully consider the potential risks and benefits before administering Feraheme to these patients.

Only administer Feraheme when personnel and therapies are immediately available for the treatment of anaphylaxis and other hypersensitivity reactions. Closely observe patients for signs and symptoms of hypersensitivity including monitoring of blood pressure and pulse during and after Feraheme administration for at least 30 minutes and until clinically stable following completion of each infusion [*see Adverse Reactions (6.2)*].

In clinical studies predominantly in patients with CKD, serious hypersensitivity reactions were reported in 0.2% (3/1,726) of subjects receiving Feraheme. Other adverse reactions potentially associated with hypersensitivity (e.g., pruritus, rash, urticaria or wheezing) were reported in 3.7% (63/1,726) of these subjects. In other trials excluding patients with Stages 4 and 5 CKD, moderate to severe hypersensitivity reactions were reported in 2.6% (26/1014) of patients treated with Feraheme.

In the post-marketing experience, fatal and serious anaphylactic type reactions presenting with cardiac/ cardiorespiratory arrest, clinically significant hypotension, syncope, and unresponsiveness have been reported. Elderly patients with multiple or serious co-morbidities who experience hypersensitivity reactions and/or hypotension following administration of

Feraheme may have more severe outcomes [*see Boxed Warning, Adverse Reactions (6.2) and Use in Specific Populations (8.5)*].

5.2 Hypotension

Severe adverse reactions of clinically significant hypotension have been reported. In clinical studies, hypotension was reported in 1.9% (33/1,726) of subjects, including three patients with serious hypotensive reactions. Hypotension has also been reported in the post-marketing experience [*see Adverse Reactions from Post-marketing Spontaneous Reports (6.2)*]. Monitor patients for signs and symptoms of hypotension following each Feraheme administration [*see Dosage and Administration (2) and Warnings and Precautions (5.1)*].

5.3 Iron Overload

Excessive therapy with parenteral iron can lead to excess storage of iron with the possibility of iatrogenic hemosiderosis. Regularly monitor the hematologic response during parenteral iron therapy [*see Dosage and Administration (2)*]. Do not administer Feraheme to patients with iron overload.

In the 24 hours following administration of Feraheme, laboratory assays may overestimate serum iron and transferrin bound iron by also measuring the iron in the Feraheme complex.

5.4 Magnetic Resonance (MR) Imaging

Administration of Feraheme may transiently affect the diagnostic ability of MR imaging. Anticipated MR imaging studies should be conducted prior to the administration of Feraheme. Alteration of MR imaging studies may persist for up to 3 months following the last Feraheme dose. If MR imaging is required within 3 months after Feraheme administration, use T1- or proton density-weighted MR pulse sequences to minimize the Feraheme effects; MR imaging using T2-weighted pulse sequences should not be performed earlier than 4 weeks after the administration of Feraheme. Maximum alteration of vascular MR imaging is anticipated to be evident for 1 – 2 days following Feraheme administration [*see Clinical Pharmacology (12.3)*].

Feraheme will not interfere with X-ray, computed tomography (CT), positron emission tomography (PET), single photon emission computed tomography (SPECT), ultrasound or nuclear medicine imaging.

6 ADVERSE REACTIONS

Feraheme administration may cause serious hypersensitivity reactions and hypotension [*see Warnings and Precautions (5.1),(5.2)*].

In clinical studies, 1,726 subjects were exposed to Feraheme; 1,562 of these had CKD and 164 did not have CKD. Of these subjects 46% were male and the median age was 63 years (range of 18 to 96 years).

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug may not reflect the rates observed in practice.

6.1 Adverse Reactions in Clinical Studies

Across the three randomized clinical trials [Trial 1, 2, and 3, *see Clinical Studies (14)*], a total of 605 patients were exposed to two injections of 510 mg of Feraheme and a total of 280 patients were exposed to 200 mg/day of oral iron for 21 days. Most patients received their second Feraheme injection 3 to 8 days after the first injection.

Adverse reactions related to Feraheme and reported by $\geq 1\%$ of Feraheme-treated patients in the randomized clinical trials are listed in Table 1. Diarrhea (4.0%), constipation (2.1%) and hypertension (1.0%) have also been reported in Feraheme-treated patients.

Table 1: Adverse Reactions to Feraheme Reported in $\geq 1\%$ of Patients with CKD

Adverse Reactions	Feraheme 2 x 510 mg (n = 605)	Oral Iron (n = 280)
Nausea	3.1%	7.5%
Dizziness	2.6%	1.8%
Hypotension	2.5%	0.4%
Peripheral Edema	2.0%	3.2%
Headache	1.8%	2.1%
Edema	1.5%	1.4%
Vomiting	1.5%	5.0%
Abdominal Pain	1.3%	1.4%
Chest Pain	1.3%	0.7%
Cough	1.3%	1.4%
Pruritus	1.2%	0.4%
Pyrexia	1.0%	0.7%
Back Pain	1.0%	0%
Muscle Spasms	1.0%	1.4%
Dyspnea	1.0%	1.1%
Rash	1.0%	0.4%

In clinical trials, adverse reactions leading to treatment discontinuation and occurring in ≥ 2 Feraheme-treated patients included hypotension, infusion site swelling, increased serum ferritin level, chest pain, diarrhea, dizziness, ecchymosis, pruritus, chronic renal failure, and urticaria.

Following completion of the controlled phase of the trials, 69 patients received two additional 510 mg intravenous injections of Feraheme (for a total cumulative dose of 2.04 g). Adverse reactions following this repeat Feraheme dosing were similar in character and frequency to those observed following the first two intravenous injections.

In a placebo-controlled, cross-over trial, 713 patients with CKD received a single 510 mg dose of Feraheme. Adverse reactions reported by these patients were similar in character and frequency to those observed in other clinical trials.

6.2 Postmarketing Experience

Because adverse reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The following serious adverse reactions have been reported from the post-marketing experience with Feraheme: fatal, life-threatening, and serious anaphylactic-type reactions, cardiac/cardiorespiratory arrest, clinically significant hypotension, syncope, unresponsiveness, loss of consciousness, tachycardia/rhythm abnormalities, angioedema, ischemic myocardial events, congestive heart failure, pulse absent, and cyanosis. These adverse reactions have usually occurred within 30 minutes after the administration of Feraheme. Reactions have occurred following the first dose or subsequent doses of Feraheme.

7 DRUG INTERACTIONS

Drug-drug interaction studies with Feraheme were not conducted. Feraheme may reduce the absorption of concomitantly administered oral iron preparations.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C

There are no studies of Feraheme in pregnant women. In animal studies, ferumoxytol caused fetal malformations and decreased fetal weights at maternally toxic doses of 6 times the estimated human daily dose. Use Feraheme during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Administration of ferumoxytol during organogenesis, at doses of 31.6 mg Fe/kg/day in rats and 16.5 mg Fe/kg/day in rabbits, did not result in maternal or fetal effects. These doses are approximately 2 times the estimated human daily dose based on body surface area. In rats, administration of ferumoxytol during organogenesis at a maternally toxic dose of 100 mg Fe/kg/day, approximately 6 times the estimated human daily dose based on body surface area, caused a decrease in fetal weights. In rabbits, administration of ferumoxytol during organogenesis at a maternally toxic dose of 45 mg Fe/kg/day, approximately 6 times the estimated human daily dose based on body surface area, was associated with external and/or soft tissue fetal malformations and decreased fetal weights.

8.3 Nursing Mothers

It is not known whether Feraheme is present in human milk. Because many drugs are excreted in human milk and because of the potential for adverse reactions in nursing infants, a decision should be made whether to discontinue nursing or to avoid Feraheme, taking into account the importance of Feraheme to the mother and the known benefits of nursing.

8.4 Pediatric Use

The safety and effectiveness of Feraheme in pediatric patients (less than 18 years old) have not been established.

8.5 Geriatric Use

In controlled clinical trials, 330 patients \geq 65 years of age were treated with Feraheme. No overall differences in safety and efficacy were observed between older and younger patients in these trials, but greater sensitivity of older individuals cannot be ruled out. In general, dose administration to an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy. Elderly patients with multiple or serious co-morbidities who experience hypersensitivity reactions and/or hypotension following administration of Feraheme may have more severe outcomes. The potential risks and benefits of Feraheme administration should be carefully considered in these patients [*see Dosage and Administration (2) Serious Hypersensitivity Reactions (5.1) and Clinical Studies (14)*].

10 OVERDOSAGE

Limited data are available regarding overdosage of Feraheme in humans.

Excessive dosages of Feraheme may lead to accumulation of iron in storage sites potentially leading to hemosiderosis. Do not administer Feraheme to patients with iron overload [*Warnings and Precautions (5.3)*].

11 DESCRIPTION

Feraheme, an iron replacement product, is a non-stoichiometric magnetite (superparamagnetic iron oxide) coated with polyglucose sorbitol carboxymethylether. The overall colloidal particle size is 17-31 nm in diameter. The chemical formula of Feraheme is $\text{Fe}_{5874}\text{O}_{8752}\text{-C}_{11719}\text{H}_{18682}\text{O}_{9933}\text{Na}_{414}$ with an apparent molecular weight of 750 kDa.

Feraheme Injection is an aqueous colloidal product that is formulated with mannitol. It is a black to reddish brown liquid, and is provided in single use vials containing 510 mg of elemental iron. Each mL of the sterile colloidal solution of Feraheme Injection contains 30 mg of elemental iron and 44 mg of mannitol, and has low bleomycin-detectable iron. The formulation is isotonic with an osmolality of 270-330 mOsm/kg. The product contains no preservatives, and has a pH of 6 to 8.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Feraheme consists of a superparamagnetic iron oxide that is coated with a carbohydrate shell, which helps to isolate the bioactive iron from plasma components until the iron-carbohydrate complex enters the reticuloendothelial system macrophages of the liver, spleen and bone marrow. The iron is released from the iron-carbohydrate complex within vesicles in the macrophages. Iron then either enters the intracellular storage iron pool (e.g., ferritin) or is transferred to plasma transferrin for transport to erythroid precursor cells for incorporation into hemoglobin.

12.2 Pharmacodynamics

Cardiac Electrophysiology

In a randomized, positive- and placebo-controlled, parallel-group study, healthy subjects received a supratherapeutic regimen of Feraheme (1.02 g given as two 510 mg doses within 24 hours), placebo or a single dose of 400 mg moxifloxacin (positive control). Results demonstrated no effect of Feraheme on QT interval durations. No clinically meaningful effect of Feraheme on heart rate was observed.

12.3 Pharmacokinetics

The pharmacokinetic (PK) behavior of Feraheme has been examined in healthy subjects and in patients with CKD stage 5D on hemodialysis. Feraheme exhibited dose-dependent, capacity-limited elimination from plasma with a half life of approximately 15 hours in humans. The clearance (CL) was decreased by increasing the dose of Feraheme. Volume of distribution (Vd) was consistent with plasma volume, and the mean maximum observed plasma concentration (C_{max}) and terminal half-life ($t_{1/2}$) values increased with dose. The estimated values of CL and Vd following two 510 mg doses of Feraheme administered intravenously within 24 hours were 69.1 mL/hr and 3.16 L, respectively. The C_{max} and time of maximum concentration (t_{max}) were 206 mcg/mL and 0.32 hr, respectively. Rate of infusion had no influence on Feraheme PK parameters. No gender differences in Feraheme PK parameters were observed. Feraheme is not removed by hemodialysis.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Ferumoxytol was not tested for carcinogenic effects. In standard genotoxicity tests, ferumoxytol showed no evidence of mutagenic activity in an *in vitro* Ames test or clastogenic activity in either an *in vitro* chromosomal aberration assay or an *in vivo* micronucleus assay.

No adverse effects on fertility or general reproductive performance were noted in animal studies. Ferumoxytol had no effect on male or female fertility or general reproductive function in rats.

14 CLINICAL STUDIES

The safety and efficacy of Feraheme for the episodic treatment of iron deficiency anemia in patients with CKD were assessed in three randomized, open-label, controlled clinical trials (Trial 1, 2 and 3). These trials also included an uncontrolled, follow-up phase in which patients with persistent iron deficiency anemia could receive two additional 510 mg intravenous injections of Feraheme. The major efficacy results from the controlled phase of each study are shown in Table 2.

In all three trials, patients with CKD and iron deficiency anemia were randomized to treatment with Feraheme or oral iron. Feraheme was administered as two 510 mg intravenous single doses and oral iron (ferrous fumarate) was administered as a total daily dose of 200 mg elemental iron daily for 21 days. The major trial outcomes assessed the change in hemoglobin from baseline to Day 35. Trial 1 and 2 enrolled patients with non-dialysis dependent CKD and Trial 3 enrolled patients who were undergoing hemodialysis.

In Trial 1, the mean age of patients was 66 years (range, 23 to 95); 60% were female; 65% were Caucasian, 32% were Black, and 2% were other races. In the Feraheme and oral iron groups, 42% and 44% of patients, respectively, were receiving erythropoiesis stimulating agents (ESAs) at baseline.

In Trial 2, the mean age of patients was 65 years (range, 31 to 96); 61% were female; 58% were Caucasian, 35% were Black, and 7% were other races. In the Feraheme and oral iron groups, 36% and 43% of patients, respectively, were receiving ESAs at baseline.

In Trial 3, the mean age of patients was 60 years (range, 24 to 87); 43% were female; 34% were Caucasian, 59% were Black, and 7% were other races. All patients were receiving ESAs.

Table 2 shows the Baseline and mean change to Day 35 in hemoglobin (Hgb, g/dL), transferrin saturation (TSAT, %) and ferritin (ng/mL) in each treatment group for Trial 1, 2, and 3.

Table 2: Changes from Baseline to Day 35 in Hemoglobin, Transferrin Saturation and Ferritin (Intent to Treat Population)

ENDPOINT	Trial 1 Non-Dialysis CKD		Trial 2 Non-Dialysis CKD		Trial 3 CKD on Dialysis	
	Feraheme n = 226	Oral Iron n = 77	Feraheme n = 228	Oral Iron n = 76	Feraheme n = 114	Oral Iron n = 116
Baseline Hgb (mean ± SD, g/dL)	9.9 ± 0.8	9.9 ± 0.7	10.0 ± 0.7	10.0 ± 0.8	10.6 ± 0.7	10.7 ± 0.6
Hgb change from Baseline at Day 35 (mean ± SD, g/dL)	1.2* ± 1.3	0.5 ± 1.0	0.8* ± 1.2	0.2 ± 1.0	1.0* ± 1.1	0.5 ± 1.1
Baseline TSAT	9.8	10.4	11.3	10.1	15.7	15.9

(mean ± SD, %)	± 5.4	± 5.2	± 6.1	± 5.5	± 7.2	± 6.3
TSAT change from Baseline at Day 35 (mean ± SD, %)	9.2 ± 9.4	0.3 ± 4.7	9.8 ± 9.2	1.3 ± 6.4	6.4 ± 12.6	0.6 ± 8.3
Baseline ferritin (mean ± SD, ng/mL)	123.7 ± 125.4	146.2 ± 136.3	146.1 ± 173.6	143.5 ± 144.9	340.5 ± 159.1	357.6 ± 171.7
Ferritin change from Baseline at Day 35 (mean ± SD, ng/mL)	300.7 ± 214.9	0.3 ± 82.0	381.7 ± 278.6	6.9 ± 60.1	233.9 ± 207.0	-59.2 ± 106.2

* p≤0.001 for main efficacy endpoint

Following completion of the controlled phase of each of the Phase 3 trials, patients who were iron deficient and anemic could receive two additional 510 mg intravenous injections of Feraheme for a total cumulative dose of 2.04 g. Overall, 69 patients received two additional 510 mg intravenous injections of Feraheme, and on Day 35 following these additional injections, the majority of these patients (70%) experienced an increase in hemoglobin and iron parameters (TSAT and ferritin). The mean change (±SD) in hemoglobin level from the retreatment baseline for patients with an increase in hemoglobin was 0.86 (± 0.68) g/dL and was 0.5 (± 0.8) g/dL for all patients.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Feraheme is available in single use vials in the following package sizes (Table 3).

Table 3: Feraheme Packaging Description

NDC Code	Dose / Total volume per vial	Vials / Carton
NDC 59338-775-01	510 mg/ 17 mL	1
NDC 59338-775-10	510 mg/ 17 mL	10

16.2 Stability and Storage

Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° – 30°C (59° – 86°F) [see USP controlled room temperature].

17 PATIENT COUNSELING INFORMATION

Refer patients to the FDA approved Patient Package Insert.

Prior to Feraheme administration:

- Question patients regarding a history of allergy to intravenous iron or any medications.
- Advise patients of the serious risks associated with Feraheme.
- Advise patients to immediately report any signs and symptoms of hypersensitivity that may develop during and following Feraheme administration, such as rash, itching, dizziness, lightheadedness, swelling and breathing problems. Advise patients to seek immediate medical attention if these occur [*see Warnings and Precautions (5)*].

U.S Patents: 6,599,498 B1; 7,553,479 B2; 7,871,597 B2; 8,501,158 B2; 8,591,864 B2;
8,926,947 B2

Distributed by: AMAG Pharmaceuticals, Inc. Waltham, MA 02451